

# ON $L$ -PACKETS AND DEPTH FOR $\mathrm{SL}_2(K)$ AND ITS INNER FORM

ANNE-MARIE AUBERT, SERGIO MENDES, ROGER PLYMEN,  
AND MAARTEN SOLLEVELD

ABSTRACT. We consider the group  $\mathrm{SL}_2(K)$ , where  $K$  a local non-archimedean field of characteristic two. We prove that the depth of any irreducible representation of  $\mathrm{SL}_2(K)$  is larger than the depth of the corresponding Langlands parameter, with equality if and only if the  $L$ -parameter is essentially tame.

We also work out a classification of all  $L$ -packets for  $\mathrm{SL}_2(K)$  and for its non-split inner form, and we provide explicit formulae for the depths of their  $L$ -parameters.

## CONTENTS

1. Introduction	1
2. Depth of $L$ -parameters	4
3. $L$ -packets	8
3.1. Stability	9
3.2. $L$ -packets of cardinality one	10
3.3. Supercuspidal $L$ -packets of cardinality two	11
3.4. Supercuspidal $L$ -packets of cardinality four	12
3.5. Principal series $L$ -packets of cardinality two	13
Appendix A. Artin-Schreier symbol	15
A.1. Explicit formula for the Artin-Schreier symbol	16
A.2. Ramification	18
References	21

## 1. INTRODUCTION

Let  $K$  be a non-archimedean local field and let  $K_s$  be a separable closure of  $K$ . A central role in the representation theory of reductive  $K$ -groups is played by the local Langlands correspondence (LLC). It is known to exist in particular for the inner forms of the groups  $\mathrm{GL}_n(K)$  or  $\mathrm{SL}_n(K)$ , and to preserve interesting arithmetic information, like local  $L$ -functions and  $\epsilon$ -factors.

Another invariant that makes sense on both sides of the LLC is *depth*. The *depth*  $d(\pi)$  of an irreducible smooth representation  $\pi$  of a reductive  $p$ -adic group  $\mathcal{G}$  was defined by Moy and Prasad [MoPr] in terms of filtrations  $\mathcal{G}_{x,r}$  ( $r \in \mathbb{R}_{\geq 0}$ ) of its

---

*Date:* October 12, 2016.

*Key words and phrases.* Representation theory, local field,  $L$ -packets, depth.

Maarten Solleveld is supported by a NWO Vidi grant "A Hecke algebra approach to the local Langlands correspondence" (nr. 639.032.528).

parahoric subgroups  $\mathcal{G}_x$ . The depth of a Langlands parameter  $\phi$  is defined to be the smallest number  $d(\phi) \geq 0$  such that  $\phi$  is trivial on  $\text{Gal}(F_s/F)^r$  for all  $r > d(\phi)$ , where  $\text{Gal}(K_s/K)^r$  be the  $r$ -th ramification subgroup of the absolute Galois group of  $K$ .

Let  $D$  be a division algebra with centre  $K$ , of dimension  $d^2$  over  $K$ . Then  $\text{GL}_m(D)$  is an inner form of  $\text{GL}_n(K)$  with  $n = dm$ . There is a reduced norm map  $\text{Nrd}: \text{GL}_m(D) \rightarrow K^\times$  and the derived group  $\text{SL}_m(D) := \ker(\text{Nrd}: \mathcal{G} \rightarrow K^\times)$  is an inner form of  $\text{SL}_n(K)$ . Every inner form of  $\text{GL}_n(K)$  or  $\text{SL}_n(K)$  is isomorphic to one of this kind. When  $n = 2$ , the only possibilities for  $d$  are 1 or 2, and so the inner forms are, up to isomorphism,  $\text{GL}_2(K)$  and  $D^\times$ , and  $\text{SL}_2(K)$  and  $\text{SL}_1(D)$ .

The LLC for  $\text{GL}_m(D)$  preserves the depth, that is, for every smooth irreducible representation  $\pi$  of  $\text{GL}_m(D)$ , we have  $d(\pi) = d(\varphi_\pi)$ , where  $\varphi_\pi$  corresponds to  $\pi$  by the LLC [ABPS1, Theorem 2.9].

The situation is different for  $\text{SL}_m(D)$ . All the irreducible representations in a given L-packet  $\Pi_\phi$  have the same depth, so the depth is an invariant of the L-packet, say  $d(\Pi_\phi)$ . We have  $d(\Pi_\phi) = d(\varphi)$  where  $\varphi$  is a lift of  $\phi$  which has minimal depth among the lifts of  $\phi$ , and the following holds:

$$(1) \quad d(\phi) \leq d(\Pi_\phi)$$

for any Langlands parameter  $\phi$  for  $\text{SL}_m(D)$  [ABPS1, Proposition 3.4 and Corollary 3.4]. Moreover (1) is an equality if  $\phi$  is *essentially tame*, that is, if the image by  $\phi$  of the wild inertia subgroup  $\mathbf{P}_K$  of the Weil group  $\mathbf{W}_K$  of  $K$  lies in a maximal torus of  $\text{PGL}_n(\mathbb{C})$ .

We observe that this notion of essentially tameness is consistent with the usual notion for Langlands parameters for  $\text{GL}_n(K)$ . Indeed, any lift  $\varphi: \mathbf{W}_K \rightarrow \text{GL}_n(\mathbb{C})$  of  $\phi$ , is called essentially tame if its restriction to  $\mathbf{P}_K$  is a direct sum of characters. Clearly  $\varphi$  is essentially tame if and only if  $\varphi(\mathbf{P}_K)$  lies in a maximal torus of  $\text{GL}_n(\mathbb{C})$ , which in turn is equivalent to  $\phi(\mathbf{P}_K)$  lying in a maximal torus of  $\text{PGL}_n(\mathbb{C})$ .

We denote by  $t(\varphi)$  the *torsion number* of  $\varphi$ , that is, the number of unramified characters  $\chi$  of  $\mathbf{W}_K$  such  $\varphi\chi \cong \varphi$ . Then  $\phi$  and  $\varphi$  are essentially tame if and only if the residual characteristic  $p$  of  $K$  does not divide  $n/t(\varphi)$  [BuHe2, Appendix].

In this article we take  $K$  to be a local non-archimedean field  $K$  of characteristic 2. In positive characteristic,  $K$  is of the form  $K = \mathbb{F}_q((t))$ , the field of Laurent series with coefficients in  $\mathbb{F}_q$ , with  $q = 2^f$ . This case is particularly interesting because there are countably many quadratic extensions of  $\mathbb{F}_q((t))$ . These quadratic extensions are parametrised by the cosets in  $K/\wp(K)$  where  $\wp$  is the map, familiar from Artin-Schreier theory, given by  $\wp(X) = X^2 - X$ .

We first show that equality holds in (1) only if  $\phi$  is essentially tame (*i.e.*,  $t(\varphi) = 2$ ):

**Theorem 1.1.** *Let  $K$  be a non-archimedean local field of characteristic 2, and let  $\pi$  be an irreducible representation of an inner form of  $\text{SL}_2(K)$ , with Langlands parameter  $\phi$ . If  $\phi$  is not essentially tame then we have*

$$d(\pi) > d(\phi).$$

Let  $\varphi$  be a lift of  $\phi$  with minimal depth among the lifts of  $\phi$ . In the proof we distinguish the cases where  $\varphi$  is imprimitive, respectively primitive.

An irreducible Langlands parameter  $\varphi: \mathbf{W}_K \rightarrow \text{GL}_2(\mathbb{C})$  is called *imprimitive* if there exists a separable quadratic extension  $L$  of  $K$  and a character  $\xi$  of  $L^\times$  such

that  $\varphi \simeq \text{ind}_{\mathbf{W}_L^K}^{\mathbf{W}_K}(\xi)$ . Then the depth of  $\varphi$  and  $\phi$  may be expressed in terms of that of  $\xi$  and  $\xi^2$ , respectively, as

$$d(\varphi) = (d(\xi) + d(L/K))/2 \quad \text{and} \quad d(\phi) = (d(\xi^2) + d(L/K))/2,$$

where  $\mathfrak{p}_K^{1+d(L/K)}$  is the relative discriminant of  $L/K$ . Let  $\mathfrak{T}(\varphi)$  be the group of characters  $\chi$  of  $\mathbf{W}_K$  such that  $\chi \otimes \varphi \simeq \varphi$ . As in [BuHe1, 41.4], we call  $\varphi$  *totally ramified* if  $\mathfrak{T}(\phi)$  does not contain any unramified character. If  $\varphi$  is not essentially tame, then it is totally ramified. We check that if this case we have  $d(\xi) > d(\xi^2)$ , and hence  $d(\Pi_\phi) > d(\phi)$ .

We obtain in Proposition 3.2 the following characterization of  $L$ -packets for  $\text{SL}_2(K)$  or  $\text{SL}_1(D)$ : an  $L$ -packet is a minimal set of irreducible representations from which a stable distribution can be constructed.

Next we give the explicit classification of the  $L$ -packets for both  $\text{SL}_2(K)$  and  $\text{SL}_1(D)$ .

In particular, to each biquadratic extension  $L/K$ , there is attached a Langlands parameter  $\phi = \phi_{L/K}$ , and an  $L$ -packet  $\Pi_\phi$  of cardinality 4. The depth of the parameter  $\phi_{L/K}$  depends on the extension  $L/K$ . More precisely, the numbers  $d(\phi)$  depend on the breaks in the upper ramification filtration of the Galois group  $\text{Gal}(L/K) = \mathbb{Z}/2\mathbb{Z} \times \mathbb{Z}/2\mathbb{Z}$ . Let  $D$  be a central division algebra of dimension 4 over  $K$ . The parameter  $\phi$  is relevant for the inner form  $\text{SL}_1(D)$ , which admits singleton  $L$ -packets.

**Theorem 1.2.** *Let  $L/K$  be a biquadratic extension, let  $\phi$  be the Langlands parameter  $\phi_{L/K}$ . If the highest break in the upper ramification of the Galois group  $\text{Gal}(L/K)$  is  $t$  then we have  $d(\phi) = t$ . For every  $\pi \in \Pi_\phi(\text{SL}_2(K)) \cup \Pi_\phi(\text{SL}_1(D))$  these integers provide lower bounds:*

$$d(\pi) \geq d(\phi).$$

*Depending on the extension  $L/K$ , all the odd numbers 1, 3, 5, 7, ... are achieved as such breaks.*

This contrasts strikingly with the case of  $\text{SL}_2(\mathbb{Q}_p)$  with  $p > 2$ . Here there is a unique biquadratic extension  $L/K$ , and a unique tamely ramified discrete parameter  $\phi : \text{Gal}(L/K) \rightarrow \text{SO}_3(\mathbb{R})$  of depth zero.

Let  $E/K$  be the quadratic extension given by

$$E = K(\wp^{-1}(\varpi^{-2n-1}))$$

with  $\varpi$  a uniformizer and  $n = 0, 1, 2, 3, \dots$  and let  $\phi_E$  be the associated  $L$ -parameter. We prove in Subsection 3.4 that the depth of  $\phi_E$  is given by

$$d(\phi_E) = 2n + 1.$$

For the  $L$ -packets considered in this article, the depths  $d(\pi)$  can be arbitrarily large.

We have included an Appendix on aspects of the Artin-Schreier theory. This Appendix goes a little further than the exposition in [FeVo, p.146–151] and the article of Dalawat [Da]. We have the occasion to refer to the Appendix at several points in our article.

We thank Chandan Dalawat for a valuable exchange of emails and for bringing the reference [Da] to our attention.

2. DEPTH OF  $L$ -PARAMETERS

The field  $K$  possesses a central division algebra  $D$  of dimension 4 and, up to isomorphism, only one. The group  $D^\times$  is locally profinite and is compact modulo its centre  $K^\times$ , see [BuHe1, p.325]. Let  $\text{Nrd}$  denote the reduced norm on  $D^\times$ . Define

$$\text{SL}_1(D) = \{x \in D^\times : \text{Nrd}(x) = 1\}.$$

Then  $\text{SL}_1(D)$  is an inner form of  $\text{SL}_2(K)$ . The articles [HiSa, ABPS2] finalize the local Langlands correspondence for any inner form of  $\text{SL}_n$  over all local fields.

*Depth of an  $L$ -parameter for  $\text{GL}_2(K)$ .* Let  $\mathbf{W}_K$  denote the Weil group of  $K$ , and let  $\Phi(\text{GL}_2(K))$  be the set of  $L$ -parameters  $\varphi: \mathbf{W}_K \times \text{SL}_2(\mathbb{C}) \rightarrow \text{GL}_2(\mathbb{C})$  for inner forms of  $\text{GL}_2(K)$ . Let  $t$  be a real number,  $t \geq 0$ , let  $\text{Gal}(K_s/K)^t$  be the  $t$ -th ramification subgroup of the absolute Galois group of  $K$ . We define

$$(2) \quad \Phi_t(\text{GL}_2(K)) := \{\varphi \in \Phi(\text{GL}_2(K)) : \text{Gal}(K_s/K)^t \subset \ker(\varphi)\}.$$

Notice that  $\Phi_{t'}(\text{GL}_2(K)) \subset \Phi_t(\text{GL}_2(K))$ , if  $t' \leq t$ . It is known that the set of  $t$ 's at which  $\text{Gal}(F_s/F)^t$  breaks consists of rational numbers and is discrete [Ser, Chap. IV, §3]. In particular there exists a unique rational number  $d(\varphi)$ , called the *depth* of  $\varphi$ , such that

$$(3) \quad \varphi \notin \Phi_{d(\varphi)}(\text{GL}_2(K)) \quad \text{and} \quad \varphi \in \Phi_t(\text{GL}_2(K)) \quad \text{for any } t > d(\varphi).$$

*Depth of an  $L$ -parameter for  $\text{SL}_2(K)$ .* The depth of an  $L$ -parameter  $\phi: \mathbf{W}_K \times \text{SL}_2(\mathbb{C}) \rightarrow \text{PGL}_2(\mathbb{C})$  for an inner form of  $\text{SL}_2(K)$  is defined as:

$$(4) \quad d(\phi) = \inf\{t \in \mathbb{R}_{\geq 0} \mid \text{Gal}(K_s/K)^{t+} \subset \ker \phi\},$$

where

$$\text{Gal}(K_s/K)^{t+} := \bigcap_{r>t} G^r.$$

Each projective representation  $\phi: \mathbf{W}_K \rightarrow \text{PGL}_2(\mathbb{C})$  lifts to a Galois representation

$$\varphi: \mathbf{W}_K \rightarrow \text{GL}_2(\mathbb{C}).$$

For any such lift  $\varphi$  of  $\phi$  we have  $\ker(\varphi) \subset \ker \phi$ , so

$$(5) \quad d(\varphi) \geq d(\phi).$$

Let  $\varphi: \mathbf{W}_K \rightarrow \text{GL}_2(\mathbb{C})$  be a 2-dimensional irreducible representation of  $\mathbf{W}_K$ , and let  $\mathfrak{T}(\varphi)$  be the group of characters  $\chi$  of  $\mathbf{W}_K$  such that  $\chi \otimes \varphi \simeq \varphi$ . Then  $\varphi$  is primitive if  $\mathfrak{T}(\varphi) = \{1\}$ , *simply imprimitive* if  $\mathfrak{T}(\varphi)$  has order 2, and *triply imprimitive* if  $\mathfrak{T}(\varphi)$  has order 4, as in [BuHe1, 41.3]. Comparing determinants, we see that every nontrivial element of  $\mathfrak{T}(\varphi)$  has order 2.

As in [BuHe1, 41.4], we call  $\phi$  and  $\varphi$  *unramified* if  $\mathfrak{T}(\varphi) \setminus \{1\}$  contains an unramified character, and *totally ramified* if  $\mathfrak{T}(\varphi) \setminus \{1\}$  does not contain any unramified character. By definition, a primitive representation is totally ramified. Thus every imprimitive irreducible representation of dimension 2 of  $\mathbf{W}_K$  which is not totally ramified is essentially tame.

Let  $\phi: \mathbf{W}_K \times \text{SL}_2(\mathbb{C}) \rightarrow \text{PGL}_2(\mathbb{C})$  with trivial restriction to  $\text{SL}_2(\mathbb{C})$ , and such that  $\varphi$  is a lift of  $\phi$ . If  $\varphi$  is essentially tame and has minimal depth among the lifts of  $\phi$ , then we have  $d(\phi) = d(\varphi)$  [ABPS1, Theorem 3.8]. Thus we are reduced to computing the depths of the projective representations of  $\mathbf{W}_K$  which lift to totally ramified representations.

We recall how the depth of an irreducible representation  $(\varphi, V)$  of  $\mathbf{W}_K$  can be computed. Put  $E = (K_s)^{\ker \varphi}$ , so that  $\phi$  factors through  $\text{Gal}(E/K)$ . Let  $g_j$  be the order of the ramification subgroup  $\text{Gal}(E/K)_j$  (in the lower numbering). The Artin conductor  $a(\varphi) = a(V)$  is given by

$$(6) \quad a(\varphi) = g_0^{-1} \sum_{j \geq 0} g_j \dim(V/V^{\text{Gal}(E/K)_j}) \in \mathbb{Z}_{\geq 0}.$$

Since  $(\varphi, V)$  is irreducible and  $\text{Gal}(E/K)_j$  is normal in  $\text{Gal}(E/K)$ ,  $V^{\text{Gal}(E/K)_j} = 0$  whenever  $g_j > 1$ . Thus (6) simplifies to the formula [GrRe, (1)]:

$$(7) \quad a(\varphi) = \frac{\dim V}{g_0} \sum_{j \geq 0: g_j > 1} g_j = \dim V + \frac{\dim V}{g_0} \sum_{j \geq 1: g_j > 1} g_j$$

It was shown in [ABPS2, Lemma 4.1] that

$$(8) \quad d(\varphi) := \begin{cases} 0 & \text{if } \mathbf{I}_F \subset \ker(\phi), \\ \frac{a(\varphi)}{\dim V} - 1 & \text{otherwise.} \end{cases}$$

Let  $\varphi: \mathbf{W}_K \rightarrow \text{GL}_2(\mathbb{C})$  be a totally ramified irreducible representation. Let  $\phi: \mathbf{W}_K \rightarrow \text{PGL}_2(\mathbb{C})$  be its projection. We will show that  $d(\varphi) > d(\phi)$ . To this end we may and will assume that  $\varphi$  has minimal depth among the lifts of  $\phi$ .

**Theorem 2.1.** *Let  $\varphi$  be an irreducible totally ramified representation  $\mathbf{W}_K \rightarrow \text{GL}_2(\mathbb{C})$ , let  $\phi: \mathbf{W}_K \rightarrow \text{PGL}_2(\mathbb{C})$  be its projection. Then we have*

$$d(\varphi) > d(\phi).$$

*Proof. Primitive representations.* Let  $\varphi$  be primitive. Put  $E = K_s^{\ker \phi}$  and  $E^+ = K_s^{\ker \varphi}$ . By [BuHe1, §42.3] there exists a unique intermediate field  $K \subset L \subset E$  such that  $E/L$  is a wildly ramified biquadratic extension. Then  $\phi(\text{Gal}(E/L))$  is a subgroup of  $\text{PGL}_2(\mathbb{C})$  isomorphic to the Klein four group. Up to conjugacy  $\text{PGL}_2(\mathbb{C})$  has only one such subgroup. After a suitable change of basis, we may assume that it is

$$(9) \quad D_2 := \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & i \\ i & 0 \end{pmatrix}, \begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \right\} \subset \text{PGL}_2(\mathbb{C}).$$

The three subextensions of  $E/L$  are conjugate under  $\text{Gal}(E/K)$  because the conjugation action of  $A_4$  on its normal subgroup  $V_4$  of order four is transitive on the nontrivial elements of  $V_4$ . Hence there is a unique  $r \in \mathbb{Z}$  such that  $\text{Gal}(E/L)_r = \text{Gal}(E/L)$  and  $\text{Gal}(E/L)_{r+1} = \{1\}$ . In section A.2 we will see that  $r$  is odd. We call this  $r$  the *ramification depth* of  $E/L$ .

The nontrivial elements of  $\text{Gal}(E/L)$  are the deepest elements of  $\text{Gal}(E/K)$  outside the kernel of  $\phi$ , and therefore the depth of  $\phi$  can be expressed in terms of  $r$ .

Let us compare this to what happens for the lift  $\varphi$  of  $\phi$ . Since  $\text{SL}_2(\mathbb{C}) \rightarrow \text{PGL}_2(\mathbb{C})$  is a surjection with kernel of order 2, the preimage of  $\phi(\mathbf{W}_K)$  in  $\text{SL}_2(\mathbb{C})$  has order  $2|\phi(\mathbf{W}_K)|$ . The matrices in (9) do not yet form a group in  $\text{GL}_2(\mathbb{C})$ , for that we really need the nontrivial element of  $\ker(\text{SL}_2(\mathbb{C}) \rightarrow \text{PGL}_2(\mathbb{C}))$ . In other words,  $\text{SL}_2(\mathbb{C})$  contains a unique subgroup of order  $2[E : K]$  which projects onto  $\phi(\mathbf{W}_K)$ . As  $\varphi$  has minimal depth among the lifts of  $\phi$ ,  $\varphi(\mathbf{W}_K)$  is precisely this subgroup. Thus  $[E^+ : E] = 2$  and  $\text{Gal}(E^+/K)$  is a nontrivial index two central extension of  $\text{Gal}(E/K)$ . In particular  $\text{Gal}(E^+/L)$  is isomorphic to the quaternion group of order eight.

Choose a subset  $\{w_1 = 1, w_2, w_3, w_4\} \subset \text{Gal}(E^+/L)$  which projects onto  $\text{Gal}(E/L)$ . We may assume that the  $\varphi(w_i)$  are ordered as in (9). As  $\ker(\text{GL}_2(\mathbb{C}) \rightarrow \text{PGL}_2(\mathbb{C}))$  is central,

$$[\varphi(w_3), \varphi(w_4)] = \left[ \begin{pmatrix} -i & 0 \\ 0 & i \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \right] = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \in \text{GL}_2(\mathbb{C}).$$

Write

$$(10) \quad z = [w_3, w_4] \in \text{Gal}(E^+/L),$$

so that  $\varphi(z) = \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix}$ . It follows from the definition of  $r$  and the condition on  $\varphi$  that

$$\text{Gal}(E^+/L)_r = \text{Gal}(E^+/L) \text{ and } \text{Gal}(E^+/L)_{r+1} = \text{Gal}(E^+/E).$$

By [Ser, Proposition IV.2.10]  $z \in \text{Gal}(E^+/L)_{2r+1}$ . Now  $z \notin \ker(\varphi)$  and it lies deeper in  $\text{Gal}(E^+/K)$  than  $w_2, w_3$  and  $w_4$ . On the other hand,  $z$  does lie in the kernel of  $\phi$ , which explains why  $\varphi$  has larger depth than  $\phi$ .

In the sequel of this section, we assume that the depth of the element  $z$  defined in (10) is exactly  $2r+1$ . This is allowed because, in the above setting, it constitutes the worst possible case for the theorem.

*Octahedral representations.* Let  $\varphi$  be octahedral, that is, it is primitive and  $\phi(\mathbf{W}_K) \cong S_4$ . Let  $\text{Ad}$  denote the adjoint representation of  $\text{PGL}_2(\mathbb{C})$  on  $\mathfrak{sl}_2(\mathbb{C}) = \text{Lie}(\text{PGL}_2(\mathbb{C}))$ . Then  $\text{Ad} \circ \phi$  is an irreducible 3-dimensional representation of  $\mathbf{W}_K$ . Since  $\text{PGL}_2(\mathbb{C})$  is the adjoint group of  $\mathfrak{sl}_2(\mathbb{C})$ ,  $\text{Ad} \circ \phi$  has the same kernel and hence the same depth as  $\phi$ .

By [BuHe1, Theorem 42.2]  $L/K$  is Galois with automorphism group  $S_3$  and residue degree 2. Thus  $\text{Ad}(\phi(\mathbf{I}_K)) \subset \text{Ad}(\phi(\mathbf{W}_K))$  is a normal subgroup of index two, isomorphic to  $A_4$ . As  $L/K$  has tame ramification index 3, the image of the wild inertia subgroup  $\mathbf{P}_K$  under  $\text{Ad} \circ \phi$  equals the image of  $\text{Gal}(E/L)$ . By our convention (9) it is  $\text{Ad}(D_2)$ . By the definition of  $r$  as the ramification depth of  $E/L$ , we have

$$g_0 = 12, g_1 = \cdots = g_r = 4 \text{ and } g_{r+1} = 1$$

With the formula (7) we find

$$a(\text{Ad} \circ \phi) = \frac{3}{12}(12 + r \cdot 4) = 3 + r,$$

and from (8) we conclude that

$$d(\phi) = d(\text{Ad} \circ \phi) = r/3.$$

On the other hand,  $\varphi$  is an irreducible two-dimensional representation of  $\mathbf{W}_K$ , and we must base our calculations on the Galois group of  $E^+/K$ . The numbers

$$g_j = |\text{Gal}(E^+/K)_j| = |\varphi(\text{Gal}(E^+/K)_j)|$$

can be computed from those for  $\phi$  by means of the twofold covering  $\varphi(\mathbf{W}_K) \rightarrow \phi(\mathbf{W}_K)$ . We find

$$g_0 = 24, g_1 = \cdots = g_r = 8 \text{ and } g_{r+1} = \cdots = g_{2r+1} = 2.$$

Assuming that the depth of  $z$  is precisely  $2r+1$  (see above), we can also say that  $g_{2r+2} = 1$ . Then (7) gives

$$a(\varphi) = \frac{2}{24}(24 + r \cdot 8 + (r+1) \cdot 2) = 2 + \frac{5r+1}{6}.$$

Now (8) says that

$$d(\varphi) = (5r + 1)/12.$$

We note that this is strictly larger than  $d(\phi) = r/3$ . (As  $a(\phi) \in \mathbb{Z}_{\geq 0}$ , we must have  $r - 1 \in 6\mathbb{Z}$ . This means that above not all biquadratic extensions can occur.)

*Tetrahedral representations.* Let  $\varphi$  be tetrahedral, that is, it is primitive and  $\phi(\mathbf{W}_K) \cong A_4$ . By [BuHe1, Theorem 42.2]  $L/K$  is a cubic Galois extension. It is of prime order, so either it is unramified or it is totally ramified.

First we consider the case that  $L/K$  ramifies totally. Then  $\mathbf{I}_K$  surjects onto  $\text{Gal}(E/K)$ , so  $\varphi(\mathbf{I}_K) = \varphi(\mathbf{W}_K)$ . This means that within  $\mathbf{I}_K$  everything is similar to octahedral representations. The same calculations as above show that

$$d(\phi) = r/3 < d(\varphi) = (5r + 1)/12.$$

Now we look at the case where  $L/K$  is unramified. Then

$$\phi(\mathbf{I}_K) = \phi(\text{Gal}(E/K)) = D_2.$$

To compute the depth, we replace  $\phi$  by the 3-dimensional representation  $\text{Ad} \circ \phi$  of  $\mathbf{W}_K$  on  $\mathfrak{sl}_2(\mathbb{C})$ . With  $r$  as before,  $g_0 = \cdots = g_r = 4$  and  $g_{r+1} = 1$ . With (7) and (8) we calculate

$$\begin{aligned} a(\text{Ad} \circ \phi) &= \frac{3}{4}((r+1) \cdot 4) = 3(r+1), \\ d(\phi) &= d(\text{Ad} \circ \phi) = \frac{3(r+1)}{3} - 1 = r. \end{aligned}$$

Like in the octahedral case, the numbers  $\text{Gal}(E^+/K)_j$  for  $\varphi$  are related to those for  $\phi$  via the twofold covering  $\text{SL}_2(\mathbb{C}) \rightarrow \text{PGL}_2(\mathbb{C})$ . We find

$$g_0 = \cdots = g_r = 8 \text{ and } g_{r+1} = \cdots = g_{2r+1} = 2.$$

Moreover  $g_{2r+2} = 1$  if we assume that the depth of  $z$  is  $2r + 1$ . Now (7) says

$$a(\varphi) = \frac{2}{8}((r+1) \cdot 8 + (r+1) \cdot 2) = 5(r+1)/2 \in \mathbb{Z},$$

and from (8) we obtain

$$d(\varphi) = \frac{5(r+1)}{2 \cdot 2} - 1 = \frac{5r+1}{4}.$$

Again, this is larger than  $d(\phi) = r$ .

*Imprimitive representations.* Consider an imprimitive totally ramified representation  $\varphi : \mathbf{W}_K \rightarrow \text{GL}_2(\mathbb{C})$ . By [BuHe1, §41.4] there exists a separable totally ramified quadratic extension  $L/K$  and a character  $\xi$  of  $\mathbf{W}_L$  such that  $\varphi = \text{ind}_{\mathbf{W}_L}^{\mathbf{W}_K}(\xi)$ . Let  $\mathfrak{p}_K^{1+d(L/K)}$  be the discriminant of  $L/K$ . If  $L \cong K[X]/(X^2 + X + b)$ , then one deduces from [BuHe1, §41.1] that  $d(L/K) = -\nu_K(b) > 0$ .

From the proof of [BuHe1, Lemma 41.5] one sees that the level of  $\varphi$  equals  $d(\xi) + d(E/F)$ . By construction the level of a  $n$ -dimensional irreducible representation of  $\mathbf{W}_K$  equals  $n$  times its depth, so

$$(11) \quad d(\varphi) = (d(\xi) + d(L/K))/2.$$

As before we assume that  $\varphi$  is minimal among the lifts of  $\phi$ . Then [BuHe1, §41.4] says that  $d(\xi) > d(L/K)$ , and in particular  $d(\xi) \geq 2$ . Since  $\text{Gal}(K_s/L)^2$  is a pro-2-group, the image of  $\xi$  in  $\mathbb{C}^\times$  is a subgroup of even order.

Let  $\sigma$  be the nontrivial element of  $\text{Gal}(L/K)$ , so that the restriction of  $\varphi$  to  $\mathbf{W}_L$  is  $\xi \oplus \sigma(\xi)$ . If  $\xi(w) = -1$ , then also  $\xi(\sigma(w)) = -1$ . As  $\xi(\mathbf{W}_L)$  is even, this means that  $\begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \in \phi(\mathbf{W}_L)$ . We note that, as every  $\mathbf{W}_K \setminus \mathbf{W}_L$  interchanges  $\xi$  and  $\sigma(\xi)$ , the kernel of  $\phi$  equals the kernel of  $\xi \oplus \sigma(\xi)$  composed with the projection  $\text{GL}_2(\mathbb{C}) \rightarrow \text{PGL}_2(\mathbb{C})$ . Thus the kernel of  $\phi$  contains the kernel of  $\varphi$  with index two. More precisely

$$\ker(\phi) = (\xi \oplus \sigma(\xi))^{-1} \left\{ \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} -1 & 0 \\ 0 & -1 \end{pmatrix} \right\} = \xi^{-1} \{1, -1\} = \ker(\xi^2).$$

By the same argument as above also  $\ker(\text{ind}_{\mathbf{W}_L}^{\mathbf{W}_K} \xi^2) = \ker(\xi^2)$ . Hence  $\phi$  and  $\text{ind}_{\mathbf{W}_L}^{\mathbf{W}_K}(\xi^2)$  have the same kernel, and in particular the same depth. With (11) we can express it as

$$(12) \quad d(\phi) = (d(\xi^2) + d(L/K))/2.$$

The depth (or level) of  $\xi$  is the least  $l$  such that  $\xi$  (or rather its composition with the Artin reciprocity isomorphism) is nontrivial on the higher units group  $U_L^l = 1 + \mathfrak{p}_L^l \subset L^\times$ . For  $l > 0$  the group  $U_L^l/U_L^{l+1}$  has exponent 2, so  $\xi(U_L^{d(\xi)}) = \{1, -1\}$ . Consequently  $U_L^{d(\xi)} \subset \ker \xi^2$  and  $d(\xi^2) < d(\xi)$ . Comparing (11) and (12), we get

$$d(\varphi) - d(\phi) = (d(\xi) - d(\xi^2))/2 > 0. \quad \square$$

### 3. $L$ -PACKETS

According to a classical result of Shelstad [She, p.200], for  $F$  of characteristic zero all the  $L$ -packets  $\Pi_\varphi(\text{SL}_2(F))$  have cardinality 1, 2 or 4. We will check below, after (15), that the same holds for the  $L$ -packets for  $\text{SL}_2(K)$ . It will follow from the classification in this section that  $L$ -packets for  $\text{SL}_1(D)$  have cardinality 1 or 2.

**Theorem 3.1.** [ABPS1] *Let  $\phi : \mathbf{W}_K \times \text{SL}_2(\mathbb{C}) \rightarrow \text{PGL}_2(\mathbb{C})$  be an  $L$ -parameter for  $\text{SL}_2(K)$ , and let  $\varphi : \mathbf{W}_K \times \text{SL}_2(\mathbb{C}) \rightarrow \text{GL}_2(\mathbb{C})$  be a lift of minimal depth. For any  $\pi$  in one of the  $L$ -packets  $\Pi_\varphi(\text{GL}_2(K))$ ,  $\Pi_\varphi(\text{GL}_1(D))$ ,  $\Pi_\phi(\text{SL}_2(K))$  and  $\Pi_\phi(\text{SL}_1(D))$ :*

$$d(\phi) \leq d(\varphi) = d(\pi).$$

Moreover  $d(\phi) = d(\varphi) = d(\pi)$  if  $\varphi$  is essentially tame, in particular whenever  $\varphi$  is unramified.

We define the groups

$$(13) \quad \begin{aligned} C(\phi) &:= Z_{\text{SL}_2(\mathbb{C})}(\text{im } \phi), \\ \mathcal{S}_\phi &:= C(\phi)/C(\phi)^\circ = \pi_0(Z_{\text{SL}_2(\mathbb{C})}(\phi)), \\ \mathcal{Z}_\phi &:= Z(\text{SL}_2(\mathbb{C}))/Z(\text{SL}_2(\mathbb{C})) \cap C(\phi)^\circ, \\ \mathcal{S}_\phi &:= \pi_0(Z_{\text{PGL}_2(\mathbb{C})}(\phi)). \end{aligned}$$

The group  $\mathcal{S}_\phi$  is abelian,  $\mathcal{S}_\phi$  can be nonabelian, and there is a short exact sequence

$$(14) \quad 1 \rightarrow \mathcal{Z}_\phi \rightarrow \pi_0(Z_{\text{SL}_2(\mathbb{C})}(\phi)) \rightarrow \pi_0(Z_{\text{PGL}_2(\mathbb{C})}(\phi)) \rightarrow 1.$$



It is easily seen that  $|\mathcal{Z}_\phi| = 2$  if and only if  $\phi$  is relevant for  $\mathrm{SL}_1(D)$ . By [ABPS2, Theorem 3.3] there are bijections

$$(15) \quad \begin{aligned} \mathbf{Irr}(\pi_0(Z_{\mathrm{PGL}_2(\mathbb{C})}(\phi))) &\longleftrightarrow \Pi_\phi(\mathrm{SL}_2(K)), \\ \mathbf{Irr}(\pi_0(Z_{\mathrm{SL}_2(\mathbb{C})}(\phi))) &\longleftrightarrow \Pi_\phi(\mathrm{SL}_2(K)) \cup \Pi_\phi(\mathrm{SL}_1(D)). \end{aligned}$$

We remark for  $\mathrm{SL}_2(F)$  with  $\mathrm{char}(F) = 0$ , (15) was shown in [GeKn, Theorem 4.2] and [HiSa, Theorem 12.7]. Recall that  $\mathfrak{T}(\varphi)$  is the abelian group of characters  $\chi$  of  $\mathbf{W}_K$  with  $\varphi \otimes \chi \cong \varphi$ . By [GeKn, Theorem 4.3] and by [ABPS2, (21)]

$$(16) \quad \mathfrak{T}(\varphi) \cong \pi_0(Z_{\mathrm{PGL}_2(\mathbb{C})}(\phi)).$$

By [BuHe1, Proposition 41.3], and by the classification of L-parameters for the principal series in Subsection 3.2,  $\mathfrak{T}(\varphi)$  has order dividing four. This shows that all L-packets for  $\mathrm{SL}_2(K)$  have order 1, 2 or 4.

### 3.1. Stability.

Before we proceed with the classification of  $L$ -packets, some remarks about the stability of the associated distributions are in order. In this subsection  $K$  can be any local non-archimedean field. Recall that a class function on an algebraic  $K$ -group  $\mathcal{G}(K)$  is called stable if it is constant on the intersection of any  $\mathcal{G}(K_s)$ -conjugacy class with  $\mathcal{G}(K)$ . For an invariant distribution on  $\mathcal{G}(K)$  one would like to use a similar definition of stability, but that does not work well in general. Instead, stable distributions are usually defined in terms of stable orbital integrals. But, whenever an invariant distribution  $\delta$  on  $\mathcal{G}(K)$  is represented by a class function on an open dense subset of  $\mathcal{G}(K)$ , we can use the easier criterion for stability of functions to determine whether or not  $\delta$  is stable.

Harish-Chandra proved that the trace of an admissible representation is a distribution which is represented by a locally constant function on the set of regular semisimple elements of  $\mathcal{G}(K)$ , see [DBHCS]. So the study the stability of traces of  $\mathcal{G}(K)$ -representations, it suffices to look at (regular) semisimple elements of  $\mathcal{G}(K)$ .

For semisimple elements in  $\mathrm{GL}_2(K)$  conjugacy is the same as stable conjugacy, it is determined by characteristic polynomials. Hence every irreducible (admissible) representation of  $\mathrm{GL}_2(K)$  defines a stable distribution.

The semisimple conjugacy classes in  $\mathrm{GL}_1(D)$  are naturally in bijection with the elliptic conjugacy classes in  $\mathrm{GL}_2(K)$ , i.e. those semisimple classes for which the characteristic polynomials are irreducible over  $K$ . Moreover any irreducible essentially square-integrable representation of  $\mathrm{GL}_2(K)$  is already determined by the values of its trace on elliptic elements. These observations constitute some of the foundations of the Jacquet–Langlands correspondence [JaLa]. In fact the Jacquet–Langlands correspondence can be defined as the unique bijection between  $\mathbf{Irr}(\mathrm{GL}_1(D))$  and the essentially square-integrable representations in  $\mathbf{Irr}(\mathrm{GL}_2(K))$  which preserves the traces on elliptic conjugacy classes, up to a sign. Consequently the trace of any irreducible representation  $\pi$  of  $\mathrm{GL}_1(D)$  is the restriction of a stable distribution on  $\mathrm{GL}_2(K)$  to the set of elliptic elements. In particular the trace of  $\pi$  is itself a stable distribution.

**Proposition 3.2.** *Let  $\phi$  be a  $L$ -parameter for  $\mathrm{SL}_2(K)$ .*

- (a) *Write  $\Pi_\phi(\mathrm{SL}_2(K)) = \{\pi_1, \dots, \pi_m\}$ . The trace of  $\pi := \pi_1 \oplus \dots \oplus \pi_m$  is a stable distribution on  $\mathrm{SL}_2(K)$ . Any other stable distribution that can be obtained from  $\Pi_\phi(\mathrm{SL}_2(K))$  is a scalar multiple of the trace of  $\pi$ .*

- (b) *Suppose that  $\phi$  is relevant for  $\mathrm{SL}_1(D)$  and write  $\Pi_\phi(\mathrm{SL}_1(D)) = \{\pi'_1, \dots, \pi'_{m'}\}$ . The trace of  $\pi' := \pi'_1 \oplus \dots \oplus \pi'_{m'}$  is a stable distribution on  $\mathrm{SL}_1(D)$ . Any other stable distribution that can be obtained from  $\Pi_\phi(\mathrm{SL}_1(D))$  is a scalar multiple of the trace of  $\pi'$ .*

*Proof.* (a) Since the restriction of irreducible representations from  $\mathrm{GL}_2(K)$  to  $\mathrm{SL}_2(K)$  is multiplicity-free [BuKu, §1],  $\pi = \pi_1 \oplus \dots \oplus \pi_m$  is the restriction of some irreducible representation of  $\mathrm{GL}_2(K)$ . If  $\varphi : \mathbf{W}_K \times \mathrm{SL}_2(\mathbb{C}) \rightarrow \mathrm{GL}_2(\mathbb{C})$  is any lift of  $\phi$ , the image of  $\phi$  under the local Langlands correspondence is such a representation. We denote this representation of  $\mathrm{GL}_2(K)$  again by  $\pi$ . By the above remarks, its trace is a stable distribution on  $\mathrm{GL}_2(K)$ , and hence also on  $\mathrm{SL}_2(K)$ .

The different  $\pi_i$  are inequivalent, but they are  $\mathrm{GL}_2(K)$  conjugate, because  $\pi$  is irreducible. If a linear combination  $\sum_{i=1}^m \lambda_i \mathrm{tr}(\pi_i)$  is a stable distribution, then it must be invariant under conjugation by  $\mathrm{GL}_2(K)$ . Hence all the  $\lambda_i \in \mathbb{C}$  must be equal.

(b) The restriction of representations from  $\mathrm{GL}_1(D)$  to  $\mathrm{SL}_1(D)$  can have multiplicities, but still every constituent will appear with the same multiplicity [GeKn, Lemma 2.1.d]. So there exists an integer  $\mu$  such that  $\mu\pi' = \mu\pi'_1 \oplus \dots \oplus \mu\pi'_{m'}$  lifts to an irreducible representation of  $\mathrm{GL}_1(D)$ . The L-parameter of such a representation is a lift of  $\phi$ , so we can take  $\mathrm{JL}(\pi)$ , the image of  $\pi$  under the Jacquet–Langlands correspondence.

As remarked above,  $\mathrm{tr}(\mathrm{JL}(\pi))$  is stable distribution on  $\mathrm{GL}_1(D)$  and by restriction also on  $\mathrm{SL}_1(D)$ . Thus  $\mathrm{tr}(\pi') = \mu^{-1} \mathrm{tr}(\mathrm{JL}(\pi))$  is also a stable distribution on  $\mathrm{SL}_1(D)$ . By the same argument as for part (a), any linear combination of the  $\mathrm{tr}(\pi'_i)$  which is stable, must be a scalar multiple of  $\mathrm{tr}(\pi')$ .  $\square$

We remark that Proposition 3.2 also holds for inner forms of  $\mathrm{SL}_n(F)$  with  $n > 2$ . The proof is the same, one only has to replace the elliptic conjugacy classes by the conjugacy classes that correspond to elements of that particular inner form.

### 3.2. L-packets of cardinality one.

First we consider the case that  $\varphi : \mathbf{W}_K \rightarrow \mathrm{GL}_2(\mathbb{C})$  is irreducible, so the L-packet consists of supercuspidal representations. By (16) and (15),  $\Pi_\phi(\mathrm{SL}_2(K))$  is a singleton if and only if  $\varphi$  is primitive. The L-parameter  $\phi$  is relevant for  $\mathrm{SL}_1(D)$ , so  $\Pi_\phi(\mathrm{SL}_1(D))$  is nonempty. It follows from (15) and (14) that  $\mathcal{Z}_\phi \cong \pi_0(Z_{\mathrm{SL}_2(\mathbb{C})}(\phi)) \cong \mathbb{Z}/2\mathbb{Z}$ , and then from (15) that  $\Pi_\phi(\mathrm{SL}_1(D))$  is also a singleton. Any primitive representation of  $\mathbf{W}_K$  is either octahedral or tetrahedral, as in Section 2. See [BuHe1, §42] for more background.

Suppose now that  $\varphi : \mathbf{W}_K \rightarrow \mathrm{GL}_2(\mathbb{C})$  is reducible, so  $\phi$  is a L-parameter for the principal series of  $\mathrm{SL}_2(K)$ . If  $\phi(\mathbf{W}_K) = 1$  and  $\phi|_{\mathrm{SL}_2(\mathbb{C})} : \mathrm{SL}_2(\mathbb{C}) \rightarrow \mathrm{PGL}_2(\mathbb{C})$  is the canonical projection, then  $\phi$  is relevant for  $\mathrm{SL}_1(D)$ . In this case  $\Pi_\phi(\mathrm{SL}_1(D))$  is just the trivial representation of  $\mathrm{SL}_1(D)$ , and  $\Pi_\phi(\mathrm{SL}_2(K))$  consists of the Steinberg representation of  $\mathrm{SL}_2(K)$  – the unique irreducible square-integrable, non-supercuspidal representation.

All other principal series L-parameters are trivial on  $\mathrm{SL}_2(\mathbb{C})$  and are irrelevant for  $\mathrm{SL}_1(D)$ . By conjugating  $\phi$ , we may assume that its image is contained in the diagonal torus of  $\mathrm{PGL}_2(\mathbb{C})$ . One checks that  $Z_{\mathrm{PGL}_2(\mathbb{C})}(\phi)$  is connected unless the image of  $\phi$  is  $\{1, \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}\}$ . Whenever  $Z_{\mathrm{PGL}_2(\mathbb{C})}(\phi)$  is disconnected, its L-packet has two elements, see Subsection 3.5.

If  $Z_{\mathrm{PGL}_2(\mathbb{C})}(\phi)$  is connected, then  $\Pi_\phi(\mathrm{SL}_2(K))$  consists of precisely one principal series representation. Let  $T$  be the diagonal torus of  $\mathrm{SL}_2(K)$ , and let  $\chi_\phi$  be the character of  $T$  determined by local class field theory. Then  $\Pi_\phi(\mathrm{SL}_2(K))$  is the Langlands quotient of the parabolic induction of  $\chi_\phi$ , and the depth of that representation equals the depth of  $\chi_\phi$ .

### 3.3. Supercuspidal $L$ -packets of cardinality two.

For such  $L$ -parameters (16) shows that

$$\mathfrak{T}(\varphi) \cong \pi_0(Z_{\mathrm{PGL}_2(\mathbb{C})}(\phi)) \cong \mathbb{Z}/2\mathbb{Z}.$$

The  $L$ -parameter  $\phi$  is relevant for  $\mathrm{SL}_1(D)$ , so by (14)  $|\pi_0(Z_{\mathrm{SL}_2(\mathbb{C})}(\phi))| = 4$ . Then  $\pi_0(Z_{\mathrm{SL}_2(\mathbb{C})}(\phi))$  is either  $\mathbb{Z}/4\mathbb{Z}$  or  $(\mathbb{Z}/2\mathbb{Z})^2$ . In any case, it is abelian and has precisely four inequivalent characters. Now (15) says that

$$|\Pi_\phi(\mathrm{SL}_1(D))| = |\Pi_\phi(\mathrm{SL}_2(K))| = 2.$$

Now we classify the discrete  $L$ -parameters  $\phi$  for which the packet  $\Pi_\phi(\mathrm{SL}_2(K))$  is not a singleton. We note that every  $L$ -parameter for a supercuspidal representation of  $\mathrm{SL}_2(K)$  has to be trivial on  $\mathrm{SL}_2(\mathbb{C})$ . For if it were nontrivial on  $\mathrm{SL}_2(\mathbb{C})$ , then the image of  $\mathbf{W}_K$  would be in the centre of  $\mathrm{PGL}_2(\mathbb{C})$ , and we would get the  $L$ -parameter for the Steinberg representation, as discussed in the previous subsection. Since we want  $\phi$  to be discrete, it has to be an irreducible projective two-dimensional representation of  $\mathbf{W}_K$ .

Let  $\varphi$  be an irreducible two-dimensional representation of  $\mathbf{W}_K$  which lifts  $\phi$ . By (16) and (15) the associated  $L$ -packet  $\Pi_\phi(\mathrm{SL}_2(K))$  has more than one element if and only if  $\varphi$  is imprimitive. By [BuHe1, §41.3]  $\varphi$  is imprimitive if and only if there exists a separable quadratic extension  $E/K$  and a character  $\xi$  of  $E^\times$  such that  $\varphi \cong \mathrm{Ind}_{E/K}\xi$ . By the irreducibility  $\xi^\sigma \neq \xi$ , where  $\sigma$  is the nontrivial automorphism of  $E$  over  $K$ .

**Lemma 3.3.** *Let  $\phi$  and  $\varphi \cong \mathrm{Ind}_{E/K}\xi$  be as above.*

- (a) *Suppose that the character  $\xi^\sigma \xi^{-1}$  of  $E^\times$  has order two. Then  $\varphi$  is triply imprimitive and there exists a biquadratic extension  $L/K$  such that  $\ker(\phi) = \mathbf{W}_L$  and  $L \supset E$ .*
- (b) *Suppose that  $\xi^\sigma \xi^{-1}$  has order  $> 2$ . Then  $\varphi$  is simply imprimitive.*

*Proof.* Let  $\chi_E$  be the unique character of  $\mathbf{W}_K$  with kernel  $\mathbf{W}_E$ . Then  $\chi_E \in \mathfrak{T}(\varphi)$ , this holds in general for induction of irreducible representations from subgroups of index two. In particular  $|\mathfrak{T}(\varphi)| \in \{2, 4\}$ . From [BuHe1, Corollary 41.3] we see that  $\mathfrak{T}(\varphi) = \{1, \chi_E\}$  if and only if the character  $\xi^\sigma \xi^{-1}$  of  $\mathbf{W}_E$  cannot be lifted to a character of  $\mathbf{W}_F$ . Since the target group  $\mathbb{C}^\times$  is divisible, this happens if and only if  $\xi^\sigma \xi^{-1}$  does not equal

$$(\xi^\sigma \xi^{-1})^\sigma = \xi \xi^{-\sigma} = (\xi^\sigma \xi^{-1})^{-1}.$$

We conclude that the representation  $\varphi = \mathrm{Ind}_{E/K}\xi$  is triply imprimitive if  $\xi^\sigma \xi^{-1}$  has order two and is simply imprimitive otherwise.

Now we focus on the triply imprimitive case. By local class field theory there exists a unique separable quadratic extension  $L/E$  such that  $\xi^\sigma \xi^{-1}$  is the associated character  $\chi_L$  of  $E^\times$ . We consider it also as a character of  $\mathbf{W}_E$ . Then

$$\mathbf{W}_L = \ker(\chi_L) = \{w \in \mathbf{W}_K : \varphi(w) \in Z(\mathrm{GL}_2(\mathbb{C}))\}.$$

Hence  $\mathbf{W}_L = \ker(\phi)$  is a normal subgroup of  $\mathbf{W}_K$ , which means that  $L/K$  is a Galois extension. The explicit form of  $\varphi$  entails that the image of  $\phi$  is the Klein four group. Consequently

$$(17) \quad \text{Gal}(L/K) \cong \mathbf{W}_K/\mathbf{W}_L \cong \phi(\mathbf{W}_K) \cong (\mathbb{Z}/2\mathbb{Z})^2,$$

which says that  $L/K$  is biquadratic.  $\square$

We remark that the depth of  $\varphi = \text{Ind}_{E/K}\xi$  can be computed in the same way as for the imprimitive representations in Section 2, see in particular (11).

**3.4. Supercuspidal  $L$ -packets of cardinality four.** We continue with the case when  $\varphi$  is triply imprimitive, as in (17). This means that we have a biquadratic extension  $L/K$  and the Langlands parameter

$$(18) \quad \phi : W_K \rightarrow \text{Gal}(L/K) \cong (\mathbb{Z}/2\mathbb{Z})^2 \subset \text{PGL}_2(\mathbb{C}).$$

We also have

$$Z_{\text{PGL}_2(\mathbb{C})}(\text{im } \phi) = \pi_0(Z_{\text{PGL}_2(\mathbb{C})}(\text{im } \phi)) = S_\phi \cong (\mathbb{Z}/2\mathbb{Z})^2.$$

This implies, by (15), that  $\Pi_\phi(\text{SL}_2(K))$  is a supercuspidal packet of cardinality four.

We note the isomorphism  $\text{PGL}_2(\mathbb{C}) = \text{PSL}_2(\mathbb{C})$ , and the morphism

$$\text{SL}_2(\mathbb{C}) \rightarrow \text{PSL}_2(\mathbb{C}).$$

As in [We, §14], the pull-back  $\mathcal{S}_\phi$  of  $S_\phi$  is isomorphic to the quaternion group of order eight. This group admits four characters and one irreducible representation of degree two. Only the two-dimensional representation has nontrivial central character.

The parameter  $\phi$  creates a packet with five elements, which are allocated to  $\text{SL}_2(K)$  or  $\text{SL}_1(D)$  according to central characters. So  $\phi$  gives rise to an  $L$ -packet  $\Pi_\phi(\text{SL}_2(K))$  with four elements, and a singleton packet to the inner form  $\text{SL}_1(D)$ .

**Theorem 3.4.** *Let  $L/K$  be a biquadratic extension, let  $\phi$  be the Langlands parameter (18). If  $t$  is the highest break in the upper ramification of  $\text{Gal}(L/K)$  then  $d(\phi) = t$ . The allowed values of  $d(\phi)$  are 1, 3, 5, 7, ... except in Case 2.2 (see Appendix A.2), when the allowed values are 3, 5, 7, ...*

*Proof.* From the inclusion  $L \subset K_s$  we obtain a natural surjection

$$\pi_{L/K} : \text{Gal}(K_s/K) \rightarrow \text{Gal}(L/K).$$

Let  $K_{\text{ur}}$  be the maximal unramified extension of  $K$  in  $K_s$  and let  $K_{\text{ab}}$  be the maximal abelian extension of  $K$  in  $K_s$ . We have a commutative diagram, where the horizontal maps are the canonical maps and the vertical maps are the natural projections

$$\begin{array}{ccccccccc} 1 & \longrightarrow & I_{K_s/K} & \xrightarrow{\iota_1} & \text{Gal}(K_s/K) & \xrightarrow{p_1} & \text{Gal}(K_{\text{ur}}/K) & \longrightarrow & 1 \\ & & \alpha_1 \downarrow & & \pi_1 \downarrow & & id \downarrow & & \\ 1 & \longrightarrow & I_{K_{\text{ab}}/K} & \xrightarrow{\iota_2} & \text{Gal}(K_{\text{ab}}/K) & \xrightarrow{p_2} & \text{Gal}(K_{\text{ur}}/K) & \longrightarrow & 1 \\ & & \alpha_2 \downarrow & & \pi_2 \downarrow & & \beta \downarrow & & \\ 1 & \longrightarrow & \mathbf{I}_{L/K} & \xrightarrow{\iota_3} & \text{Gal}(L/K) & \xrightarrow{p_3} & \text{Gal}(L \cap K_{\text{ur}}/K) & \longrightarrow & 1 \end{array}$$

In the above notation, we have  $\pi_{L/K} = \pi_2 \circ \pi_1$ . Let

$$(19) \quad \dots \subset \mathbf{I}^{(2)} \subset \mathbf{I}^{(1)} \subset \mathbf{I}^{(0)} \subset G = \text{Gal}(L/K)$$

be the filtration of the relative inertia subgroup  $\mathbf{I}^{(0)} = \mathbf{I}_{L/K}$  of  $\text{Gal}(L/K)$ ,  $\mathbf{I}^{(1)}$  is the wild inertia subgroup, and so on. Note that  $\mathbf{I}^{(r)}$  is the restriction of the filtration  $G^r$  of  $G = \text{Gal}(L/K)$  to the subgroup  $\mathbf{I}_{L/K}$ , i.e,  $\mathbf{I}^{(r)} = \iota_3(G^r)$ . Let

$$(20) \quad \dots \subset I^{(2)} \subset I^{(1)} \subset I^{(0)} \subset G = \text{Gal}(\overline{K}/K)$$

be the filtration of the absolute inertia subgroup  $I^{(0)} = I_{K^s/K}$  of  $\text{Gal}(K^s/K)$ ,  $I^{(1)}$  is the wild inertia subgroup, and so on.

We have

$$(21) \quad (\forall r) \pi_{L/K} I^{(r)} = \mathbf{I}^{(r)}$$

This follows immediately from the above diagram. Here, we identify  $I^{(r)}$  with  $\iota_1(I^{(r)})$  and  $\mathbf{I}^{(r)}$  with  $\iota_3(\mathbf{I}^{(r)})$ . (Note that  $\alpha$  is *injective*. Therefore, by (21), we have

$$\phi(I^{(r)}) = 1 \iff (\alpha \circ \pi_{L/K})(I^{(r)}) = 1 \iff \alpha(\mathbf{I}^{(r)}) = 1 \iff \mathbf{I}^{(r)} = 1.$$

The *highest break*  $t$  has the property that  $I^{(t+1)} = 1$  and  $I^{(t)} \neq I^{(t+1)}$ . It follows that  $d(\phi) = t$ .

**Case 1:** There are two ramification breaks occurring at  $-1$  and some odd integer  $t > 0$ :

$$\{1\} = \dots = \mathbf{I}^{(t+1)} \subset \mathbf{I}^{(t)} = \dots = \mathbf{I}^{(0)} = \mathbf{I}_{L/K} \subset \text{Gal}(L/K), \quad d(\phi) = t.$$

The allowed depths are 1, 3, 5, 7, ...

**Case 2.1:** One single ramification break occurs at some odd integer  $t > 0$ :

$$\{1\} = \dots = \mathbf{I}^{(t+1)} \subset \mathbf{I}^{(t)} = \dots = \mathbf{I}^{(0)} = \mathbf{I}_{L/K} = \text{Gal}(L/K); \quad d(\phi) = t.$$

The allowed depths are 1, 3, 5, 7, ...

**Case 2.2:** There are two ramification breaks occurring at some odd integers  $t_1 < t_2$  (with  $\mathbf{I}^{(0)} = \mathbf{I}_{L/K}$ ):

$$\{1\} = \dots = \mathbf{I}^{(t_2+1)} \subset \mathbf{I}^{(t_2)} = \dots = \mathbf{I}^{(t_1+1)} \subset \mathbf{I}^{(t_1)} = \dots = \mathbf{I}^{(0)} = \mathbf{I}_{L/K} = \text{Gal}(L/K);$$

$$d(\phi) = t_2.$$

The allowed depths are 3, 5, 7, 9, ... □

Theorem 3.4 contrasts with the case of  $\text{SL}_2(\mathbb{Q}_p)$  with  $p > 2$ . Here there is a unique biquadratic extension  $L/K$ , and the associated L-parameter  $\phi : \text{Gal}(L/K) \rightarrow \text{SO}_3(\mathbb{R})$  has depth zero.

### 3.5. Principal series $L$ -packets of cardinality two.

Recall from Subsection 3.2 that a principal series L-parameter whose  $L$ -packet is not a singleton has image  $\{1, \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}\}$  in the diagonal torus  $T^\vee$  of  $\text{PGL}_2(\mathbb{C})$ . Thus it comes from a character  $\mathbf{W}_K \rightarrow \mathbb{C}^\times$  of order two. Define

$$\mathbf{W}_K \times \text{SL}_2(\mathbb{C}) \rightarrow K^\times$$

to be the projection  $(g, M) \mapsto g$  followed by the Artin reciprocity map

$$\mathbf{a}_K : \mathbf{W}_K \rightarrow K^\times.$$

Let  $E/K$  be a quadratic extension and let  $\chi_E$  be the associated quadratic character of  $K^\times$ . Consider the map

$$K^\times \rightarrow \mathrm{PGL}_2(\mathbb{C}), \quad \alpha \mapsto \begin{pmatrix} \chi_E(\alpha) & 0 \\ 0 & 1 \end{pmatrix}$$

The composite map

$$\phi_E: \mathbf{W}_K \times \mathrm{SL}_2(\mathbb{C}) \rightarrow K^\times \rightarrow \mathrm{PGL}_2(\mathbb{C})$$

is then an  $L$ -parameter attached to  $\chi_E$ . For the centralizer of the image, we have

$$Z_{\mathrm{PGL}_2(\mathbb{C})}(\mathrm{im} \phi_E) = N_{\mathrm{PGL}_2(\mathbb{C})}(T^\vee), \quad S_\phi \cong \mathcal{S}_\phi = \{1, w\},$$

where  $w$  generates the Weyl group of the dual group  $\mathrm{PGL}_2(\mathbb{C})$ . As there are two characters  $1, \epsilon$  of  $W = \{1, w\}$ , (15) says that the  $L$ -packet has cardinality two. There are two enhanced parameters  $(\phi_E, 1)$  and  $(\phi_E, \epsilon)$ , which parametrize the two elements in the  $L$ -packet  $\Pi_{\phi_E} = \Pi_{\phi_E}(\mathrm{SL}_2(K))$ . We will write

$$(22) \quad \Pi_{\phi_E} = \{\pi_E^1, \pi_E^2\}.$$

If  $\gamma \in K_s$  is a root of  $X^2 - X - \beta \in K[X]$ , the quadratic extension  $K(\gamma)$  is denoted also by  $K(\wp^{-1}(\beta))$ , with  $\beta \in K$ , where  $\wp(X) = X^2 - X$ . So the quadratic character

$$\chi_{n,j} = (-, u_j \wp^{-2n-1} + \wp(K)]$$

is associated with the quadratic extension  $E = K(\wp^{-1}(u_j \wp^{-2n-1}))$ , see (27) in the Appendix.

Let  $E/K$  be a quadratic extension. There are two kinds: the unramified one  $E_0 = K(\gamma_0)$  and countably many totally (and wildly) ramified  $E = K(\gamma)$ . The unramified quadratic extension has a single ramification break for  $t = -1$ .

Let  $E/K$  be a quadratic totally ramified extension. According to [Da, Proposition 11, p.411 and Proposition 14, p.413], there is a single ramification break for  $t = 2n + 1$ . Each value  $2n + 1$  occurs as a break, with  $n \geq 0, 1, 2, 3, \dots$ . By Theorem 3.4, adapted to the present case, we have

$$d(\phi_E) = 2n + 1.$$

Fix a basis  $\mathcal{B} = \{u_1, \dots, u_f\}$  of  $\mathbb{F}_q/\mathbb{F}_2$  and let  $u_j \in \mathcal{B}$ . The next result shows how to realise the extension  $E/K$ .

**Theorem 3.5.** *If  $E = K(\wp^{-1}(u_j \wp^{-2n-1}))$  then*

$$d(\phi_E) = 2n + 1$$

*with  $n = 0, 1, 2, 3, 4, \dots$*

*Proof.* Let  $\mathbf{a}_K : \mathbf{W}_K \rightarrow K^\times$  be the Artin reciprocity map. Then we have [ABPS1, Theorem 3.6]:

$$\mathbf{a}_K(\mathrm{Gal}(K_s/K)^l) = U^{[l]}$$

for all  $l \geq 0$ , where  $[l]$  denotes the least integer greater than or equal to  $l$ , and  $U_K^i$  is the  $i$ th higher unit group.

We are concerned here with the quadratic character  $\chi = \chi_E$  and the associated  $L$ -parameter  $\phi = \phi_E$ . The level  $\ell(\chi)$  of  $\chi$  is the least integer  $n \geq 0$  for which  $\chi(U_K^{n+1}) = 1$ . Call this integer  $N$ . For this integer  $N$ , we have

$$N < l \leq N + 1 \implies \mathbf{a}_K(\mathrm{Gal}(K_s/K)^l) = U_K^{[l]} = U_K^{N+1} \text{ on which } \chi \text{ is trivial}$$

$N - 1 < l \leq N \implies \mathbf{a}_K(\text{Gal}(K_s/K)^l) = U_K^{[l]} = U_K^N$  on which  $\chi$  is nontrivial

The  $L$ -parameter  $\phi$  will factor through  $K^\times$  and we have to consider its depth  $d(\phi)$ . Recall: the depth of  $\phi$  is the smallest number  $d(\phi) \geq 0$  such that  $\phi$  is trivial on  $\text{Gal}(K_s/K)^l$  for all  $l > d(\phi)$ . Then  $d(\phi) = N$  in view of the above two implications. We infer that

$$(23) \quad \ell(\chi_E) = d(\phi_E).$$

If  $\chi$  is the unramified quadratic character given by  $\chi(x) = (-1)^{\text{val}_K(x)}$  then we will have to allow  $N = -1$  in which case  $\phi$  has negative depth.

If  $E = K(\wp^{-1}(u_j \wp^{-2n-1}))$  then  $\chi_E = \chi_{n,j}$  and so we have

$$(24) \quad \ell(\chi_E) = \ell(\chi_{n,j}).$$

We now compute the level of the quadratic character  $\chi_{n,j}$  defined in (27). Every  $\alpha \in U_K^i$  has the form  $\alpha = 1 + \varepsilon \wp^i$ , with  $\varepsilon \in \mathfrak{o}$ , and can be expanded in the convergent product

$$\alpha = \prod_{i \geq 1} (1 + \theta_i \wp^i)$$

for unique  $\theta_i \in \mathbb{F}_q$ . As we can see in the proof of Theorem A.2,

$$d_\wp(1 + \theta_{2n+1} \wp^{2n+1}, u_j \wp^{-2n-1}) = \text{Tr}_{\mathbb{F}_q/\mathbb{F}_2}(u_j \theta_{2n+1})$$

and

$$d_\wp(1 + \theta_i \wp^i, u_j \wp^{-2n-1}) = 0$$

if  $i \nmid 2n + 1$ . There exists, therefore, an element  $\alpha \in U_K^{2n+1}$  such that  $\chi_{n,j}(\alpha) \neq 0$  and  $\chi_{n,j}(U_K^{2n+2}) = 1$ . We infer that

$$(25) \quad \ell(\chi_{n,j}) = 2n + 1.$$

The theorem now follows from (23), (24) and (25).  $\square$

We conclude that, if  $E = K(\wp^{-1}(u_j \wp^{-2n-1}))$ , then

$$d(\pi_E^i) \geq 2n + 1$$

with  $i = 1, 2$ .

It follows that the depths of the irreducible representations  $\pi_E^1, \pi_E^2$  in the  $L$ -packet  $\Pi_{\phi_E}$  can be arbitrarily large. For representations of enormous depth, such as the ones encountered in this article, the term *hadopelagic* commends itself, in contrast to the currently accepted term *epipelagic* for representations of modest depth, see [en.wikipedia.org/wiki/Epipelagic](http://en.wikipedia.org/wiki/Epipelagic).

#### APPENDIX A. ARTIN-SCHREIER SYMBOL

Let  $K$  be a local field of characteristic  $p$  with finite residue field  $k$ . The field of constants  $k = \mathbb{F}_q$  is a finite extension of  $\mathbb{F}_p$ , with degree  $[k : \mathbb{F}_p] = f$  and  $q = p^f$ . Let  $\mathfrak{o}$  be the ring of integers in  $K$  and  $\mathfrak{p} \subset \mathfrak{o}$  the maximal ideal. A choice of uniformizer  $\wp \in \mathfrak{o}$  determines isomorphisms  $K \cong \mathbb{F}_q((\wp))$ ,  $\mathfrak{o} \cong \mathbb{F}_q[[\wp]]$  and  $\mathfrak{p} = \wp \mathfrak{o} \cong \wp \mathbb{F}_q[[\wp]]$ . The group of units is denoted by  $\mathfrak{o}^\times$  and  $\nu$  represents a normalized valuation on  $K$ , so that  $\nu(\wp) = 1$  and  $\nu(K) = \mathbb{Z}$ .

Following [FeVo, IV.4 - IV.5], we have the reciprocity map

$$\Psi_K : K^\times \rightarrow \text{Gal}(K_{\text{ab}}/K)$$

We define the map (Artin-Schreier symbol)

$$(-, -] : K^\times \times K \rightarrow \mathbb{F}_p$$

by the formula

$$(\alpha, \beta] = \Psi_K(\alpha)(\gamma) - \gamma$$

where  $\gamma$  is a root of the polynomial  $X^p - X - \beta$ . The polynomial  $X^p - X$  is denoted  $\wp(X)$ . According to [FeVo, p.148] the pairing  $(-, -]$  determines the nondegenerate pairing

$$(26) \quad K^\times / K^{\times p} \times K / \wp(K) \rightarrow \mathbb{F}_p.$$

Let us fix a coset  $\beta + \wp(K) \in K / \wp(K)$ . According to (26), this coset determines an element of  $\text{Hom}(K^\times / K^{\times p}, \mathbb{F}_p)$ .

Now specialise to  $p = 2$ . We will identify the additive group  $\mathbb{F}_2$  with the multiplicative group  $\mu_2(\mathbb{C}) = \{1, -1\} \subset \mathbb{C}$ . In that case, the elements of  $\text{Hom}(K^\times / K^{\times 2}, \mathbb{F}_2)$  are precisely the quadratic characters of  $K^\times$ . Since the pairing (26) is nondegenerate, the quadratic characters are parametrised by the cosets  $\beta + \wp(K) \in K / \wp(K)$ . Now the index of  $\wp(K)$  in  $K$  is infinite; in fact, the powers  $\{\varpi^{-2n-1} : n \geq 0\}$  are distinct coset representatives, see [FeVo, p.146].

**Lemma A.1.** *For  $K = \mathbb{F}_2((\varpi))$  the set of powers  $\{\varpi^{-2n-1} : n \geq 0\}$  is a complete set of coset representatives.*

That is not the case when  $K = \mathbb{F}_q((\varpi))$  has residue degree  $f > 1$ . Let  $\mathcal{B} = \{u_1, \dots, u_f\}$  denote a basis of the  $\mathbb{F}_2$ -linear space  $\mathbb{F}_q$ . Then,

$$\{u_j \varpi^{-2n-1} : n \geq 0, j = 1, \dots, f\}$$

is a complete set of coset representatives of  $K / \wp(K)$ , see §5 and §6 of [Da].

The pairing (26) creates a sequence of quadratic characters

$$(27) \quad \chi_{n,j}(\alpha) := (\alpha, u_j \varpi^{-2n-1} + \wp(K)]$$

with  $n \geq 0$  and  $j = 1, \dots, f$ .

#### A.1. Explicit formula for the Artin-Schreier symbol.

In [FeVo, Corollary 5.5, p.148], the authors introduce the map  $d_\varpi$  which we now describe. Let  $\varpi$  be a fixed uniformizer. Using the isomorphism  $K = \mathbb{F}_q((\varpi))$ , where  $q = 2^f$ , every element  $\alpha \in K$  can be uniquely expanded as

$$(28) \quad \alpha = \sum_{i \geq i_a} \vartheta_i \varpi^i, \quad \vartheta_i \in \mathbb{F}_q.$$

Put

$$\frac{d\alpha}{d\varpi} = \sum_{i \geq i_a} i \vartheta_i \varpi^{i-1}, \quad \text{res}_\varpi(\alpha) = \vartheta_{-1}.$$

Define the pairing

$$(29) \quad d_\varpi : K^\times \times K \rightarrow \mathbb{F}_2, \quad d_\varpi(\alpha, \beta) = \text{Tr}_{\mathbb{F}_q/\mathbb{F}_2} \text{res}_\varpi(\beta \alpha^{-1} \frac{d\alpha}{d\varpi})$$

By [FeVo, Theorem 5.6. p.149], the pairing  $(-, -]$  coincides with the pairing defined in (29). In particular,  $d_\varpi$  does not depend on the choice of uniformizer.



We conclude that every quadratic character  $\chi_{n,j}$  from (27) is completely described by

$$(30) \quad d_{\varpi}(\alpha, u_j \varpi^{-2n-1}) = \mathrm{Tr}_{\mathbb{F}_q/\mathbb{F}_2} \mathrm{res}_{\varpi}(u_j \varpi^{-2n-1} \alpha^{-1} \frac{d\alpha}{d\varpi}), \quad n \geq 0.$$

We seek a formula more explicit than (30).

By [FeVo, Proposition 5.10, p. 17], for every  $\alpha \in K^\times$  there exist uniquely determined  $k \in \mathbb{Z}$  and  $\theta_i \in \mathbb{F}_q$  for  $i \geq 0$  such that  $\alpha$  can be expanded in the convergent product

$$(31) \quad \alpha = \varpi^k \theta_0 \prod_{i \geq 1} (1 + \theta_i \varpi^i)$$

We have

$$\begin{aligned} d_{\varpi}(\varpi^k \theta_0 \prod_{i \geq 1} (1 + \theta_i \varpi^i), u_j \varpi^{-2n-1}) &= \\ &= d_{\varpi}(\theta_0 \varpi^k, u_j \varpi^{-2n-1}) + d_{\varpi}(\prod_{i \geq 1} (1 + \theta_i \varpi^i), u_j \varpi^{-2n-1}) \end{aligned}$$

Now,  $d_{\varpi}(\theta_0 \varpi^k, u_j \varpi^{-2n-1})$  is easy to compute:

$$\begin{aligned} d_{\varpi}(\theta_0 \varpi^k, u_j \varpi^{-2n-1}) &= \mathrm{Tr}_{\mathbb{F}_q/\mathbb{F}_2} \mathrm{res}_{\varpi}(u_j \varpi^{-2n-1} \theta_0^{-1} \varpi^{-k} \frac{d(\theta_0 \varpi^k)}{d\varpi}) \\ &= \mathrm{Tr}_{\mathbb{F}_q/\mathbb{F}_2} \mathrm{res}_{\varpi}(k u_j \varpi^{-2n-2}) \\ &= 0. \end{aligned}$$

On the other hand,

$$\begin{aligned} d_{\varpi}(\prod_{i \geq 1} (1 + \theta_i \varpi^i), u_j \varpi^{-2n-1}) &= \sum_{i \geq 1} d_{\varpi}(1 + \theta_i \varpi^i, u_j \varpi^{-2n-1}) \\ &= \sum_{i=1}^{2n+1} d_{\varpi}(1 + \theta_i \varpi^i, u_j \varpi^{-2n-1}) \end{aligned}$$

since  $d_{\varpi}(1 + \theta_i \varpi^i, u_j \varpi^{-2n-1}) = 0$  if  $i > 2n+1$  (see [FeVo, p. 150], proof of Corollary). Moreover, by the same proof of Corollary in [FeVo, p. 150], we have

$$(32) \quad \begin{aligned} d_{\varpi}(1 + \theta_{2n+1} \varpi^{2n+1}, u_j \varpi^{-2n-1}) &= \mathrm{Tr}_{\mathbb{F}_q/\mathbb{F}_2}((2n+1) u_j \theta_{2n+1}) \\ &= \mathrm{Tr}_{\mathbb{F}_q/\mathbb{F}_2}(u_j \theta_{2n+1}). \end{aligned}$$

This last formula is a particular case of a more general formula we are about to prove.

In order to compute  $d_{\varpi}(1 + \theta_i \varpi^i, u_j \varpi^{-2n-1})$  for  $i = 1, \dots, 2n+1$ , we need to find the Laurent series expansion of  $(1 + \theta_i \varpi^i)^{-1}$ . This can be done by expanding the geometric series

$$(1 + \theta_i \varpi^i)^{-1} = \sum_{j \geq 0} (-\theta_i \varpi^i)^j = 1 - \theta_i \varpi^i + \theta_i^2 \varpi^{2i} - \theta_i^3 \varpi^{3i} + \dots$$

We have

$$u_j \varpi^{-2n-1} (1 + \theta_i \varpi^i)^{-1} \frac{d}{d\varpi} (1 + \theta_i \varpi^i) = \\ i u_j \theta_i \varpi^{-2n-1+i-1} (1 - \theta_i \varpi^i + \theta_i^2 \varpi^{2i} - \theta_i^3 \varpi^{3i} + \dots + (-1)^r \theta_i^r \varpi^{ri} + \dots)$$

The residue will be nonzero if

$$-2n - 1 + i - 1 + ri = -1 \Leftrightarrow r = \frac{2n+1}{i} - 1$$

Hence,  $d_\varpi(1 + \theta_i \varpi^i, u_j \varpi^{-2n-1}) = 0$  if  $i \nmid 2n+1$ . In particular,  $i$  must be odd.

We have:

$$d_\varpi(1 + \theta_i \varpi^i, u_j \varpi^{-2n-1}) = \begin{cases} 0 & , \quad \text{if } i \nmid 2n+1 \\ \text{Tr}_{\mathbb{F}_q/\mathbb{F}_2}(u_j \theta_i^{(2n+1)/i}) & , \quad \text{if } i \mid 2n+1 \end{cases}$$

In particular, we recover formula (32) by taking  $i = 2n+1$ .

From the above, we have established the following explicit formula.

**Theorem A.2.** *Let  $K$  be a local function field of characteristic 2 with residue degree  $f$ , and let  $\chi_{n,j}$  denote the quadratic character from (27). Then we have the explicit formula*

$$\chi_{n,j}(\alpha) = \sum_{i \mid 2n+1} \text{Tr}_{\mathbb{F}_q/\mathbb{F}_2}(u_j \theta_i^{(2n+1)/i})$$

where  $\alpha = \varpi^k \theta_0 \prod_{i \geq 1} (1 + \theta_i \varpi^i) \in K^\times$ ,  $n \geq 0$  and  $j = 1, \dots, f$ .

For example, we have

$$\chi_{0,1}(\alpha) = \text{Tr}_{\mathbb{F}_q/\mathbb{F}_2} \theta_1, \quad \chi_{1,1}(\alpha) = \text{Tr}_{\mathbb{F}_q/\mathbb{F}_2}(\theta_1^3 + \theta_3), \quad \chi_{2,1}(\alpha) = \text{Tr}_{\mathbb{F}_q/\mathbb{F}_2}(\theta_1^5 + \theta_5),$$

where  $\{1, u_2, \dots, u_f\}$  is a basis of  $\mathbb{F}_q/\mathbb{F}_2$ .

## A.2. Ramification.

Quadratic extensions  $L/K$  are obtained by adjoining an  $\mathbb{F}_2$ -line  $D \subset K/\wp(K)$ . Therefore,  $L = K(\wp^{-1}(D)) = K(\gamma)$  where  $D = \text{span}\{\beta + \wp(K)\}$ , with  $\gamma^2 - \gamma = \beta$ . In particular, if  $\beta_0 \in \mathfrak{o} \setminus \mathfrak{p}$  such that the image of  $\beta_0$  in  $\mathfrak{o}/\mathfrak{p}$  has nonzero trace in  $\mathbb{F}_2$ , the  $\mathbb{F}_2$ -line  $V_0 = \text{span}\{\beta_0 + \wp(K)\}$  contains all the cosets  $\beta_i + \wp(K)$  where  $\beta_i$  is an integer and so  $K(\wp^{-1}(\mathfrak{o})) = K(\wp^{-1}(V_0)) = K(\gamma_0)$  where  $\gamma_0^2 - \gamma_0 = \beta_0$  gives the unramified quadratic extension, see [Da, Proposition 12, p. 412].

Biquadratic extensions are computed the same way, by considering  $\mathbb{F}_2$ -planes  $W = \text{span}\{\beta_1 + \wp(K), \beta_2 + \wp(K)\} \subset K/\wp(K)$ . Therefore, if  $\beta_1 + \wp(K)$  and  $\beta_2 + \wp(K)$  are  $\mathbb{F}_2$ -linearly independent then  $K(\wp^{-1}(W)) := K(\gamma_1, \gamma_2)$  is biquadratic, where  $\gamma_1^2 - \gamma_1 = \beta_1$  and  $\gamma_2^2 - \gamma_2 = \beta_2$ ,  $\gamma_1, \gamma_2 \in K^s$ . Therefore,  $K(\gamma_1, \gamma_2)/K$  is biquadratic if  $\beta_2 - \beta_1 \notin \wp(K)$ .

A biquadratic extension containing the line  $V_0$  is of the form  $K(\gamma_0, \gamma)/K$ . There are countably many quadratic extensions  $L_0/K$  containing the unramified quadratic extension. They have ramification index  $e(L_0/K) = 2$ . And there are countably many biquadratic extensions  $L/K$  which do not contain the unramified quadratic extension. They have ramification index  $e(L/K) = 4$ .

So, there is a plentiful supply of biquadratic extensions  $K(\gamma_1, \gamma_2)/K$ .

The space  $K/\wp(K)$  comes with a filtration

$$(33) \quad 0 \subset_1 V_0 \subset_f V_1 = V_2 \subset_f V_3 = V_4 \subset_f \dots \subset K/\wp(K)$$

where  $V_0$  is the image of  $\mathfrak{o}_K$  and  $V_i$  ( $i > 0$ ) is the image of  $\mathfrak{p}^{-i}$  under the canonical surjection  $K \rightarrow K/\wp(K)$ . For  $K = \mathbb{F}_q((\varpi))$  and  $i > 0$ , each inclusion  $V_{2i} \subset_f V_{2i+1}$  is a sub- $\mathbb{F}_2$ -space of codimension  $f$ . The  $\mathbb{F}_2$ -dimension of  $V_n$  is

$$(34) \quad \dim_{\mathbb{F}_2} V_n = 1 + \lceil n/2 \rceil f,$$

for every  $n \in \mathbb{N}$ , where  $\lceil x \rceil$  is the smallest integer bigger than  $x$ .

Let  $L/K$  denote a Galois extension with Galois group  $G$ . For each  $i \geq -1$  we define the  $i^{\text{th}}$ -ramification subgroup of  $G$  (in the lower numbering) to be:

$$G_i = \{\sigma \in G : \sigma(x) - x \in \mathfrak{p}_L^{i+1}, \forall x \in \mathfrak{o}_L\}.$$

An integer  $t$  is a *break* for the filtration  $\{G_i\}_{i \geq -1}$  if  $G_t \neq G_{t+1}$ . The study of ramification groups  $\{G_i\}_{i \geq -1}$  is equivalent to the study of breaks of the filtration.

There is another decreasing filtration with upper numbering  $\{G^i\}_{i \geq -1}$  and defined by the *Hasse-Herbrand function*  $\psi = \psi_{L/K}$ :

$$G^u = G_{\psi(u)}.$$

In particular,  $G^{-1} = G_{-1} = G$  and  $G^0 = G_0$ , since  $\psi(-1) = -1$  and  $\psi(0) = 0$ .

Now, in analogy with the lower notation, a real number  $t \geq -1$  is a *break* for the filtration  $\{G^i\}_{i \geq -1}$  if

$$(35) \quad \forall \varepsilon > 0, G^t \neq G^{t+\varepsilon}.$$

We define

$$(36) \quad G^{t+} := \bigcap_{r>t} G^r.$$

Then  $t$  is a break of the filtration if and only if  $G^{t+} \neq G^t$ . The set of breaks of the filtration is countably infinite and need not consist of integers.

If  $G$  is abelian, it follows from Hasse-Arf theorem [FeVo, p.91] that the breaks are integers and (35) is equivalent to

$$G^t \neq G^{t+1}.$$

Let  $G_2 = \text{Gal}(K_2/K)$  be the Galois group of the maximal abelian extension of exponent 2,  $K_2 = K(\wp^{-1}(K))$ . Since  $G_2 \cong K^\times/K^{\times 2}$ , the nondegenerate pairing (26) coincides with the pairing  $G_2 \times K/\wp(K) \rightarrow \mathbb{Z}/2\mathbb{Z}$ .

The profinite group  $G_2$  comes equipped with a ramification filtration  $(G_2^u)_{u \geq -1}$  in the upper numbering, see [Da, p.409]. For  $u \geq 0$ , we have an orthogonal relation [Da, Proposition 17, p.415]

$$(37) \quad (G_2^u)^\perp = \overline{\mathfrak{p}^{-\lceil u \rceil + 1}} = V_{\lceil u \rceil - 1}$$

under the pairing  $G_2 \times K/\wp(K) \rightarrow \mathbb{Z}/2\mathbb{Z}$ .

Since the upper filtration is more suitable for quotients, we will compute the upper breaks. By using the Hasse-Herbrand function it is then possible to compute the lower breaks in order to obtain the lower ramification filtration.

According to [Da, Proposition 17], the positive breaks in the filtration  $(G^v)_v$  occur precisely at integers prime to  $p$ . So, for  $ch(K) = 2$ , the positive breaks will occur at odd integers. The lower numbering breaks are also integers. If  $G$  is cyclic of prime order, then there is a unique break for any decreasing filtration  $(G^v)_v$  (see [Da, Proposition 14]). In general, the number of breaks depends on the possible filtration of the Galois group.

Given a plane  $W \subset K/\wp(K)$ , the filtration (33)  $(V_i)_i$  on  $K/\wp(K)$  induces a filtration  $(W_i)_i$  on  $W$ , where  $W_i = W \cap V_i$ . There are three possibilities for the filtration breaks on a plane and we will consider each case individually.

**Case 1:**  $W$  contains the line  $V_0$ , i.e.  $L_0 = K(\wp^{-1}(W))$  contains the unramified quadratic extension  $K(\wp^{-1}(V_0)) = K(\alpha_0)$  of  $K$ . The extension has residue degree  $f(L_0/K) = 2$  and ramification index  $e(L_0/K) = 2$ . In this case, there is an integer  $t > 0$ , necessarily odd, such that the filtration  $(W_i)_i$  looks like

$$0 \subset_1 W_0 = W_{t-1} \subset_1 W_t = W.$$

By the orthogonality relation (37), the upper ramification filtration on  $G = \text{Gal}(L_0/K)$  looks like

$$\{1\} = \dots = G^{t+1} \subset_1 G^t = \dots = G^0 \subset_1 G^{-1} = G$$

Therefore, the upper ramification breaks occur at  $-1$  and  $t$ .

The number of such  $W$  is equal to the number of planes in  $V_t$  containing the line  $V_0$  but not contained in the subspace  $V_{t-1}$ . This number can be computed and equals the number of biquadratic extensions of  $K$  containing the unramified quadratic extensions and with a pair of upper ramification breaks  $(-1, t)$ ,  $t > 0$  and odd. Here is an example.

**Example A.3.** *The number of biquadratic extensions containing the unramified quadratic extension and with a pair of upper ramification breaks  $(-1, 1)$  is equal to the number of planes in an  $1+f$ -dimensional  $\mathbb{F}_2$ -space, containing the line  $V_0$ . There are precisely*

$$1 + 2 + 2^2 + \dots + 2^{f-1} = \frac{1 - 2^f}{1 - 2} = q - 1$$

*of such biquadratic extensions.*

**Case 2.1:**  $W$  does not contains the line  $V_0$  and the induced filtration on the plane  $W$  looks like

$$0 = W_{t-1} \subset_2 W_t = W$$

for some integer  $t$ , necessarily odd.

The number of such  $W$  is equal to the number of planes in  $V_t$  whose intersection with  $V_{t-1}$  is  $\{0\}$ . Note that, there are no such planes when  $f = 1$ . So, for  $K = \mathbb{F}_2((\varpi))$ , **case 2.1** does not occur.

Suppose  $f > 1$ . By the orthogonality relation, the upper ramification filtration on  $G = \text{Gal}(L/K)$  looks like

$$\{1\} = \dots = G^{t+1} \subset_2 G^t = \dots = G^{-1} = G$$

Therefore, there is a single upper ramification break occurring at  $t > 0$  and is necessarily odd.

For  $f = 1$  there is no such biquadratic extension. For  $f > 1$ , the number of these biquadratic extensions equals the number of planes  $W$  contained in an  $\mathbb{F}_2$ -space of dimension  $1 + fi$ ,  $t = 2i - 1$ , which are transverse to a given codimension- $f$   $\mathbb{F}_2$ -space.

**Case 2.2:**  $W$  does not contains the line  $V_0$  and the induced filtration on the plane  $W$  looks like

$$0 = W_{t_1-1} \subset_1 W_{t_1} = W_{t_2-1} \subset_1 W_{t_2} = W$$

for some integers  $t_1$  and  $t_2$ , necessarily odd, with  $0 < t_1 < t_2$ .

The orthogonality relation for this case implies that the upper ramification filtration on  $G = \text{Gal}(L/K)$  looks like

$$\{1\} = \dots = G^{t_2+1} \subset_1 G^{t_2} = \dots = G^{t_1+1} \subset_1 G^{t_1} = \dots = G$$

The upper ramification breaks occur at odd integers  $t_1$  and  $t_2$ .

There is only a finite number of such biquadratic extensions, for a given pair of upper breaks  $(t_1, t_2)$ .

#### REFERENCES

- [ABPS1] A.-M. Aubert, P. Baum, R.J. Plymen, M. Solleveld, Depth and the local Langlands correspondence, Arbeitstagung Bonn 2013, Progress in Math. 319, Birkhauser 2016, [arxiv.org/abs/1311.1606](https://arxiv.org/abs/1311.1606)
- [ABPS2] A.-M. Aubert, P. Baum, R.J. Plymen, M. Solleveld, The local Langlands correspondence for inner forms of  $\text{SL}_n$ , Res. Math. Sci., to appear, [arxiv.org/abs/1305.2638](https://arxiv.org/abs/1305.2638)
- [BuHe1] C. Bushnell, G. Henniart, *The local Langlands conjecture for  $\text{GL}(2)$* , Grundlehren der math. Wissenschaften, **335**, Springer, 2006.
- [BuHe2] C. Bushnell, G. Henniart, *The essentially tame local Langlands correspondence, III: the general case*, Proc. Lond. Math. Soc. (3) **101** (2010), no. 2, 497–553.
- [BuKu] C.J. Bushnell, P.C. Kutzko, The admissible dual of  $\text{SL}(N)$  II, Proc. London Math. Soc. (3) **68** (1994), 317–379.
- [Da] C.S. Dalawat, Further remarks on local discriminants, J. Ramanujan Math. Soc., (4) 25 (2010) 393–417.
- [DBHCS] S. DeBacker, Harish-Chandra, P.J. Sally, *Admissible invariant distributions on reductive  $p$ -adic groups*, University Lecture Series **16**, American Mathematical Society, Providence RI, 1991.
- [Del] P. Deligne, “Les corps locaux de caractéristique  $p$ , limites de corps locaux de caractéristique 0”, pp. 119–157 in: *Représentations des groupes réductifs sur un corps local*, Travaux en cours, Hermann, 1984.
- [FeVo] I. B. Fesenko and S.V. Vostokov, *Local fields and their extensions*, 2nd ed., AMS Math. Monograph Translation 121, 2002.
- [GeKn] S.S. Gelbart, A.W. Knap, L-indistinguishability and R groups for the special linear group, Adv. in Math. **43** (1982), 101–121.
- [GrRe] B.H. Gross, M. Reeder, Arithmetic invariants of discrete Langlands parameters, Duke Math. J. **154** (2010), 431–508.
- [HiSa] K. Hiraga, H. Saito, On L-packets for inner forms of  $\text{SL}_n$ , Mem. Amer. Math. Soc. **1013**, Vol. **215**, 2012.
- [JaLa] H. Jacquet, R. Langlands, *Automorphic forms on  $\text{GL}(2)$* , Lecture Notes in Mathematics **114**, Springer-Verlag, Berlin, 1970.
- [MoPr] A. Moy, G. Prasad, Unrefined minimal  $K$ -types for  $p$ -adic groups, Inv. Math. **116** (1994), 393–408.
- [Ne] J. Neukirch, Algebraic Number Theory, Springer-Verlag, Berlin, 1999.
- [Ser] J.-P. Serre, *Corps locaux*. Hermann, Paris, 1962.
- [She] D. Shelstad, Notes on  $L$ -indistinguishability, Proc. Symp. Pure Math. **33**, part.2 (1979), 193–203.
- [We] A. Weil, Exercices dyadiques, Invent. Math. 27 (1974) 1–22.

INSTITUT DE MATHÉMATIQUES DE JUSSIEU – PARIS RIVE GAUCHE, U.M.R. 7586 DU C.N.R.S.,  
U.P.M.C., 4 PLACE JUSSIEU 75005 PARIS, FRANCE

*E-mail address:* `anne-marie.aubert@imj-prg.fr`

ISCTE - LISBON UNIVERSITY INSTITUTE, AV. DAS FORÇAS ARMADAS, 1649-026, LISBON,  
PORTUGAL

*E-mail address:* `sergio.mendes@iscte.pt`

SCHOOL OF MATHEMATICS, SOUTHAMPTON UNIVERSITY, SOUTHAMPTON SO17 1BJ, ENGLAND  
and SCHOOL OF MATHEMATICS, MANCHESTER UNIVERSITY, MANCHESTER M13 9PL, ENGLAND

*E-mail address:* `r.j.plymen@soton.ac.uk` `plymen@manchester.ac.uk`

IMAPP, RADBOUD UNIVERSITEIT, HEYENDAALSEWEG 135, 6525AJ, NIJMEGEN, THE NETHER-  
LANDS

*E-mail address:* `m.solleveld@science.ru.nl`