

Number Fields

Errata, version 2023-1
update February 20, 2024

- page 52 line -14: $p = 2$ and $m \equiv 3 \pmod{4}$, in which case p ramifies.
- page 64 exercise 18: with $p \nmid \text{disc}(f)$. Let $K = \mathbb{Q}(\alpha)$. Show that
- page 64 exercise 19: and $p^2 \nmid \text{disc}(f)$. Let $K = \mathbb{Q}(\alpha)$. Prove that the ideal
- page 127 line 13: $(\psi(\varepsilon_1, \dots, \psi(\varepsilon_{r+s-1}))$
- page 142 exercises 5 and 6: Add: Put $P = \max(\mathcal{O}_K) \setminus \{\mathfrak{p}\}$. Replace $K_{\mathfrak{p}}$ and $K_{\mathfrak{p}}^*$ by K_P and K_P^* respectively.
- page 142 exercise 8: Dedekind domains
- page 152 line 1: $3\gamma + \text{Tr}_{\mathbb{Q}}^L(\gamma) \in \mathbb{Z}[\zeta_3] + \mathbb{Z}[\alpha] + \mathbb{Z}[\zeta_3\alpha] + \mathbb{Z}[\zeta_3^2\alpha]$
- page 209 line -7: is the next lemma.
- page 485 line -9: of right cosets of U in G .
- page 485 line -8: a partition of $U \setminus G$ into orbits of cosets.
- page 488: The proof of Proposition 18.43 refers to Theorem 18.34. However, this theorem applies only to abelian groups. For the following proof this condition is not needed.

PROOF. For $U \in \Sigma(G)$ we have $r_U + 2s_U = [L^U : \mathbb{Q}]$. So

$$\sum_{U \in \Sigma(G)} n_U(r_U + 2s_U)\#(U) = \sum_{U \in \Sigma(G)} n_U[L^U : \mathbb{Q}][L : L^U] = [L : \mathbb{Q}] \sum_{U \in \Sigma(G)} n_U = 0$$

In the proof of Theorem 7.53 the splitting of a prime ideal in an intermediate field of a Galois extension is used. This applies equally well to the splitting of infinite primes. Let \mathfrak{q} be an infinite prime of L . Note that for infinite primes the inertia groups coincide with the decomposition groups. In Theorem 18.38 it is shown that $\sum_U n_U \#(U) t_{\mathfrak{p}, U} = 0$, where t_U is the number of prime ideals of \mathcal{O}_{L^U} above a given prime ideal \mathfrak{p} of K with a given residue class degree. For infinite primes the same

holds, all residue class degrees being 1. Summation over all infinite primes of K yields

$$\begin{aligned} \sum_{U \in \Sigma(G)} n_U \#(U)(r_U + s_U) &= \sum_{U \in \Sigma(G)} n_U \#(U) \sum_{\mathfrak{p} \in \mathcal{P}_\infty(K)} t_{\mathfrak{p},U} \\ &= \sum_{\mathfrak{p} \in \mathcal{P}_\infty(K)} \sum_{U \in \Sigma(G)} n_U \#(U) t_{\mathfrak{p},U} = 0. \end{aligned}$$