

Entanglement of completely symmetric quantum states

Hans Maassen, Radboud University Nijmegen, the Netherlands

QP-NCG-QI, Edinburgh January 18, 2012.

Collaboration with Burkhard Kümmerner and Bram Petri

Statement of the problem

Statement of the problem

We consider a quantum system consisting of n identical, but distinguishable subsystems ("particles") described by Hilbert spaces of dimension d .

Statement of the problem

We consider a quantum system consisting of n identical, but distinguishable subsystems ("particles") described by Hilbert spaces of dimension d .

A state on such a system is called **completely symmetric** if it is symmetric both for the global rotation of all the individual Hilbert spaces together ("**Werner state**") and for permutations of the particles.

Statement of the problem

We consider a quantum system consisting of n identical, but distinguishable subsystems ("particles") described by Hilbert spaces of dimension d .

A state on such a system is called **completely symmetric** if it is symmetric both for the global rotation of all the individual Hilbert spaces together ("**Werner state**") and for permutations of the particles.

The state is called **entangled** if it can not be written as a convex combination of product states.

Statement of the problem

We consider a quantum system consisting of n identical, but distinguishable subsystems ("particles") described by Hilbert spaces of dimension d .

A state on such a system is called **completely symmetric** if it is symmetric both for the global rotation of all the individual Hilbert spaces together ("**Werner state**") and for permutations of the particles.

The state is called **entangled** if it can not be written as a convex combination of product states.

For a given completely symmetric state, we want to find out if it is entangled or not, and, if so, to quantify how entangled it is.

Motivation

Motivation

Entanglement is a central issue in quantum information theory.

Motivation

Entanglement is a central issue in quantum information theory.

The study of n party-entanglement is considered difficult. It is complicated by the fact that the state space of n systems of size d has a large dimension: $d^{2n} - 1$.

Motivation

Entanglement is a central issue in quantum information theory.

The study of n party-entanglement is considered difficult. It is complicated by the fact that the state space of n systems of size d has a large dimension: $d^{2^n} - 1$.

The number of parameters is greatly reduced by requiring the state to be completely symmetric. The dimension d drops out entirely, and the number of parameters becomes (one less than) the number of possible partitions of the n particles.

Motivation

Entanglement is a central issue in quantum information theory.

The study of n party-entanglement is considered difficult. It is complicated by the fact that the state space of n systems of size d has a large dimension: $d^{2n} - 1$.

The number of parameters is greatly reduced by requiring the state to be completely symmetric. The dimension d drops out entirely, and the number of parameters becomes (one less than) the number of possible partitions of the n particles.

For example, for 2 quantum identical systems of arbitrary size d there is only one parameter.

Motivation

Entanglement is a central issue in quantum information theory.

The study of n party-entanglement is considered difficult. It is complicated by the fact that the state space of n systems of size d has a large dimension: $d^{2^n} - 1$.

The number of parameters is greatly reduced by requiring the state to be completely symmetric. The dimension d drops out entirely, and the number of parameters becomes (one less than) the number of possible partitions of the n particles.

For example, for 2 quantum identical systems of arbitrary size d there is only one parameter.

An advantage of this restraint is that we can lean on a vast body of results from classical mathematics: the representation theory of S_n and $SU(d)$, as pioneered by Frobenius, Schur, Weyl, Littlewood,

Motivation

Entanglement is a central issue in quantum information theory.

The study of n party-entanglement is considered difficult. It is complicated by the fact that the state space of n systems of size d has a large dimension: $d^{2n} - 1$.

The number of parameters is greatly reduced by requiring the state to be completely symmetric. The dimension d drops out entirely, and the number of parameters becomes (one less than) the number of possible partitions of the n particles.

For example, for 2 quantum identical systems of arbitrary size d there is only one parameter.

An advantage of this restraint is that we can lean on a vast body of results from classical mathematics: the representation theory of S_n and $SU(d)$, as pioneered by Frobenius, Schur, Weyl, Littlewood,

But also some recent work in pure mathematics turns out to be surprisingly relevant to our question.

Overview of the talk

Overview of the talk

- ▶ Entanglement of Werner states for $n = 2$;

Overview of the talk

- ▶ Entanglement of Werner states for $n = 2$;
- ▶ Completely symmetric states on n particles;

Overview of the talk

- ▶ Entanglement of Werner states for $n = 2$;
- ▶ Completely symmetric states on n particles;
- ▶ Extremal separable completely symmetric states and immanants;

Overview of the talk

- ▶ Entanglement of Werner states for $n = 2$;
- ▶ Completely symmetric states on n particles;
- ▶ Extremal separable completely symmetric states and immanants;
- ▶ The case $n = 3$;

Overview of the talk

- ▶ Entanglement of Werner states for $n = 2$;
- ▶ Completely symmetric states on n particles;
- ▶ Extremal separable completely symmetric states and immanants;
- ▶ The case $n = 3$;
- ▶ The 'shadow' of the product states and its behaviour for general n ;

Overview of the talk

- ▶ Entanglement of Werner states for $n = 2$;
- ▶ Completely symmetric states on n particles;
- ▶ Extremal separable completely symmetric states and immanants;
- ▶ The case $n = 3$;
- ▶ The 'shadow' of the product states and its behaviour for general n ;
- ▶ Schur's inequality and Lieb's conjecture;

Overview of the talk

- ▶ Entanglement of Werner states for $n = 2$;
- ▶ Completely symmetric states on n particles;
- ▶ Extremal separable completely symmetric states and immanants;
- ▶ The case $n = 3$;
- ▶ The 'shadow' of the product states and its behaviour for general n ;
- ▶ Schur's inequality and Lieb's conjecture;
- ▶ The case $n = 4$;

Overview of the talk

- ▶ Entanglement of Werner states for $n = 2$;
- ▶ Completely symmetric states on n particles;
- ▶ Extremal separable completely symmetric states and immanants;
- ▶ The case $n = 3$;
- ▶ The 'shadow' of the product states and its behaviour for general n ;
- ▶ Schur's inequality and Lieb's conjecture;
- ▶ The case $n = 4$;
- ▶ The case $n = 5$: Hope crashed.

Overview of the talk

- ▶ Entanglement of Werner states for $n = 2$;
- ▶ Completely symmetric states on n particles;
- ▶ Extremal separable completely symmetric states and immanants;
- ▶ The case $n = 3$;
- ▶ The 'shadow' of the product states and its behaviour for general n ;
- ▶ Schur's inequality and Lieb's conjecture;
- ▶ The case $n = 4$;
- ▶ The case $n = 5$: Hope crashed.
- ▶ A measure of entanglement.

Symmetries in the two-particle case

Symmetries in the two-particle case

$$\mathcal{H} = \mathbb{C}^d \otimes \mathbb{C}^d .$$

Symmetries in the two-particle case

$$\mathcal{H} = \mathbb{C}^d \otimes \mathbb{C}^d .$$

Symmetry group: $u \in SU(d)$ acting as $u \otimes u$.

Symmetries in the two-particle case

$$\mathcal{H} = \mathbb{C}^d \otimes \mathbb{C}^d .$$

Symmetry group: $u \in SU(d)$ acting as $u \otimes u$.

This action commutes with the "Flip" operator:

$$F : \varphi \otimes \psi \mapsto \psi \otimes \varphi .$$

Symmetries in the two-particle case

$$\mathcal{H} = \mathbb{C}^d \otimes \mathbb{C}^d .$$

Symmetry group: $u \in SU(d)$ acting as $u \otimes u$.

This action commutes with the "Flip" operator:

$$F : \varphi \otimes \psi \mapsto \psi \otimes \varphi .$$

Eigenspaces \mathcal{H}_+ and \mathcal{H}_- of F are invariant for $SU(d)$:

Symmetries in the two-particle case

$$\mathcal{H} = \mathbb{C}^d \otimes \mathbb{C}^d .$$

Symmetry group: $u \in SU(d)$ acting as $u \otimes u$.

This action commutes with the "Flip" operator:

$$F : \varphi \otimes \psi \mapsto \psi \otimes \varphi .$$

Eigenspaces \mathcal{H}_+ and \mathcal{H}_- of F are invariant for $SU(d)$:

$$\mathcal{H}_+ := \text{span}\{\psi \otimes \psi \mid \psi \in \mathbb{C}^d\}$$

Symmetries in the two-particle case

$$\mathcal{H} = \mathbb{C}^d \otimes \mathbb{C}^d .$$

Symmetry group: $u \in SU(d)$ acting as $u \otimes u$.

This action commutes with the "Flip" operator:

$$F : \varphi \otimes \psi \mapsto \psi \otimes \varphi .$$

Eigenspaces \mathcal{H}_+ and \mathcal{H}_- of F are invariant for $SU(d)$:

$$\mathcal{H}_+ := \text{span}\{ \psi \otimes \psi \mid \psi \in \mathbb{C}^d \}$$

$$\text{basis: } \{ e_i \otimes e_i \mid 0 \leq i \leq d \} \cup \left\{ \frac{1}{\sqrt{2}}(e_i \otimes e_j + e_j \otimes e_i) \mid 0 \leq i < j \leq d \right\} .$$

Symmetries in the two-particle case

$$\mathcal{H} = \mathbb{C}^d \otimes \mathbb{C}^d .$$

Symmetry group: $u \in SU(d)$ acting as $u \otimes u$.

This action commutes with the "Flip" operator:

$$F : \varphi \otimes \psi \mapsto \psi \otimes \varphi .$$

Eigenspaces \mathcal{H}_+ and \mathcal{H}_- of F are invariant for $SU(d)$:

$$\mathcal{H}_+ := \text{span}\{\psi \otimes \psi \mid \psi \in \mathbb{C}^d\}$$

$$\text{basis: } \{e_i \otimes e_i \mid 0 \leq i \leq d\} \cup \left\{ \frac{1}{\sqrt{2}}(e_i \otimes e_j + e_j \otimes e_i) \mid 0 \leq i < j \leq d \right\} .$$

$$\dim \mathcal{H}_+ = \frac{d(d+1)}{2} = \binom{d+1}{2} ;$$

Symmetries in the two-particle case

$$\mathcal{H} = \mathbb{C}^d \otimes \mathbb{C}^d .$$

Symmetry group: $u \in SU(d)$ acting as $u \otimes u$.

This action commutes with the "Flip" operator:

$$F : \varphi \otimes \psi \mapsto \psi \otimes \varphi .$$

Eigenspaces \mathcal{H}_+ and \mathcal{H}_- of F are invariant for $SU(d)$:

$$\mathcal{H}_+ := \text{span}\{\psi \otimes \psi \mid \psi \in \mathbb{C}^d\}$$

$$\text{basis: } \{e_i \otimes e_i \mid 0 \leq i \leq d\} \cup \left\{ \frac{1}{\sqrt{2}}(e_i \otimes e_j + e_j \otimes e_i) \mid 0 \leq i < j \leq d \right\} .$$

$$\dim \mathcal{H}_+ = \frac{d(d+1)}{2} = \binom{d+1}{2} ;$$

$$\text{basis of } \mathcal{H}_- : \left\{ \frac{1}{\sqrt{2}}(e_i \otimes e_j - e_j \otimes e_i) \mid 0 \leq i < j \leq d \right\} .$$

Symmetries in the two-particle case

$$\mathcal{H} = \mathbb{C}^d \otimes \mathbb{C}^d .$$

Symmetry group: $u \in SU(d)$ acting as $u \otimes u$.

This action commutes with the "Flip" operator:

$$F : \varphi \otimes \psi \mapsto \psi \otimes \varphi .$$

Eigenspaces \mathcal{H}_+ and \mathcal{H}_- of F are invariant for $SU(d)$:

$$\mathcal{H}_+ := \text{span}\{\psi \otimes \psi \mid \psi \in \mathbb{C}^d\}$$

$$\text{basis: } \{e_i \otimes e_i \mid 0 \leq i \leq d\} \cup \left\{ \frac{1}{\sqrt{2}}(e_i \otimes e_j + e_j \otimes e_i) \mid 0 \leq i < j \leq d \right\} .$$

$$\dim \mathcal{H}_+ = \frac{d(d+1)}{2} = \binom{d+1}{2} ;$$

$$\text{basis of } \mathcal{H}_- : \left\{ \frac{1}{\sqrt{2}}(e_i \otimes e_j - e_j \otimes e_i) \mid 0 \leq i < j \leq d \right\} .$$

$$\dim \mathcal{H}_- = \frac{d(d-1)}{2} = \binom{d}{2} .$$

Werner states

Werner states

The group $SU(d)$ acts irreducibly on \mathcal{H}_+ and \mathcal{H}_- .

Werner states

The group $SU(d)$ acts irreducibly on \mathcal{H}_+ and \mathcal{H}_- .

Hence the action of $SU(d)$ has commutant

$$\{u \otimes u \mid u \in SU(d)\}' = \{\lambda p_+ + \mu p_- \mid \lambda, \mu \in \mathbb{C}\} = \{F\}'' ,$$

Werner states

The group $SU(d)$ acts irreducibly on \mathcal{H}_+ and \mathcal{H}_- .

Hence the action of $SU(d)$ has commutant

$$\{u \otimes u \mid u \in SU(d)\}' = \{\lambda p_+ + \mu p_- \mid \lambda, \mu \in \mathbb{C}\} = \{F\}'' ,$$

and the only symmetric projections are

$$p_{\pm} := \frac{\mathbb{1} \pm F}{2} = \text{projection onto } \mathcal{H}_{\pm} .$$

Werner states

The group $SU(d)$ acts irreducibly on \mathcal{H}_+ and \mathcal{H}_- .

Hence the action of $SU(d)$ has commutant

$$\{u \otimes u \mid u \in SU(d)\}' = \{\lambda p_+ + \mu p_- \mid \lambda, \mu \in \mathbb{C}\} = \{F\}'' ,$$

and the only symmetric projections are

$$p_{\pm} := \frac{\mathbb{1} \pm F}{2} = \text{projection onto } \mathcal{H}_{\pm} .$$

The only $SU(d)$ -symmetric states (i.e. **Werner states**) are convex combinations of

$$\rho_{\pm} := (\text{anti-})\text{symmetric state: } x \mapsto \frac{\text{tr} p_{\pm} x}{\text{tr} p_{\pm}} = \frac{\text{tr} \left(\frac{\mathbb{1} \pm F}{2} x \right)}{\dim \mathcal{H}_{\pm}} .$$

Werner states

The group $SU(d)$ acts irreducibly on \mathcal{H}_+ and \mathcal{H}_- .

Hence the action of $SU(d)$ has commutant

$$\{u \otimes u \mid u \in SU(d)\}' = \{\lambda p_+ + \mu p_- \mid \lambda, \mu \in \mathbb{C}\} = \{F\}'' ,$$

and the only symmetric projections are

$$p_{\pm} := \frac{\mathbb{1} \pm F}{2} = \text{projection onto } \mathcal{H}_{\pm} .$$

The only $SU(d)$ -symmetric states (i.e. **Werner states**) are convex combinations of

$$\rho_{\pm} := (\text{anti-})\text{symmetric state: } x \mapsto \frac{\text{tr} \rho_{\pm} x}{\text{tr} \rho_{\pm}} = \frac{\text{tr} \left(\frac{\mathbb{1} \pm F}{2} x \right)}{\dim \mathcal{H}_{\pm}} .$$

The Werner states are given by

$$\rho = \lambda \rho_+ + (1 - \lambda) \rho_- , \quad 0 \leq \lambda \leq 1 .$$

Entanglement of Werner states for $n = 2$

Entanglement of Werner states for $n = 2$

For $x \in M_d \otimes M_d$, let P_X denote its projection onto the center \mathcal{Z} :

$$P_X := \int_{SU(d)} (u \otimes u)^* x (u \otimes u) du ,$$

where du denotes the Haar measure on $SU(d)$.

Entanglement of Werner states for $n = 2$

For $x \in M_d \otimes M_d$, let P_X denote its projection onto the center \mathcal{Z} :

$$P_X := \int_{SU(d)} (u \otimes u)^* x (u \otimes u) du ,$$

where du denotes the Haar measure on $SU(d)$.

For states: $P^* \vartheta : x \mapsto \vartheta(P_X)$: projection of ϑ onto the Werner states,

Entanglement of Werner states for $n = 2$

For $x \in M_d \otimes M_d$, let P_X denote its projection onto the center \mathcal{Z} :

$$P_X := \int_{SU(d)} (u \otimes u)^* x (u \otimes u) du ,$$

where du denotes the Haar measure on $SU(d)$.

For states: $P^* \vartheta : x \mapsto \vartheta(P_X)$: projection of ϑ onto the Werner states, $P^* \vartheta$ is symmetric and coincides with ϑ on \mathcal{Z} .

Entanglement of Werner states for $n = 2$

For $x \in M_d \otimes M_d$, let P_X denote its projection onto the center \mathcal{Z} :

$$P_X := \int_{SU(d)} (u \otimes u)^* x (u \otimes u) du ,$$

where du denotes the Haar measure on $SU(d)$.

For states: $P^* \vartheta : x \mapsto \vartheta(P_X)$: projection of ϑ onto the Werner states, $P^* \vartheta$ is symmetric and coincides with ϑ on \mathcal{Z} .

Theorem

A Werner state ρ on $M_d \otimes M_d$ is separable iff

$$\rho(F) \geq 0 .$$

Entanglement of Werner states for $n = 2$

For $x \in M_d \otimes M_d$, let P_X denote its projection onto the center \mathcal{Z} :

$$P_X := \int_{SU(d)} (u \otimes u)^* x (u \otimes u) du ,$$

where du denotes the Haar measure on $SU(d)$.

For states: $P^* \vartheta : x \mapsto \vartheta(P_X)$: projection of ϑ onto the Werner states, $P^* \vartheta$ is symmetric and coincides with ϑ on \mathcal{Z} .

Theorem

A Werner state ρ on $M_d \otimes M_d$ is separable iff

$$\rho(F) \geq 0 .$$

This is a simple but optimal **Bell inequality**.

Entanglement of Werner states for $n = 2$

For $x \in M_d \otimes M_d$, let P_X denote its projection onto the center \mathcal{Z} :

$$P_X := \int_{SU(d)} (u \otimes u)^* x (u \otimes u) du ,$$

where du denotes the Haar measure on $SU(d)$.

For states: $P^* \vartheta : x \mapsto \vartheta(P_X)$: projection of ϑ onto the Werner states, $P^* \vartheta$ is symmetric and coincides with ϑ on \mathcal{Z} .

Theorem

A Werner state ρ on $M_d \otimes M_d$ is separable iff

$$\rho(F) \geq 0 .$$

This is a simple but optimal **Bell inequality**.

Entanglement of Werner states for $n = 2$

For $x \in M_d \otimes M_d$, let P_X denote its projection onto the center \mathcal{Z} :

$$P_X := \int_{SU(d)} (u \otimes u)^* x (u \otimes u) du ,$$

where du denotes the Haar measure on $SU(d)$.


For states: $P^* \vartheta : x \mapsto \vartheta(P_X)$: projection of ϑ onto the Werner states, $P^* \vartheta$ is symmetric and coincides with ϑ on \mathcal{Z} .

Theorem

A Werner state ρ on $M_d \otimes M_d$ is separable iff

$$\rho(F) \geq 0 .$$

This is a simple but optimal **Bell inequality**.



Entanglement of Werner states for $n = 2$

For $x \in M_d \otimes M_d$, let P_X denote its projection onto the center \mathcal{Z} :

$$P_X := \int_{SU(d)} (u \otimes u)^* x (u \otimes u) du ,$$

where du denotes the Haar measure on $SU(d)$.


For states: $P^* \vartheta : x \mapsto \vartheta(P_X)$: projection of ϑ onto the Werner states, $P^* \vartheta$ is symmetric and coincides with ϑ on \mathcal{Z} .

Theorem

A Werner state ρ on $M_d \otimes M_d$ is separable iff

$$\rho(F) \geq 0 .$$

This is a simple but optimal **Bell inequality**.



Entanglement of Werner states for $n = 2$

For $x \in M_d \otimes M_d$, let P_X denote its projection onto the center \mathcal{Z} :

$$P_X := \int_{SU(d)} (u \otimes u)^* x (u \otimes u) du ,$$

where du denotes the Haar measure on $SU(d)$.


For states: $P^* \vartheta : x \mapsto \vartheta(P_X)$: projection of ϑ onto the Werner states, $P^* \vartheta$ is symmetric and coincides with ϑ on \mathcal{Z} .

Theorem

A Werner state ρ on $M_d \otimes M_d$ is separable iff

$$\rho(F) \geq 0 .$$

This is a simple but optimal **Bell inequality**.



Proof.

For any pure product state $\psi \otimes \varphi \in \mathbb{C}^d \otimes \mathbb{C}^d$ the expectation of F is positive:



Proof.

For any pure product state $\psi \otimes \varphi \in \mathbb{C}^d \otimes \mathbb{C}^d$ the expectation of F is positive:

$$\langle \psi \otimes \varphi, F(\psi \otimes \varphi) \rangle = \langle \psi \otimes \varphi, \varphi \otimes \psi \rangle = \langle \psi, \varphi \rangle \langle \varphi, \psi \rangle = |\langle \psi, \varphi \rangle|^2 \geq 0.$$



Proof.

For any pure product state $\psi \otimes \varphi \in \mathbb{C}^d \otimes \mathbb{C}^d$ the expectation of F is positive:

$$\langle \psi \otimes \varphi, F(\psi \otimes \varphi) \rangle = \langle \psi \otimes \varphi, \varphi \otimes \psi \rangle = \langle \psi, \varphi \rangle \langle \varphi, \psi \rangle = |\langle \psi, \varphi \rangle|^2 \geq 0.$$

The inequality extends to all separable states by convexity.



Proof.

For any pure product state $\psi \otimes \varphi \in \mathbb{C}^d \otimes \mathbb{C}^d$ the expectation of F is positive:

$$\langle \psi \otimes \varphi, F(\psi \otimes \varphi) \rangle = \langle \psi \otimes \varphi, \varphi \otimes \psi \rangle = \langle \psi, \varphi \rangle \langle \varphi, \psi \rangle = |\langle \psi, \varphi \rangle|^2 \geq 0 .$$

The inequality extends to all separable states by convexity.

Conversely, suppose $0 \leq \rho(F) \leq 1$ for some Werner state ρ , and choose unit vectors ψ, φ with

$$|\langle \psi, \varphi \rangle|^2 = \rho(F) .$$



Proof.

For any pure product state $\psi \otimes \varphi \in \mathbb{C}^d \otimes \mathbb{C}^d$ the expectation of F is positive:

$$\langle \psi \otimes \varphi, F(\psi \otimes \varphi) \rangle = \langle \psi \otimes \varphi, \varphi \otimes \psi \rangle = \langle \psi, \varphi \rangle \langle \varphi, \psi \rangle = |\langle \psi, \varphi \rangle|^2 \geq 0.$$

The inequality extends to all separable states by convexity.

Conversely, suppose $0 \leq \rho(F) \leq 1$ for some Werner state ρ , and choose unit vectors ψ, φ with

$$|\langle \psi, \varphi \rangle|^2 = \rho(F).$$

Then the separable state

$$x \mapsto \langle \psi \otimes \varphi, P(x)\psi \otimes \varphi \rangle = \int_{SU(d)} \langle (u \otimes u)\psi \otimes \varphi, x(u \otimes u)\psi \otimes \varphi \rangle du$$

is a Werner state, and coincides with ρ on F .



Proof.

For any pure product state $\psi \otimes \varphi \in \mathbb{C}^d \otimes \mathbb{C}^d$ the expectation of F is positive:

$$\langle \psi \otimes \varphi, F(\psi \otimes \varphi) \rangle = \langle \psi \otimes \varphi, \varphi \otimes \psi \rangle = \langle \psi, \varphi \rangle \langle \varphi, \psi \rangle = |\langle \psi, \varphi \rangle|^2 \geq 0.$$

The inequality extends to all separable states by convexity.

Conversely, suppose $0 \leq \rho(F) \leq 1$ for some Werner state ρ , and choose unit vectors ψ, φ with

$$|\langle \psi, \varphi \rangle|^2 = \rho(F).$$

Then the separable state

$$x \mapsto \langle \psi \otimes \varphi, P(x)\psi \otimes \varphi \rangle = \int_{SU(d)} \langle (u \otimes u)\psi \otimes \varphi, x(u \otimes u)\psi \otimes \varphi \rangle du$$

is a Werner state, and coincides with ρ on F . Hence it equals ρ . □

The trace state moves to the boundary as $d \rightarrow \infty$

The trace state moves to the boundary as $d \rightarrow \infty$

Curious fact:

$$\lim_{d \rightarrow \infty} \tau_d \otimes \tau_d(F) = 0 .$$

The trace state moves to the boundary as $d \rightarrow \infty$

Curious fact:

$$\lim_{d \rightarrow \infty} \tau_d \otimes \tau_d(F) = 0 .$$

Indeed,

$$\tau_d \otimes \tau_d(F) = \frac{1}{d^2} \text{tr}_d \otimes \text{tr}_d(F)$$

The trace state moves to the boundary as $d \rightarrow \infty$

Curious fact:

$$\lim_{d \rightarrow \infty} \tau_d \otimes \tau_d(F) = 0.$$

Indeed,

$$\tau_d \otimes \tau_d(F) = \frac{1}{d^2} \text{tr}_d \otimes \text{tr}_d(F) = \frac{1}{d^2} \sum_{i,j=1}^d \langle e_i \otimes e_j, F(e_i \otimes e_j) \rangle$$

The trace state moves to the boundary as $d \rightarrow \infty$

Curious fact:

$$\lim_{d \rightarrow \infty} \tau_d \otimes \tau_d(F) = 0 .$$

Indeed,

$$\begin{aligned} \tau_d \otimes \tau_d(F) &= \frac{1}{d^2} \text{tr}_d \otimes \text{tr}_d(F) = \frac{1}{d^2} \sum_{i,j=1}^d \langle e_i \otimes e_j, F(e_i \otimes e_j) \rangle \\ &= \frac{1}{d^2} \sum_{i,j=1}^d \langle e_i \otimes e_j, e_j \otimes e_i \rangle = \frac{1}{d^2} \sum_{i,j=1}^d \delta_{ij} = \frac{1}{d} . \end{aligned}$$


The trace state moves to the boundary as $d \rightarrow \infty$

Curious fact:

$$\lim_{d \rightarrow \infty} \tau_d \otimes \tau_d(F) = 0.$$

Indeed,

$$\begin{aligned} \tau_d \otimes \tau_d(F) &= \frac{1}{d^2} \text{tr}_d \otimes \text{tr}_d(F) = \frac{1}{d^2} \sum_{i,j=1}^d \langle e_i \otimes e_j, F(e_i \otimes e_j) \rangle \\ &= \frac{1}{d^2} \sum_{i,j=1}^d \langle e_i \otimes e_j, e_j \otimes e_i \rangle = \frac{1}{d^2} \sum_{i,j=1}^d \delta_{ij} = \frac{1}{d}. \end{aligned}$$




The trace state moves to the boundary as $d \rightarrow \infty$

Curious fact:

$$\lim_{d \rightarrow \infty} \tau_d \otimes \tau_d(F) = 0.$$

Indeed,

$$\begin{aligned} \tau_d \otimes \tau_d(F) &= \frac{1}{d^2} \text{tr}_d \otimes \text{tr}_d(F) = \frac{1}{d^2} \sum_{i,j=1}^d \langle e_i \otimes e_j, F(e_i \otimes e_j) \rangle \\ &= \frac{1}{d^2} \sum_{i,j=1}^d \langle e_i \otimes e_j, e_j \otimes e_i \rangle = \frac{1}{d^2} \sum_{i,j=1}^d \delta_{ij} = \frac{1}{d}. \end{aligned}$$




The trace state moves to the boundary as $d \rightarrow \infty$

Curious fact:

$$\lim_{d \rightarrow \infty} \tau_d \otimes \tau_d(F) = 0.$$

Indeed,

$$\begin{aligned} \tau_d \otimes \tau_d(F) &= \frac{1}{d^2} \text{tr}_d \otimes \text{tr}_d(F) = \frac{1}{d^2} \sum_{i,j=1}^d \langle e_i \otimes e_j, F(e_i \otimes e_j) \rangle \\ &= \frac{1}{d^2} \sum_{i,j=1}^d \langle e_i \otimes e_j, e_j \otimes e_i \rangle = \frac{1}{d^2} \sum_{i,j=1}^d \delta_{ij} = \frac{1}{d}. \end{aligned}$$




The trace state moves to the boundary as $d \rightarrow \infty$

Curious fact:

$$\lim_{d \rightarrow \infty} \tau_d \otimes \tau_d(F) = 0.$$

Indeed,

$$\begin{aligned} \tau_d \otimes \tau_d(F) &= \frac{1}{d^2} \text{tr}_d \otimes \text{tr}_d(F) = \frac{1}{d^2} \sum_{i,j=1}^d \langle e_i \otimes e_j, F(e_i \otimes e_j) \rangle \\ &= \frac{1}{d^2} \sum_{i,j=1}^d \langle e_i \otimes e_j, e_j \otimes e_i \rangle = \frac{1}{d^2} \sum_{i,j=1}^d \delta_{ij} = \frac{1}{d}. \end{aligned}$$



The trace state moves to the boundary as $d \rightarrow \infty$

Curious fact:

$$\lim_{d \rightarrow \infty} \tau_d \otimes \tau_d(F) = 0.$$

Indeed,

$$\begin{aligned} \tau_d \otimes \tau_d(F) &= \frac{1}{d^2} \text{tr}_d \otimes \text{tr}_d(F) = \frac{1}{d^2} \sum_{i,j=1}^d \langle e_i \otimes e_j, F(e_i \otimes e_j) \rangle \\ &= \frac{1}{d^2} \sum_{i,j=1}^d \langle e_i \otimes e_j, e_j \otimes e_i \rangle = \frac{1}{d^2} \sum_{i,j=1}^d \delta_{ij} = \frac{1}{d}. \end{aligned}$$



General $n \in \mathbb{N}$: Schur-Weyl duality

General $n \in \mathbb{N}$: Schur-Weyl duality

On the Hilbert space

$$\mathcal{H} := \mathbb{C}^d \otimes \mathbb{C}^d \otimes \dots \otimes \mathbb{C}^d \quad (n \text{ times})$$

General $n \in \mathbb{N}$: Schur-Weyl duality

On the Hilbert space

$$\mathcal{H} := \mathbb{C}^d \otimes \mathbb{C}^d \otimes \dots \otimes \mathbb{C}^d \quad (n \text{ times})$$

there are representations of **two groups**: S_n and $SU(d)$:

General $n \in \mathbb{N}$: Schur-Weyl duality

On the Hilbert space

$$\mathcal{H} := \mathbb{C}^d \otimes \mathbb{C}^d \otimes \dots \otimes \mathbb{C}^d \quad (n \text{ times})$$

there are representations of **two groups**: S_n and $SU(d)$:

$$S_n \ni \sigma \quad : \quad \pi(\sigma)\psi_1 \otimes \psi_2 \otimes \dots \otimes \psi_n := \psi_{\sigma^{-1}(1)} \otimes \psi_{\sigma^{-1}(2)} \otimes \dots \otimes \psi_{\sigma^{-1}(n)}$$

General $n \in \mathbb{N}$: Schur-Weyl duality

On the Hilbert space

$$\mathcal{H} := \mathbb{C}^d \otimes \mathbb{C}^d \otimes \dots \otimes \mathbb{C}^d \quad (n \text{ times})$$

there are representations of **two groups**: S_n and $SU(d)$:

$$S_n \ni \sigma \quad : \quad \pi(\sigma)\psi_1 \otimes \psi_2 \otimes \dots \otimes \psi_n := \psi_{\sigma^{-1}(1)} \otimes \psi_{\sigma^{-1}(2)} \otimes \dots \otimes \psi_{\sigma^{-1}(n)}$$

$$SU(d) \ni u \quad : \quad \pi'(u)\psi_1 \otimes \psi_2 \otimes \dots \otimes \psi_n := u\psi_1 \otimes u\psi_2 \otimes \dots \otimes u\psi_n$$

General $n \in \mathbb{N}$: Schur-Weyl duality

On the Hilbert space

$$\mathcal{H} := \mathbb{C}^d \otimes \mathbb{C}^d \otimes \dots \otimes \mathbb{C}^d \quad (n \text{ times})$$

there are representations of **two groups**: S_n and $SU(d)$:

$$S_n \ni \sigma \quad : \quad \pi(\sigma)\psi_1 \otimes \psi_2 \otimes \dots \otimes \psi_n := \psi_{\sigma^{-1}(1)} \otimes \psi_{\sigma^{-1}(2)} \otimes \dots \otimes \psi_{\sigma^{-1}(n)}$$

$$SU(d) \ni u \quad : \quad \pi'(u)\psi_1 \otimes \psi_2 \otimes \dots \otimes \psi_n := u\psi_1 \otimes u\psi_2 \otimes \dots \otimes u\psi_n$$

The classical **Schur-Weyl duality theorem** states that these two group actions do not only commute, but the algebras they generate are actually **each other's commutant**.

General $n \in \mathbb{N}$: Schur-Weyl duality

On the Hilbert space

$$\mathcal{H} := \mathbb{C}^d \otimes \mathbb{C}^d \otimes \dots \otimes \mathbb{C}^d \quad (n \text{ times})$$

there are representations of **two groups**: S_n and $SU(d)$:

$$S_n \ni \sigma \quad : \quad \pi(\sigma)\psi_1 \otimes \psi_2 \otimes \dots \otimes \psi_n := \psi_{\sigma^{-1}(1)} \otimes \psi_{\sigma^{-1}(2)} \otimes \dots \otimes \psi_{\sigma^{-1}(n)}$$

$$SU(d) \ni u \quad : \quad \pi'(u)\psi_1 \otimes \psi_2 \otimes \dots \otimes \psi_n := u\psi_1 \otimes u\psi_2 \otimes \dots \otimes u\psi_n$$

The classical **Schur-Weyl duality theorem** states that these two group actions do not only commute, but the algebras they generate are actually **each other's commutant**. In particular they have **the same center**:

$$\mathcal{Z} := \mathcal{Z}(n, d) := \pi(S_n)' \cap \pi'(SU(d))' .$$

General $n \in \mathbb{N}$: Schur-Weyl duality

On the Hilbert space

$$\mathcal{H} := \mathbb{C}^d \otimes \mathbb{C}^d \otimes \dots \otimes \mathbb{C}^d \quad (n \text{ times})$$

there are representations of **two groups**: S_n and $SU(d)$:

$$S_n \ni \sigma \quad : \quad \pi(\sigma)\psi_1 \otimes \psi_2 \otimes \dots \otimes \psi_n := \psi_{\sigma^{-1}(1)} \otimes \psi_{\sigma^{-1}(2)} \otimes \dots \otimes \psi_{\sigma^{-1}(n)}$$

$$SU(d) \ni u \quad : \quad \pi'(u)\psi_1 \otimes \psi_2 \otimes \dots \otimes \psi_n := u\psi_1 \otimes u\psi_2 \otimes \dots \otimes u\psi_n$$

The classical **Schur-Weyl duality theorem** states that these two group actions do not only commute, but the algebras they generate are actually **each other's commutant**. In particular they have **the same center**:

$$\mathcal{Z} := \mathcal{Z}(n, d) := \pi(S_n)' \cap \pi'(SU(d))' .$$

The minimal projections in this center cut **both group representations** into their irreducible components, and they are labeled by **Young diagrams**.

General $n \in \mathbb{N}$: Schur-Weyl duality

On the Hilbert space

$$\mathcal{H} := \mathbb{C}^d \otimes \mathbb{C}^d \otimes \dots \otimes \mathbb{C}^d \quad (n \text{ times})$$

there are representations of **two groups**: S_n and $SU(d)$:

$$S_n \ni \sigma \quad : \quad \pi(\sigma)\psi_1 \otimes \psi_2 \otimes \dots \otimes \psi_n := \psi_{\sigma^{-1}(1)} \otimes \psi_{\sigma^{-1}(2)} \otimes \dots \otimes \psi_{\sigma^{-1}(n)}$$

$$SU(d) \ni u \quad : \quad \pi'(u)\psi_1 \otimes \psi_2 \otimes \dots \otimes \psi_n := u\psi_1 \otimes u\psi_2 \otimes \dots \otimes u\psi_n$$

The classical **Schur-Weyl duality theorem** states that these two group actions do not only commute, but the algebras they generate are actually **each other's commutant**. In particular they have **the same center**:

$$\mathcal{Z} := \mathcal{Z}(n, d) := \pi(S_n)' \cap \pi'(SU(d))' .$$

The minimal projections in this center cut **both group representations** into their irreducible components, and they are labeled by **Young diagrams**.

For example

$$\begin{aligned} 3 \otimes 3 \otimes 3 &= (10 \otimes 1_+) \oplus (8 \otimes 2) \oplus (1 \otimes 1_-) . \\ &= \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array} \oplus \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} \oplus \begin{array}{|c|} \hline \square \\ \hline \end{array} . \end{aligned}$$

The group algebra of S_n

The group algebra of S_n

Let \mathcal{A}_n denote the **group algebra** of S_n :

The group algebra of S_n

Let \mathcal{A}_n denote the **group algebra** of S_n :

$$f : S_n \rightarrow \mathbb{C} \quad \text{to be viewed as} \quad \sum_{\sigma \in S_n} f(\sigma)\sigma .$$

The group algebra of S_n

Let \mathcal{A}_n denote the **group algebra** of S_n :

$$f : S_n \rightarrow \mathbb{C} \quad \text{to be viewed as} \quad \sum_{\sigma \in S_n} f(\sigma)\sigma .$$

Multiplication in \mathcal{A}_n is convolution:

$$(f * g)(\sigma) = \sum_{\tau \in S_n} f(\tau)g(\tau^{-1}\sigma) .$$

The group algebra of S_n

Let \mathcal{A}_n denote the **group algebra** of S_n :

$$f : S_n \rightarrow \mathbb{C} \quad \text{to be viewed as} \quad \sum_{\sigma \in S_n} f(\sigma)\sigma .$$

Multiplication in \mathcal{A}_n is convolution:

$$(f * g)(\sigma) = \sum_{\tau \in S_n} f(\tau)g(\tau^{-1}\sigma) .$$

The unit is δ_e , where e is the identity element of S_n .

The group algebra of S_n

Let \mathcal{A}_n denote the **group algebra** of S_n :

$$f : S_n \rightarrow \mathbb{C} \quad \text{to be viewed as} \quad \sum_{\sigma \in S_n} f(\sigma)\sigma .$$

Multiplication in \mathcal{A}_n is convolution:

$$(f * g)(\sigma) = \sum_{\tau \in S_n} f(\tau)g(\tau^{-1}\sigma) .$$

The unit is δ_e , where e is the identity element of S_n .

Adjoint operation:

$$f^*(\sigma) = \overline{f(\sigma^{-1})} .$$

The group algebra of S_n

Let \mathcal{A}_n denote the **group algebra** of S_n :

$$f : S_n \rightarrow \mathbb{C} \quad \text{to be viewed as} \quad \sum_{\sigma \in S_n} f(\sigma)\sigma .$$

Multiplication in \mathcal{A}_n is convolution:

$$(f * g)(\sigma) = \sum_{\tau \in S_n} f(\tau)g(\tau^{-1}\sigma) .$$

The unit is δ_e , where e is the identity element of S_n .

Adjoint operation:

$$f^*(\sigma) = \overline{f(\sigma^{-1})} .$$

Every unitary representation of S_n automatically extends to a representation of \mathcal{A}_n .

The group algebra of S_n

Let \mathcal{A}_n denote the **group algebra** of S_n :

$$f : S_n \rightarrow \mathbb{C} \quad \text{to be viewed as} \quad \sum_{\sigma \in S_n} f(\sigma)\sigma .$$

Multiplication in \mathcal{A}_n is convolution:

$$(f * g)(\sigma) = \sum_{\tau \in S_n} f(\tau)g(\tau^{-1}\sigma) .$$

The unit is δ_e , where e is the identity element of S_n .

Adjoint operation:

$$f^*(\sigma) = \overline{f(\sigma^{-1})} .$$

Every unitary representation of S_n automatically extends to a representation of \mathcal{A}_n .

In our case

$$\pi(f) : \quad \psi_1 \otimes \dots \otimes \psi_n \mapsto \sum_{\sigma \in S_n} f(\sigma)\psi_{\sigma^{-1}(1)} \otimes \dots \otimes \psi_{\sigma^{-1}n} .$$

The (left) regular representation of S_n

The (left) regular representation of S_n

We let $f \in \mathcal{A}_n$ act on the Hilbert space $l^2(S_n)$ by convolution on the left:

$$h \mapsto f * h .$$

The (left) regular representation of S_n

We let $f \in \mathcal{A}_n$ act on the Hilbert space $l^2(S_n)$ by convolution on the left:

$$h \mapsto f * h .$$

The trace on this "Hilbert space" is of a particularly simple form:

$$\mathrm{tr}_{\mathrm{reg}}(f) := \sum_{\sigma \in S_n} \langle \delta_\sigma, f * \delta_\sigma \rangle = \sum_{\sigma \in S_n} (f * \delta_\sigma)(\sigma) = n! \cdot f(e) ,$$

and will be called the **regular trace**.

The (left) regular representation of S_n

We let $f \in \mathcal{A}_n$ act on the Hilbert space $l^2(S_n)$ by convolution on the left:

$$h \mapsto f * h .$$

The trace on this "Hilbert space" is of a particularly simple form:

$$\mathrm{tr}_{\mathrm{reg}}(f) := \sum_{\sigma \in S_n} \langle \delta_\sigma, f * \delta_\sigma \rangle = \sum_{\sigma \in S_n} (f * \delta_\sigma)(\sigma) = n! \cdot f(e) ,$$

and will be called the **regular trace**.

The normalized version $\tau_{\mathrm{reg}} := \frac{1}{n!} \mathrm{tr}_{\mathrm{reg}}$ is the **regular trace state**.

The center of the group algebra of S_n

The center of the group algebra of S_n

$$\mathcal{Z}_n := \mathcal{A}_n \cap \mathcal{A}'_n .$$

The center of the group algebra of S_n

$$\mathcal{Z}_n := \mathcal{A}_n \cap \mathcal{A}'_n .$$

We have $f \in \mathcal{Z}_n$ if and only if for all $\sigma, \tau \in S_n$: $f(\sigma\tau) = f(\tau\sigma)$:

The center of the group algebra of S_n

$$\mathcal{Z}_n := \mathcal{A}_n \cap \mathcal{A}'_n .$$

We have $f \in \mathcal{Z}_n$ if and only if for all $\sigma, \tau \in S_n$: $f(\sigma\tau) = f(\tau\sigma)$:
The center consists of the **class functions**.

The center of the group algebra of S_n

$$\mathcal{Z}_n := \mathcal{A}_n \cap \mathcal{A}'_n .$$

We have $f \in \mathcal{Z}_n$ if and only if for all $\sigma, \tau \in S_n$: $f(\sigma\tau) = f(\tau\sigma)$:
The center consists of the **class functions**. Hence

The center of the group algebra of S_n

$$\mathcal{Z}_n := \mathcal{A}_n \cap \mathcal{A}'_n .$$

We have $f \in \mathcal{Z}_n$ if and only if for all $\sigma, \tau \in S_n$: $f(\sigma\tau) = f(\tau\sigma)$:
The center consists of the **class functions**. Hence

$$\dim \mathcal{Z}_n = \#(\text{conjugacy classes of } S_n)$$

The center of the group algebra of S_n

$$\mathcal{Z}_n := \mathcal{A}_n \cap \mathcal{A}'_n .$$

We have $f \in \mathcal{Z}_n$ if and only if for all $\sigma, \tau \in S_n$: $f(\sigma\tau) = f(\tau\sigma)$:
The center consists of the **class functions**. Hence

$$\begin{aligned} \dim \mathcal{Z}_n &= \#(\text{conjugacy classes of } S_n) \\ &= \#(\text{partitions of } n) =: \mathcal{P}(n) . \end{aligned}$$

The center of the group algebra of S_n

$$\mathcal{Z}_n := \mathcal{A}_n \cap \mathcal{A}'_n .$$

We have $f \in \mathcal{Z}_n$ if and only if for all $\sigma, \tau \in S_n$: $f(\sigma\tau) = f(\tau\sigma)$:
The center consists of the **class functions**. Hence

$$\begin{aligned} \dim \mathcal{Z}_n &= \#(\text{conjugacy classes of } S_n) \\ &= \#(\text{partitions of } n) =: \mathcal{P}(n) . \end{aligned}$$

On the other hand, since \mathcal{Z}_n is an abelian matrix algebra, it must be of the form

$$\mathcal{Z}_n = \bigoplus_{i=1}^{\mathcal{P}(n)} \mathbb{C} p_i$$

for some orthogonal set of minimal projections p_i in the center.

The center of the group algebra of S_n

$$\mathcal{Z}_n := \mathcal{A}_n \cap \mathcal{A}'_n .$$

We have $f \in \mathcal{Z}_n$ if and only if for all $\sigma, \tau \in S_n$: $f(\sigma\tau) = f(\tau\sigma)$:
The center consists of the **class functions**. Hence

$$\begin{aligned} \dim \mathcal{Z}_n &= \#(\text{conjugacy classes of } S_n) \\ &= \#(\text{partitions of } n) =: \mathcal{P}(n) . \end{aligned}$$

On the other hand, since \mathcal{Z}_n is an abelian matrix algebra, it must be of the form

$$\mathcal{Z}_n = \bigoplus_{i=1}^{\mathcal{P}(n)} \mathbb{C} p_i$$

for some orthogonal set of minimal projections p_i in the center.

The states on the center form a simplex with extreme points ρ_i given by

$$\rho_i(p_j) = \delta_{ij} .$$

minimal projections, characters, and irreducible representations

minimal projections, characters, and irreducible representations

The center \mathcal{Z}_n of the group algebra \mathcal{A}_n has dimension $\mathcal{P}(n)$.

minimal projections, characters, and irreducible representations

The center \mathcal{Z}_n of the group algebra \mathcal{A}_n has dimension $\mathcal{P}(n)$.

So it contains $\mathcal{P}(n)$ **minimal projections** $p_1, p_2, \dots, p_{\mathcal{P}(n)}$:

$$p_i(\sigma^{-1}) = \overline{p_i(\sigma)}, \quad p_i * p_j = \delta_{ij} p_i \quad \text{and} \quad \sum_{i=1}^{\mathcal{P}(n)} p_i = \delta_e .$$

minimal projections, characters, and irreducible representations

The center \mathcal{Z}_n of the group algebra \mathcal{A}_n has dimension $\mathcal{P}(n)$.

So it contains $\mathcal{P}(n)$ **minimal projections** $p_1, p_2, \dots, p_{\mathcal{P}(n)}$:

$$p_i(\sigma^{-1}) = \overline{p_i(\sigma)}, \quad p_i * p_j = \delta_{ij} p_i \quad \text{and} \quad \sum_{i=1}^{\mathcal{P}(n)} p_i = \delta_e .$$

They cut the algebra $\mathcal{A} = \mathcal{A}_n$ into factors $p_i \mathcal{A}$:

$$\mathcal{A} = \bigoplus_{i=1}^{\mathcal{P}(n)} p_i \mathcal{A} \quad \simeq \quad \bigoplus_{i=1}^{\mathcal{P}(n)} M_{d(i)} .$$

minimal projections, characters, and irreducible representations

The center \mathcal{Z}_n of the group algebra \mathcal{A}_n has dimension $\mathcal{P}(n)$.

So it contains $\mathcal{P}(n)$ **minimal projections** $p_1, p_2, \dots, p_{\mathcal{P}(n)}$:

$$p_i(\sigma^{-1}) = \overline{p_i(\sigma)}, \quad p_i * p_j = \delta_{ij} p_i \quad \text{and} \quad \sum_{i=1}^{\mathcal{P}(n)} p_i = \delta_e .$$

They cut the algebra $\mathcal{A} = \mathcal{A}_n$ into factors $p_i \mathcal{A}$:

$$\mathcal{A} = \bigoplus_{i=1}^{\mathcal{P}(n)} p_i \mathcal{A} \quad \simeq \quad \bigoplus_{i=1}^{\mathcal{P}(n)} M_{d(i)} .$$

Hence

$$d(i)^2 = \operatorname{tr}(p_i) = n! \cdot p_i(e) .$$

minimal projections, characters, and irreducible representations

The center \mathcal{Z}_n of the group algebra \mathcal{A}_n has dimension $\mathcal{P}(n)$.

So it contains $\mathcal{P}(n)$ **minimal projections** $p_1, p_2, \dots, p_{\mathcal{P}(n)}$:

$$p_i(\sigma^{-1}) = \overline{p_i(\sigma)}, \quad p_i * p_j = \delta_{ij} p_i \quad \text{and} \quad \sum_{i=1}^{\mathcal{P}(n)} p_i = \delta_e .$$

They cut the algebra $\mathcal{A} = \mathcal{A}_n$ into factors $p_i \mathcal{A}$:

$$\mathcal{A} = \bigoplus_{i=1}^{\mathcal{P}(n)} p_i \mathcal{A} \quad \simeq \quad \bigoplus_{i=1}^{\mathcal{P}(n)} M_{d(i)} .$$

Hence

$$d(i)^2 = \text{tr}(p_i) = n! \cdot p_i(e) .$$

Now define the **character** $\chi_i : \mathcal{S}_n \rightarrow \mathbb{C}$ by:

$$\chi_i(\sigma) := \frac{n!}{d(i)} p_i(\sigma) .$$

minimal projections, characters, and irreducible representations

The center \mathcal{Z}_n of the group algebra \mathcal{A}_n has dimension $\mathcal{P}(n)$.

So it contains $\mathcal{P}(n)$ **minimal projections** $p_1, p_2, \dots, p_{\mathcal{P}(n)}$:

$$p_i(\sigma^{-1}) = \overline{p_i(\sigma)}, \quad p_i * p_j = \delta_{ij} p_i \quad \text{and} \quad \sum_{i=1}^{\mathcal{P}(n)} p_i = \delta_e .$$

They cut the algebra $\mathcal{A} = \mathcal{A}_n$ into factors $p_i \mathcal{A}$:

$$\mathcal{A} = \bigoplus_{i=1}^{\mathcal{P}(n)} p_i \mathcal{A} \quad \simeq \quad \bigoplus_{i=1}^{\mathcal{P}(n)} M_{d(i)} .$$

Hence

$$d(i)^2 = \text{tr}(p_i) = n! \cdot p_i(e) .$$

Now define the **character** $\chi_i : \mathcal{S}_n \rightarrow \mathbb{C}$ by:

$$\chi_i(\sigma) := \frac{n!}{d(i)} p_i(\sigma) .$$

Then these functions form an orthonormal set in the sense that

$$\langle \chi_i, \chi_j \rangle = \sum_{\sigma} \overline{\chi_i(\sigma)} \chi_j(\sigma) = \frac{(n!)^2}{d(i)d(j)} p_i * p_j(e) = \frac{n!}{d(i)^2} \cdot n! p_i(e) \delta_{ij} = n! \cdot \delta_{ij} .$$

Interpretation of the characters χ_i

Interpretation of the characters χ_i

Lemma

The character $\chi_i(\sigma)$ is the trace of σ in its irreducible representation labelled by i .

Interpretation of the characters χ_i

Lemma

The character $\chi_i(\sigma)$ is the trace of σ in its irreducible representation labelled by i .

Proof.

Interpretation of the characters χ_i

Lemma

The character $\chi_i(\sigma)$ is the trace of σ in its irreducible representation labelled by i .

Proof.

The space $p_i \mathcal{H}_{\text{reg}}$ looks like $M_{d(i)} \simeq \mathbb{C}^{d(i)} \otimes \mathbb{C}^{d(i)}$.

Interpretation of the characters χ_i

Lemma

The character $\chi_i(\sigma)$ is the trace of σ in its irreducible representation labelled by i .

Proof.

The space $p_i \mathcal{H}_{\text{reg}}$ looks like $M_{d(i)} \simeq \mathbb{C}^{d(i)} \otimes \mathbb{C}^{d(i)}$.

On this space \mathcal{A} and its commutant \mathcal{A}' act as

$$p_i \mathcal{A} \simeq M_{d(i)} \otimes \mathbb{1} \quad \text{and} \quad p_i (\mathcal{A}') \simeq \mathbb{1} \otimes M_{d(i)} .$$

Interpretation of the characters χ_i

Lemma

The character $\chi_i(\sigma)$ is the trace of σ in its irreducible representation labelled by i .

Proof.

The space $p_i \mathcal{H}_{\text{reg}}$ looks like $M_{d(i)} \simeq \mathbb{C}^{d(i)} \otimes \mathbb{C}^{d(i)}$.

On this space \mathcal{A} and its commutant \mathcal{A}' act as

$$p_i \mathcal{A} \simeq M_{d(i)} \otimes \mathbb{1} \quad \text{and} \quad p_i(\mathcal{A}') \simeq \mathbb{1} \otimes M_{d(i)} .$$

Let us now choose a minimal projection $q_i \in p_i(\mathcal{A}')$.

Interpretation of the characters χ_i

Lemma

The character $\chi_i(\sigma)$ is the trace of σ in its irreducible representation labelled by i .

Proof.

The space $p_i \mathcal{H}_{\text{reg}}$ looks like $M_{d(i)} \simeq \mathbb{C}^{d(i)} \otimes \mathbb{C}^{d(i)}$.

On this space \mathcal{A} and its commutant \mathcal{A}' act as

$$p_i \mathcal{A} \simeq M_{d(i)} \otimes \mathbb{1} \quad \text{and} \quad p_i(\mathcal{A}') \simeq \mathbb{1} \otimes M_{d(i)} .$$

Let us now choose a minimal projection $q_i \in p_i(\mathcal{A}')$.

Then the regular representation acts irreducibly on the range of $p_i q_i$.

Interpretation of the characters χ_i

Lemma

The character $\chi_i(\sigma)$ is the trace of σ in its irreducible representation labelled by i .

Proof.

The space $p_i \mathcal{H}_{\text{reg}}$ looks like $M_{d(i)} \simeq \mathbb{C}^{d(i)} \otimes \mathbb{C}^{d(i)}$.

On this space \mathcal{A} and its commutant \mathcal{A}' act as

$$p_i \mathcal{A} \simeq M_{d(i)} \otimes \mathbb{1} \quad \text{and} \quad p_i (\mathcal{A}') \simeq \mathbb{1} \otimes M_{d(i)} .$$

Let us now choose a minimal projection $q_i \in p_i (\mathcal{A}')$.

Then the regular representation acts irreducibly on the range of $p_i q_i$.

Now, in the regular representation we may calculate

$$\chi_i(\sigma) = \frac{n!}{d(i)} p_i(\sigma) = \frac{1}{d(i)} n! (p_i * \delta_\sigma)(e) = \frac{1}{d(i)} \text{tr}_{\text{reg}}(p_i * \delta_\sigma) = \text{tr}_{\text{reg}}(p_i * q_i * \delta_\sigma) ,$$



Young frames

Young frames

The irreducible representations of S_n (and hence also the minimal central projections and the characters) are labelled by **Young frames** with n boxes:

$$Y = \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \\ \hline \square & \square & \square & \\ \hline \square & & & \\ \hline \end{array} .$$

Young frames

The irreducible representations of S_n (and hence also the minimal central projections and the characters) are labelled by **Young frames** with n boxes:

$$Y = \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \\ \hline \square & \square & \square & \\ \hline \square & & & \\ \hline \end{array} .$$

(Hook length rule)

$$d(Y) = \frac{n!}{\prod \text{hook lengths}} .$$

Young frames

The irreducible representations of S_n (and hence also the minimal central projections and the characters) are labelled by **Young frames** with n boxes:

$$Y = \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \square & \square & \square & \\ \hline \square & \square & \square & \\ \hline \square & & & \\ \hline \end{array} .$$

(Hook length rule)

$$d(Y) = \frac{n!}{\prod \text{hook lengths}} .$$

For example:

$$d \left(\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \square & \square & \\ \hline \end{array} \right) = \frac{5!}{4 \times 3 \times 2} = 5 \quad \text{hook lengths: } \begin{array}{|c|c|c|} \hline 4 & 3 & 1 \\ \hline 2 & 1 & \\ \hline \end{array} .$$

An exclusion principle

An exclusion principle

Theorem

Let $n, d \in \mathbb{N}$. Let Y denote a Young frame with n boxes and height $h(Y)$.

Then

$$\pi_{n,d}(p_Y) = 0 \quad \text{iff} \quad h(Y) > d .$$

An exclusion principle

Theorem

Let $n, d \in \mathbb{N}$. Let Y denote a Young frame with n boxes and height $h(Y)$.
Then

$$\pi_{n,d}(p_Y) = 0 \quad \text{iff} \quad h(Y) > d .$$

Proof.

This can be shown using the explicit representation of p_Y ; see, for example, B. Simon: '*Representations of finite and compact groups*'. □

An exclusion principle

Theorem

Let $n, d \in \mathbb{N}$. Let Y denote a Young frame with n boxes and height $h(Y)$.
Then

$$\pi_{n,d}(p_Y) = 0 \quad \text{iff} \quad h(Y) > d .$$

Proof.

This can be shown using the explicit representation of p_Y ; see, for example, B. Simon: *'Representations of finite and compact groups'*. □

For example, the symmetric subspace, having Young frame $\square\square\square\square$, is nonzero in $(\mathbb{C}^d)^{\otimes 4}$ for every one-particle dimension d , but, according to **Pauli's exclusion principle**, the antisymmetric subspace, with Young frame $\begin{array}{c} \square \\ \square \\ \square \\ \square \end{array}$, needs $d \geq 4$.

An exclusion principle

Theorem

Let $n, d \in \mathbb{N}$. Let Y denote a Young frame with n boxes and height $h(Y)$.
Then

$$\pi_{n,d}(p_Y) = 0 \quad \text{iff} \quad h(Y) > d .$$

Proof.

This can be shown using the explicit representation of p_Y ; see, for example, B. Simon: *'Representations of finite and compact groups'*. □

For example, the symmetric subspace, having Young frame $\square\square\square\square$, is nonzero in $(\mathbb{C}^d)^{\otimes 4}$ for every one-particle dimension d , but, according to **Pauli's exclusion**

principle, the antisymmetric subspace, with Young frame $\begin{array}{c} \square \\ \square \\ \square \end{array}$, needs $d \geq 4$.

Hence the above theorem generalizes this exclusion principle.

Completely symmetric states on $\mathcal{B}((\mathbb{C}^d)^{\otimes n})$

Completely symmetric states on $\mathcal{B}((\mathbb{C}^d)^{\otimes n})$

Observables (operators) on $\mathcal{H} := \mathbb{C}^d \otimes \dots \otimes \mathbb{C}^d$ can be 'twirled' and averaged:

$$Ta := \int_{SU(d)} (u \otimes \dots \otimes u)^* a (u \otimes \dots \otimes u) du ;$$

$$Ma := \frac{1}{n!} \sum_{\sigma \in S_n} \pi(\sigma) a \pi(\sigma) .$$

Completely symmetric states on $\mathcal{B}((\mathbb{C}^d)^{\otimes n})$

Observables (operators) on $\mathcal{H} := \mathbb{C}^d \otimes \dots \otimes \mathbb{C}^d$ can be 'twirled' and averaged:

$$Ta := \int_{SU(d)} (u \otimes \dots \otimes u)^* a (u \otimes \dots \otimes u) du ;$$
$$Ma := \frac{1}{n!} \sum_{\sigma \in S_n} \pi(\sigma) a \pi(\sigma) .$$

Clearly, $Ta \in \pi'(SU(d))'$, and in the same way $Ma \in \pi(S_n)'$.

Completely symmetric states on $\mathcal{B}((\mathbb{C}^d)^{\otimes n})$

Observables (operators) on $\mathcal{H} := \mathbb{C}^d \otimes \dots \otimes \mathbb{C}^d$ can be 'twirled' and averaged:

$$Ta := \int_{SU(d)} (u \otimes \dots \otimes u)^* a (u \otimes \dots \otimes u) du ;$$

$$Ma := \frac{1}{n!} \sum_{\sigma \in S_n} \pi(\sigma) a \pi(\sigma) .$$

Clearly, $Ta \in \pi'(SU(d))'$, and in the same way $Ma \in \pi(S_n)'$.
Hence $P := TM = MT$ projects onto the center $\pi(\mathcal{Z}_n)$,

Completely symmetric states on $\mathcal{B}((\mathbb{C}^d)^{\otimes n})$

Observables (operators) on $\mathcal{H} := \mathbb{C}^d \otimes \dots \otimes \mathbb{C}^d$ can be 'twirled' and averaged:

$$Ta := \int_{SU(d)} (u \otimes \dots \otimes u)^* a (u \otimes \dots \otimes u) du ;$$
$$Ma := \frac{1}{n!} \sum_{\sigma \in S_n} \pi(\sigma) a \pi(\sigma) .$$

Clearly, $Ta \in \pi'(SU(d))'$, and in the same way $Ma \in \pi(S_n)'$.

Hence $P := TM = MT$ projects onto the center $\pi(\mathcal{Z}_n)$,

Dually P^* takes a state ϑ , restricts it to the center, and then extends it to a completely symmetric state on $\mathcal{B}(\mathcal{H})$:

$$(P^* \vartheta)(a) := \vartheta(Pa) .$$

Completely symmetric states on $\mathcal{B}((\mathbb{C}^d)^{\otimes n})$

Observables (operators) on $\mathcal{H} := \mathbb{C}^d \otimes \dots \otimes \mathbb{C}^d$ can be 'twirled' and averaged:

$$Ta := \int_{SU(d)} (u \otimes \dots \otimes u)^* a (u \otimes \dots \otimes u) du ;$$
$$Ma := \frac{1}{n!} \sum_{\sigma \in S_n} \pi(\sigma) a \pi(\sigma) .$$

Clearly, $Ta \in \pi'(SU(d))'$, and in the same way $Ma \in \pi(S_n)'$.
Hence $P := TM = MT$ projects onto the center $\pi(\mathcal{Z}_n)$,
Dually P^* takes a state ϑ , restricts it to the center, and then extends it to a completely symmetric state on $\mathcal{B}(\mathcal{H})$:

$$(P^* \vartheta)(a) := \vartheta(Pa) .$$

Theorem (Separability of completely symmetric states)

Let ϑ be a completely symmetric state on $\mathcal{B}((\mathbb{C}^d)^{\otimes n})$. Then ϑ is separable iff its restriction to \mathcal{Z} lies in the convex hull of the restricted product states.

Completely symmetric states on $\mathcal{B}((\mathbb{C}^d)^{\otimes n})$

Observables (operators) on $\mathcal{H} := \mathbb{C}^d \otimes \dots \otimes \mathbb{C}^d$ can be 'twirled' and averaged:

$$Ta := \int_{SU(d)} (u \otimes \dots \otimes u)^* a (u \otimes \dots \otimes u) du ;$$
$$Ma := \frac{1}{n!} \sum_{\sigma \in S_n} \pi(\sigma) a \pi(\sigma) .$$

Clearly, $Ta \in \pi'(SU(d))'$, and in the same way $Ma \in \pi(S_n)'$.
Hence $P := TM = MT$ projects onto the center $\pi(\mathcal{Z}_n)$,
Dually P^* takes a state ϑ , restricts it to the center, and then extends it to a completely symmetric state on $\mathcal{B}(\mathcal{H})$:

$$(P^* \vartheta)(a) := \vartheta(Pa) .$$

Theorem (Separability of completely symmetric states)

Let ϑ be a completely symmetric state on $\mathcal{B}((\mathbb{C}^d)^{\otimes n})$. Then ϑ is separable iff its restriction to \mathcal{Z} lies in the convex hull of the restricted product states.

Conclusion: We must calculate the shadow of the product states!

Proof.



Proof.

If ϑ is separable, then it is a convex combination of product states, so its restriction to \mathcal{Z} is a convex combination of such restrictions.



Proof.

If ϑ is separable, then it is a convex combination of product states, so its restriction to \mathcal{Z} is a convex combination of such restrictions.

Conversely, if for all $z \in \mathcal{Z}$ we have

$$\vartheta(z) = \sum_i \mu_i \langle \psi_i, z \psi_i \rangle ,$$

for some positive weights μ_i with sum 1 and unit product vectors ψ_i ,



Proof.

If ϑ is separable, then it is a convex combination of product states, so its restriction to \mathcal{Z} is a convex combination of such restrictions.

Conversely, if for all $z \in \mathcal{Z}$ we have

$$\vartheta(z) = \sum_i \mu_i \langle \psi_i, z \psi_i \rangle ,$$

for some positive weights μ_i with sum 1 and unit product vectors ψ_i , then since ϑ is completely symmetric, we have for all $x \in \mathcal{B}(\mathcal{H})$,

$$\begin{aligned} \vartheta(x) &= \vartheta(Px) = \sum_i \mu_i \langle \psi_i, Px \psi_i \rangle \\ &= \frac{1}{n!} \sum_i \sum_{\sigma \in S_n} \int_{SU(d)} \mu_i \langle \pi(\sigma)(u \otimes \dots \otimes u) \psi_i, x \pi(\sigma)(u \otimes \dots \otimes u) \psi_i \rangle du , \end{aligned}$$

which is a convex integral of product states, hence separable. □

The trace state

The trace state

Theorem

The strace state moves towards the regular trace as $d \rightarrow \infty$.

The trace state

Theorem

The trace state moves towards the regular trace as $d \rightarrow \infty$.

Proof.

First we calculate:

The trace state

Theorem

The trace state moves towards the regular trace as $d \rightarrow \infty$.

Proof.

First we calculate:

$$\begin{aligned}\mathrm{tr}_d^{\otimes n}(\pi(\sigma)) &= \sum_{i_1=1}^d \cdots \sum_{i_n=1}^d \langle \mathbf{e}_{i_1} \otimes \cdots \otimes \mathbf{e}_{i_n}, \pi(\sigma) \mathbf{e}_{i_1} \otimes \cdots \otimes \mathbf{e}_{i_n} \rangle \\ &= \sum_{i_1=1}^d \cdots \sum_{i_n=1}^d \delta_{i_1 i_{\sigma^{-1}(1)}} \cdots \delta_{i_n i_{\sigma^{-1}(n)}} \cdot \\ &= d^{\#(\text{cycles of } \sigma)}.\end{aligned}$$

The trace state

Theorem

The trace state moves towards the regular trace as $d \rightarrow \infty$.

Proof.

First we calculate:

$$\begin{aligned}\mathrm{tr}_d^{\otimes n}(\pi(\sigma)) &= \sum_{i_1=1}^d \cdots \sum_{i_n=1}^d \langle \mathbf{e}_{i_1} \otimes \cdots \otimes \mathbf{e}_{i_n}, \pi(\sigma) \mathbf{e}_{i_1} \otimes \cdots \otimes \mathbf{e}_{i_n} \rangle \\ &= \sum_{i_1=1}^d \cdots \sum_{i_n=1}^d \delta_{i_1 i_{\sigma^{-1}(1)}} \cdots \delta_{i_n i_{\sigma^{-1}(n)}} \cdot \\ &= d^{\#(\text{cycles of } \sigma)}.\end{aligned}$$

since for every cycle one summation variable remains.

The trace state

Theorem

The trace state moves towards the regular trace as $d \rightarrow \infty$.

Proof.

First we calculate:

$$\begin{aligned}\mathrm{tr}_d^{\otimes n}(\pi(\sigma)) &= \sum_{i_1=1}^d \cdots \sum_{i_n=1}^d \langle \mathbf{e}_{i_1} \otimes \cdots \otimes \mathbf{e}_{i_n}, \pi(\sigma) \mathbf{e}_{i_1} \otimes \cdots \otimes \mathbf{e}_{i_n} \rangle \\ &= \sum_{i_1=1}^d \cdots \sum_{i_n=1}^d \delta_{i_1 i_{\sigma^{-1}(1)}} \cdots \delta_{i_n i_{\sigma^{-1}(n)}} \cdot \\ &= d^{\#(\text{cycles of } \sigma)}.\end{aligned}$$

since for every cycle one summation variable remains. Hence:

The trace state

Theorem

The trace state moves towards the regular trace as $d \rightarrow \infty$.

Proof.

First we calculate:

$$\begin{aligned}\mathrm{tr}_d^{\otimes n}(\pi(\sigma)) &= \sum_{i_1=1}^d \cdots \sum_{i_n=1}^d \langle \mathbf{e}_{i_1} \otimes \cdots \otimes \mathbf{e}_{i_n}, \pi(\sigma) \mathbf{e}_{i_1} \otimes \cdots \otimes \mathbf{e}_{i_n} \rangle \\ &= \sum_{i_1=1}^d \cdots \sum_{i_n=1}^d \delta_{i_1 i_{\sigma^{-1}(1)}} \cdots \delta_{i_n i_{\sigma^{-1}(n)}} \cdot \\ &= d^{\#(\text{cycles of } \sigma)}.\end{aligned}$$

since for every cycle one summation variable remains. Hence:

$$\tau_d^{\otimes n}(p_Y) = \frac{d(Y)}{n!} \sum_{\sigma \in S_n} \chi_Y(\sigma) \frac{1}{d^n} \mathrm{tr}_d^{\otimes n}(\pi(\sigma)) \rightarrow \frac{d(Y)^2}{n!}, \quad (d \rightarrow \infty).$$



The shadow of the product states

The shadow of the product states

The state space $\mathcal{S}(\mathcal{Z})$ of the center \mathcal{Z} is a simplex whose corners are the states

$$\rho_Y : \quad p_{Y'} \mapsto \delta_{YY'} .$$

The shadow of the product states

The state space $\mathcal{S}(\mathcal{Z})$ of the center \mathcal{Z} is a simplex whose corners are the states

$$\rho_Y : \quad \rho_{Y'} \mapsto \delta_{YY'} .$$

The product states throw their shadow on this simplex:

The shadow of the product states

The state space $\mathcal{S}(\mathcal{Z})$ of the center \mathcal{Z} is a simplex whose corners are the states

$$\rho_Y : \quad p_{Y'} \mapsto \delta_{YY'} .$$

The product states throw their shadow on this simplex:
the affine components of the product state $\psi_1 \otimes \dots \otimes \psi_n$ are the **weights**

$$w_\psi(Y) := \langle \psi_1 \otimes \dots \otimes \psi_n, \pi(p_Y) \psi_1 \otimes \dots \otimes \psi_n \rangle .$$

The shadow of the product states

The state space $\mathcal{S}(\mathcal{Z})$ of the center \mathcal{Z} is a simplex whose corners are the states

$$\rho_Y : \quad p_{Y'} \mapsto \delta_{YY'} .$$

The product states throw their shadow on this simplex:
the affine components of the product state $\psi_1 \otimes \dots \otimes \psi_n$ are the **weights**

$$w_\psi(Y) := \langle \psi_1 \otimes \dots \otimes \psi_n, \pi(p_Y) \psi_1 \otimes \dots \otimes \psi_n \rangle .$$

We note that the the regular trace has the following weights:

$$w_{\text{reg}}(Y) := \tau_{\text{reg}}(p_Y) = \frac{d(Y)^2}{n!} .$$

The shadow of the product states

The state space $\mathcal{S}(\mathcal{Z})$ of the center \mathcal{Z} is a simplex whose corners are the states

$$\rho_Y : \quad p_{Y'} \mapsto \delta_{YY'} .$$

The product states throw their shadow on this simplex:
the affine components of the product state $\psi_1 \otimes \dots \otimes \psi_n$ are the **weights**

$$w_\psi(Y) := \langle \psi_1 \otimes \dots \otimes \psi_n, \pi(p_Y) \psi_1 \otimes \dots \otimes \psi_n \rangle .$$

We note that the the regular trace has the following weights:

$$w_{\text{reg}}(Y) := \tau_{\text{reg}}(p_Y) = \frac{d(Y)^2}{n!} .$$

Now here's our basic connection between **entanglement** and **classical mathematics**:

The shadow of the product states

The state space $\mathcal{S}(\mathcal{Z})$ of the center \mathcal{Z} is a simplex whose corners are the states

$$\rho_Y : \quad p_{Y'} \mapsto \delta_{YY'} .$$

The product states throw their shadow on this simplex:
the affine components of the product state $\psi_1 \otimes \dots \otimes \psi_n$ are the **weights**

$$w_\psi(Y) := \langle \psi_1 \otimes \dots \otimes \psi_n, \pi(p_Y) \psi_1 \otimes \dots \otimes \psi_n \rangle .$$

We note that the the regular trace has the following weights:

$$w_{\text{reg}}(Y) := \tau_{\text{reg}}(p_Y) = \frac{d(Y)^2}{n!} .$$

Now here's our basic connection between **entanglement** and **classical mathematics**:

Theorem

*The density of a product state $\psi_1 \otimes \dots \otimes \psi_n$ with respect to the regular trace is the normalized **immanant** of the Gram matrix of $\psi_1, \psi_2, \dots, \psi_n$.*

Immanants of a matrix

Immanants of a matrix

Let A be an $n \times n$ matrix, and let Y be a Young frame with n boxes.

Immanants of a matrix

Let A be an $n \times n$ matrix, and let Y be a Young frame with n boxes.

Then the **immanant** $\text{Imm}_Y(A)$ of this matrix associated to Y is defined as

$$\text{Imm}_Y(A) := \sum_{\sigma \in S_n} \chi_Y(\sigma) a_{1\sigma(1)} a_{2\sigma(2)} \cdots a_{n\sigma(n)}.$$

Immanants of a matrix

Let A be an $n \times n$ matrix, and let Y be a Young frame with n boxes.

Then the **immanant** $\text{Imm}_Y(A)$ of this matrix associated to Y is defined as

$$\text{Imm}_Y(A) := \sum_{\sigma \in S_n} \chi_Y(\sigma) a_{1\sigma(1)} a_{2\sigma(2)} \cdots a_{n\sigma(n)}.$$

The **normalized immanant** $\tilde{\text{Imm}}_Y(A)$ is defined so as to have $\tilde{\text{Imm}}(\mathbb{1}) = 1$:

$$\tilde{\text{Imm}}_Y(A) := \frac{\text{Imm}_Y(A)}{d(Y)}.$$

Immanants of a matrix

Let A be an $n \times n$ matrix, and let Y be a Young frame with n boxes. Then the **immanant** $\text{Imm}_Y(A)$ of this matrix associated to Y is defined as

$$\text{Imm}_Y(A) := \sum_{\sigma \in S_n} \chi_Y(\sigma) a_{1\sigma(1)} a_{2\sigma(2)} \cdots a_{n\sigma(n)}.$$

The **normalized immanant** $\tilde{\text{Imm}}_Y(A)$ is defined so as to have $\tilde{\text{Imm}}(\mathbb{1}) = 1$:

$$\tilde{\text{Imm}}_Y(A) := \frac{\text{Imm}_Y(A)}{d(Y)}.$$

Note the following well-known special cases:

$$\text{Imm}_{\begin{array}{|c|} \hline \square \\ \square \\ \square \\ \square \\ \square \\ \square \\ \square \\ \square \\ \hline \end{array}}(A) = \det(A) \quad \text{and} \quad \text{Imm}_{\square\square\square\square\square\square}(A) = \text{per}(A).$$

Immanants of a matrix

Let A be an $n \times n$ matrix, and let Y be a Young frame with n boxes. Then the **immanant** $\text{Imm}_Y(A)$ of this matrix associated to Y is defined as

$$\text{Imm}_Y(A) := \sum_{\sigma \in S_n} \chi_Y(\sigma) a_{1\sigma(1)} a_{2\sigma(2)} \cdots a_{n\sigma(n)}.$$

The **normalized immanant** $\tilde{\text{Imm}}_Y(A)$ is defined so as to have $\tilde{\text{Imm}}(\mathbb{1}) = 1$:

$$\tilde{\text{Imm}}_Y(A) := \frac{\text{Imm}_Y(A)}{d(Y)}.$$

Note the following well-known special cases:

$$\text{Imm}_{\begin{array}{|c|} \hline \square \\ \square \\ \square \\ \square \\ \square \\ \square \\ \square \\ \square \\ \hline \end{array}}(A) = \det(A) \quad \text{and} \quad \text{Imm}_{\begin{array}{|c|c|c|c|c|c|} \hline \square & \square & \square & \square & \square & \square \\ \hline \end{array}}(A) = \text{per}(A).$$

We mention the following inequalities: for all positive definite matrices A and all Young frames Y :

$$\det(A) \leq \tilde{\text{Imm}}_Y(A) \leq \text{per}(A).$$

Immanants of a matrix

Let A be an $n \times n$ matrix, and let Y be a Young frame with n boxes. Then the **immanant** $\text{Imm}_Y(A)$ of this matrix associated to Y is defined as

$$\text{Imm}_Y(A) := \sum_{\sigma \in S_n} \chi_Y(\sigma) a_{1\sigma(1)} a_{2\sigma(2)} \cdots a_{n\sigma(n)}.$$

The **normalized immanant** $\tilde{\text{Imm}}_Y(A)$ is defined so as to have $\tilde{\text{Imm}}(\mathbb{1}) = 1$:

$$\tilde{\text{Imm}}_Y(A) := \frac{\text{Imm}_Y(A)}{d(Y)}.$$

Note the following well-known special cases:

$$\text{Imm}_{\begin{array}{|c|} \hline \square \\ \square \\ \square \\ \square \\ \square \\ \square \\ \square \\ \square \\ \hline \end{array}}(A) = \det(A) \quad \text{and} \quad \text{Imm}_{\begin{array}{|c|c|c|c|c|c|} \hline \square & \square & \square & \square & \square & \square \\ \hline \end{array}}(A) = \text{per}(A).$$

We mention the following inequalities: for all positive definite matrices A and all Young frames Y :

$$\det(A) \leq \tilde{\text{Imm}}_Y(A) \leq \text{per}(A).$$

The first inequality was proved by Schur in 1918, the second was **conjectured** by Elliott Lieb in 1967, and is **still open!**

The weight formula

Proof.



The weight formula

Proof.

This is not more than a concatenation of definitions connecting quantum information (**entanglement**) to algebra (**immanants**).



The weight formula

Proof.

This is not more than a concatenation of definitions connecting quantum information (**entanglement**) to algebra (**immanants**).

The weight of the extreme point ρ_Y in the expansion of the pure product state $\psi_1 \otimes \dots \otimes \psi_n$ is equal to



The weight formula

Proof.

This is not more than a concatenation of definitions connecting quantum information (**entanglement**) to algebra (**immanants**).

The weight of the extreme point ρ_Y in the expansion of the pure product state $\psi_1 \otimes \dots \otimes \psi_n$ is equal to

$$w_\psi(Y) = \langle \psi_1 \otimes \dots \otimes \psi_n, \pi(\rho_Y) \psi_1 \otimes \dots \otimes \psi_n \rangle$$



The weight formula

Proof.

This is not more than a concatenation of definitions connecting quantum information (**entanglement**) to algebra (**immanants**).

The weight of the extreme point ρ_Y in the expansion of the pure product state $\psi_1 \otimes \dots \otimes \psi_n$ is equal to

$$\begin{aligned}w_\psi(Y) &= \langle \psi_1 \otimes \dots \otimes \psi_n, \pi(\rho_Y) \psi_1 \otimes \dots \otimes \psi_n \rangle \\ &= \frac{d(Y)}{n!} \sum_{\sigma \in S_n} \chi_Y(\sigma) \langle \psi_1 \otimes \dots \otimes \psi_n, \pi(\sigma) \psi_1 \otimes \dots \otimes \psi_n \rangle\end{aligned}$$



The weight formula

Proof.

This is not more than a concatenation of definitions connecting quantum information (**entanglement**) to algebra (**immanants**).

The weight of the extreme point ρ_Y in the expansion of the pure product state $\psi_1 \otimes \dots \otimes \psi_n$ is equal to

$$\begin{aligned}w_\psi(Y) &= \langle \psi_1 \otimes \dots \otimes \psi_n, \pi(\rho_Y) \psi_1 \otimes \dots \otimes \psi_n \rangle \\ &= \frac{d(Y)}{n!} \sum_{\sigma \in S_n} \chi_Y(\sigma) \langle \psi_1 \otimes \dots \otimes \psi_n, \pi(\sigma) \psi_1 \otimes \dots \otimes \psi_n \rangle \\ &= \frac{d(Y)}{n!} \sum_{\sigma \in S_n} \chi_Y(\sigma) \langle \psi_1 \otimes \dots \otimes \psi_n, \psi_{\sigma^{-1}(1)} \otimes \dots \otimes \psi_{\sigma^{-1}(n)} \rangle\end{aligned}$$



The weight formula

Proof.

This is not more than a concatenation of definitions connecting quantum information (**entanglement**) to algebra (**immanants**).

The weight of the extreme point ρ_Y in the expansion of the pure product state $\psi_1 \otimes \dots \otimes \psi_n$ is equal to

$$\begin{aligned}w_{\psi}(Y) &= \langle \psi_1 \otimes \dots \otimes \psi_n, \pi(\rho_Y) \psi_1 \otimes \dots \otimes \psi_n \rangle \\&= \frac{d(Y)}{n!} \sum_{\sigma \in S_n} \chi_Y(\sigma) \langle \psi_1 \otimes \dots \otimes \psi_n, \pi(\sigma) \psi_1 \otimes \dots \otimes \psi_n \rangle \\&= \frac{d(Y)}{n!} \sum_{\sigma \in S_n} \chi_Y(\sigma) \langle \psi_1 \otimes \dots \otimes \psi_n, \psi_{\sigma^{-1}(1)} \otimes \dots \otimes \psi_{\sigma^{-1}(n)} \rangle \\&= \frac{d(Y)}{n!} \sum_{\sigma \in S_n} \chi_Y(\sigma) \prod_{j=1}^n \langle \psi_j, \psi_{\sigma^{-1}(j)} \rangle\end{aligned}$$



The weight formula

Proof.

This is not more than a concatenation of definitions connecting quantum information (**entanglement**) to algebra (**immanants**).


The weight of the extreme point ρ_Y in the expansion of the pure product state $\psi_1 \otimes \dots \otimes \psi_n$ is equal to

$$\begin{aligned}w_\psi(Y) &= \langle \psi_1 \otimes \dots \otimes \psi_n, \pi(\rho_Y) \psi_1 \otimes \dots \otimes \psi_n \rangle \\&= \frac{d(Y)}{n!} \sum_{\sigma \in S_n} \chi_Y(\sigma) \langle \psi_1 \otimes \dots \otimes \psi_n, \pi(\sigma) \psi_1 \otimes \dots \otimes \psi_n \rangle \\&= \frac{d(Y)}{n!} \sum_{\sigma \in S_n} \chi_Y(\sigma) \langle \psi_1 \otimes \dots \otimes \psi_n, \psi_{\sigma^{-1}(1)} \otimes \dots \otimes \psi_{\sigma^{-1}(n)} \rangle \\&= \frac{d(Y)}{n!} \sum_{\sigma \in S_n} \chi_Y(\sigma) \prod_{j=1}^n \langle \psi_j, \psi_{\sigma^{-1}(j)} \rangle \\&= \frac{d(Y)}{n!} \text{Imm}_Y(G(\psi)) = \frac{d(Y)^2}{n!} \tilde{\text{Imm}}_Y(G(\psi)).\end{aligned}$$




The simplex $\mathcal{S}(\mathcal{Z}_n)$ for $n = 3$


The simplex $\mathcal{S}(\mathcal{Z}_n)$ for $n = 3$




The simplex $\mathcal{S}(\mathcal{Z}_n)$ for $n = 3$




The simplex $\mathcal{S}(\mathcal{Z}_n)$ for $n = 3$



The simplex $\mathcal{S}(\mathcal{Z}_n)$ for $n = 3$




The simplex $\mathcal{S}(\mathcal{Z}_n)$ for $n = 3$




$$\tau_{\text{reg}} = P^*(|\psi\rangle\langle\psi|) \text{ with } G(\psi) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix};$$

The simplex $\mathcal{S}(\mathcal{Z}_n)$ for $n = 3$




$$\tau_{\text{reg}} = P^*(|\psi\rangle\langle\psi|) \text{ with } G(\psi) = \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix};$$

The simplex $\mathcal{S}(\mathcal{Z}_n)$ for $n = 3$



$$\rho_+ = P^*(|\psi\rangle\langle\psi|) \text{ with } G(\psi) = \begin{pmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{pmatrix};$$

The simplex $\mathcal{S}(\mathcal{Z}_n)$ for $n = 3$



$$\rho_{120^\circ} = P^*(|\psi\rangle\langle\psi|) \text{ with } G(\psi) = \begin{pmatrix} 1 & -\frac{1}{2} & -\frac{1}{2} \\ -\frac{1}{2} & 1 & -\frac{1}{2} \\ -\frac{1}{2} & -\frac{1}{2} & 1 \end{pmatrix};$$

Optimal Bell inequalities for $n = 3$

Optimal Bell inequalities for $n = 3$

For $n = 3$ the separable region is a polytope, having a finite number (3) of extreme points.

Optimal Bell inequalities for $n = 3$

For $n = 3$ the separable region is a polytope, having a finite number (3) of extreme points.

We need only two linear ('Bell') inequalities in order to distinguish the separable from the entangled completely symmetric states.

$$\rho(p_+ + 5p_-) \geq 1 ;$$

$$\rho(4p_+ + p_-) \geq 1 .$$

Optimal Bell inequalities for $n = 3$

For $n = 3$ the separable region is a polytope, having a finite number (3) of extreme points.

We need only two linear ('Bell') inequalities in order to distinguish the separable from the entangled completely symmetric states.

$$\rho(p_+ + 5p_-) \geq 1 ;$$

$$\rho(4p_+ + p_-) \geq 1 .$$

They correspond to the green lines in the figure.

Optimal Bell inequalities for $n = 3$

For $n = 3$ the separable region is a polytope, having a finite number (3) of extreme points.

We need only two linear ('Bell') inequalities in order to distinguish the separable from the entangled completely symmetric states.

$$\rho(p_+ + 5p_-) \geq 1 ;$$

$$\rho(4p_+ + p_-) \geq 1 .$$

They correspond to the green lines in the figure.

Questions:

Optimal Bell inequalities for $n = 3$

For $n = 3$ the separable region is a polytope, having a finite number (3) of extreme points.

We need only two linear ('Bell') inequalities in order to distinguish the separable from the entangled completely symmetric states.

$$\rho(p_+ + 5p_-) \geq 1 ;$$

$$\rho(4p_+ + p_-) \geq 1 .$$

They correspond to the green lines in the figure.

Questions:

- ▶ Is the separable completely symmetric region (the 'shadow') always a polytope?

Optimal Bell inequalities for $n = 3$

For $n = 3$ the separable region is a polytope, having a finite number (3) of extreme points.

We need only two linear ('Bell') inequalities in order to distinguish the separable from the entangled completely symmetric states.

$$\rho(p_+ + 5p_-) \geq 1 ;$$

$$\rho(4p_+ + p_-) \geq 1 .$$

They correspond to the green lines in the figure.

Questions:

- ▶ Is the separable completely symmetric region (the 'shadow') always a polytope?
- ▶ What is the general shape of this region?

Optimal Bell inequalities for $n = 3$

For $n = 3$ the separable region is a polytope, having a finite number (3) of extreme points.

We need only two linear ('Bell') inequalities in order to distinguish the separable from the entangled completely symmetric states.

$$\rho(p_+ + 5p_-) \geq 1 ;$$

$$\rho(4p_+ + p_-) \geq 1 .$$

They correspond to the green lines in the figure.

Questions:

- ▶ Is the separable completely symmetric region (the 'shadow') always a polytope?
- ▶ What is the general shape of this region?
- ▶ Does it grow or shrink with increasing n ?

General n : the shadow touches only one corner



General n : the shadow touches only one corner

Theorem

Only the state $\rho_+ = \rho_{\square\square\square\square\square\square}$ (n boxes) is separable, all other extremal states ρ_Y on $\mathcal{Z}_{n,d}$ are entangled.



General n : the shadow touches only one corner

Theorem

Only the state $\rho_+ = \rho_{\square\square\square\square\square\square}$ (n boxes) is separable, all other extremal states ρ_Y on $\mathcal{Z}_{n,d}$ are entangled.

Proof.



General n : the shadow touches only one corner

Theorem

Only the state $\rho_+ = \rho_{\square\square\square\square\square\square}$ (n boxes) is separable, all other extremal states ρ_Y on $\mathcal{Z}_{n,d}$ are entangled.

Proof.

Choose $\vartheta \in \mathbb{C}^d$.



General n : the shadow touches only one corner

Theorem

Only the state $\rho_+ = \rho_{\square\square\square\square\square\square}$ (n boxes) is separable, all other extremal states ρ_Y on $\mathcal{Z}_{n,d}$ are entangled.

Proof.

Choose $\vartheta \in \mathbb{C}^d$.

Since $\vartheta \otimes \dots \otimes \vartheta \in \mathcal{H}_+$, we have $\langle \vartheta \otimes \dots \otimes \vartheta, \rho_+ \vartheta \otimes \dots \otimes \vartheta \rangle = 1$. Hence $\rho_+ = P^*(|\vartheta \otimes \dots \otimes \vartheta\rangle\langle \vartheta \otimes \dots \otimes \vartheta|)$, a convex integral of product states.



General n : the shadow touches only one corner

Theorem

Only the state $\rho_+ = \rho_{\square\square\square\square\square\square}$ (n boxes) is separable, all other extremal states ρ_Y on $\mathcal{Z}_{n,d}$ are entangled.

Proof.

Choose $\vartheta \in \mathbb{C}^d$.

Since $\vartheta \otimes \dots \otimes \vartheta \in \mathcal{H}_+$, we have $\langle \vartheta \otimes \dots \otimes \vartheta, \rho_+ \vartheta \otimes \dots \otimes \vartheta \rangle = 1$. Hence $\rho_+ = P^*(|\vartheta \otimes \dots \otimes \vartheta\rangle\langle \vartheta \otimes \dots \otimes \vartheta|)$, a convex integral of product states.

Conversely, suppose that $Y \neq \square\square\square\square\square\square$ (n boxes), and that ρ_Y is separable:

$$\rho_Y(a) = \sum_i \langle \psi_i, a \psi_i \rangle$$



General n : the shadow touches only one corner

Theorem

Only the state $\rho_+ = \rho_{\square\square\square\square\square\square}$ (n boxes) is separable, all other extremal states ρ_Y on $\mathcal{Z}_{n,d}$ are entangled.

Proof.

Choose $\vartheta \in \mathbb{C}^d$.

Since $\vartheta \otimes \dots \otimes \vartheta \in \mathcal{H}_+$, we have $\langle \vartheta \otimes \dots \otimes \vartheta, \rho_+ \vartheta \otimes \dots \otimes \vartheta \rangle = 1$. Hence $\rho_+ = P^*(|\vartheta \otimes \dots \otimes \vartheta\rangle\langle \vartheta \otimes \dots \otimes \vartheta|)$, a convex integral of product states.

Conversely, suppose that $Y \neq \square\square\square\square\square\square$ (n boxes), and that ρ_Y is separable:

$$\rho_Y(a) = \sum_i \langle \psi_i, a \psi_i \rangle$$

Then $0 = \rho_Y(\rho_+) = \sum_i \langle \psi_i, \rho_+ \psi_i \rangle$, i.e. $\rho_+ \psi_i = 0$ for all i .



General n : the shadow touches only one corner

Theorem

Only the state $\rho_+ = \rho_{\square\square\square\square\square\square}$ (n boxes) is separable, all other extremal states ρ_Y on $\mathcal{Z}_{n,d}$ are entangled.

Proof.

Choose $\vartheta \in \mathbb{C}^d$.

Since $\vartheta \otimes \dots \otimes \vartheta \in \mathcal{H}_+$, we have $\langle \vartheta \otimes \dots \otimes \vartheta, \rho_+ \vartheta \otimes \dots \otimes \vartheta \rangle = 1$. Hence $\rho_+ = P^*(|\vartheta \otimes \dots \otimes \vartheta\rangle\langle \vartheta \otimes \dots \otimes \vartheta|)$, a convex integral of product states.

Conversely, suppose that $Y \neq \square\square\square\square\square\square$ (n boxes), and that ρ_Y is separable:

$$\rho_Y(a) = \sum_i \langle \psi_i, a\psi_i \rangle$$

Then $0 = \rho_Y(\rho_+) = \sum_i \langle \psi_i, \rho_+ \psi_i \rangle$, i.e. $\rho_+ \psi_i = 0$ for all i .
However, product vectors with this property do not exist!



General n : the shadow touches only one corner

Theorem

Only the state $\rho_+ = \rho_{\square\square\square\square\square\square}$ (n boxes) is separable, all other extremal states ρ_Y on $\mathcal{Z}_{n,d}$ are entangled.

Proof.

Choose $\vartheta \in \mathbb{C}^d$.

Since $\vartheta \otimes \dots \otimes \vartheta \in \mathcal{H}_+$, we have $\langle \vartheta \otimes \dots \otimes \vartheta, \rho_+ \vartheta \otimes \dots \otimes \vartheta \rangle = 1$. Hence $\rho_+ = P^*(|\vartheta \otimes \dots \otimes \vartheta\rangle\langle \vartheta \otimes \dots \otimes \vartheta|)$, a convex integral of product states.

Conversely, suppose that $Y \neq \square\square\square\square\square\square$ (n boxes), and that ρ_Y is separable:

$$\rho_Y(a) = \sum_i \langle \psi_i, a \psi_i \rangle$$

Then $0 = \rho_Y(\rho_+) = \sum_i \langle \psi_i, \rho_+ \psi_i \rangle$, i.e. $\rho_+ \psi_i = 0$ for all i .

However, product vectors with this property do not exist!

Indeed, suppose that the product vector $\psi_1 \otimes \dots \otimes \psi_n$ is orthogonal to $\rho_+ \mathcal{H}$.



General n : the shadow touches only one corner

Theorem

Only the state $\rho_+ = \rho_{\square\square\square\square\square\square}$ (n boxes) is separable, all other extremal states ρ_Y on $\mathcal{Z}_{n,d}$ are entangled.

Proof.

Choose $\vartheta \in \mathbb{C}^d$.

Since $\vartheta \otimes \dots \otimes \vartheta \in \mathcal{H}_+$, we have $\langle \vartheta \otimes \dots \otimes \vartheta, \rho_+ \vartheta \otimes \dots \otimes \vartheta \rangle = 1$. Hence $\rho_+ = P^*(|\vartheta \otimes \dots \otimes \vartheta\rangle\langle \vartheta \otimes \dots \otimes \vartheta|)$, a convex integral of product states.

Conversely, suppose that $Y \neq \square\square\square\square\square\square$ (n boxes), and that ρ_Y is separable:

$$\rho_Y(a) = \sum_i \langle \psi_i, a \psi_i \rangle$$

Then $0 = \rho_Y(\rho_+) = \sum_i \langle \psi_i, \rho_+ \psi_i \rangle$, i.e. $\rho_+ \psi_i = 0$ for all i .

However, product vectors with this property do not exist!

Indeed, suppose that the product vector $\psi_1 \otimes \dots \otimes \psi_n$ is orthogonal to $\rho_+ \mathcal{H}$.

Then it is orthogonal to all vectors of the form $\vartheta \otimes \dots \otimes \vartheta$ with $\vartheta \in \mathbb{C}^d$:

$$0 = \langle \vartheta \otimes \dots \otimes \vartheta, \psi_1 \otimes \dots \otimes \psi_n \rangle = \prod_{j=1}^n \langle \vartheta, \psi_j \rangle. \quad \text{But: } \bigcup_{j=1}^n \{\psi_j\}^\perp \neq \mathbb{C}^d,$$

the left hand side having Lebesgue measure 0.



Shadow moves away from the antisymmetric state



Shadow moves away from the antisymmetric state

Theorem

For all separable states ρ we have

$$\rho(p_-) \leq \frac{1}{n!}$$

with equality only for the regular trace state.



Shadow moves away from the antisymmetric state

Theorem

For all separable states ρ we have

$$\rho(p_-) \leq \frac{1}{n!}$$

with equality only for the regular trace state.

Proof.



Shadow moves away from the antisymmetric state

Theorem

For all separable states ρ we have

$$\rho(p_-) \leq \frac{1}{n!}$$

with equality only for the regular trace state.

Proof.

The determinant of the Gram matrix of an n -tuple of unit vectors is equal to

$$\begin{aligned} \det(\langle \psi_i, \psi_j \rangle) &= \det\left(\sum_{k=1}^n \langle \psi_i, \mathbf{e}_k \rangle \langle \mathbf{e}_k, \psi_j \rangle\right) \\ &= \left| \det(\langle \psi_i, \mathbf{e}_k \rangle) \right|^2 = \text{vol}(\psi_1, \psi_2, \dots, \psi_n)^2 \leq 1 ; \end{aligned}$$



Shadow moves away from the antisymmetric state

Theorem

For all separable states ρ we have

$$\rho(p_-) \leq \frac{1}{n!}$$

with equality only for the regular trace state.

Proof.

The determinant of the Gram matrix of an n -tuple of unit vectors is equal to

$$\begin{aligned} \det(\langle \psi_i, \psi_j \rangle) &= \det\left(\sum_{k=1}^n \langle \psi_i, \mathbf{e}_k \rangle \langle \mathbf{e}_k, \psi_j \rangle\right) \\ &= \left| \det(\langle \psi_i, \mathbf{e}_k \rangle) \right|^2 = \text{vol}(\psi_1, \psi_2, \dots, \psi_n)^2 \leq 1 ; \end{aligned}$$

Hence

$$\langle \psi_1 \otimes \dots \otimes \psi_n, p_- \psi_1 \otimes \dots \otimes \psi_n \rangle \leq \tau_{\text{reg}}(p_-) = \frac{1}{n!} .$$



The Schur and Lieb inequalities

We have $2\mathcal{P}(n) - 3$ inequalities, which divide the state space $\mathcal{S}(\mathcal{Z}_n)$ into compartments, and claim the the shadow of the product states falls into one of them.

Schur's 1918 inequality states that for all separables states ρ and all Young frames $Y \neq \{-\}$:

$$\rho(p_Y) \geq d(Y)^2 \rho(p_-).$$

Lieb's 1967 conjecture hopes that for all separable ρ and all Young frames

$Y \neq \{+\}$:


$$\rho(p_Y) \leq d(Y)^2 \rho(p_+).$$

The last **trivial inequality** says that for all separable ρ :

$$\rho(p_-) \leq \rho(p_+).$$


These are all Bell inequalities.

The immanant inequalities for $n = 3$ in a picture




These are all Bell inequalities, but not all optimal.

The immanant inequalities for $n = 3$ in a picture




These are all Bell inequalities, but not all optimal.

The immanant inequalities for $n = 3$ in a picture




These are all Bell inequalities, but not all optimal.

The immanant inequalities for $n = 3$ in a picture




These are all Bell inequalities, but not all optimal.

The immanant inequalities for $n = 3$ in a picture




These are all Bell inequalities, but not all optimal.

The immanant inequalities for $n = 3$ in a picture




These are all Bell inequalities, but not all optimal.

The immanant inequalities for $n = 3$ in a picture



These are all Bell inequalities, but not all optimal.

The immanent inequalities for $n = 3$ in a picture




These are all Bell inequalities, but not all optimal.

The separable region for $n = 4$


The separable region for $n = 4$

In the case $n = 4$ there are five Young diagrams:



The separable region for $n = 4$

In the case $n = 4$ there are five Young diagrams:




Theorem (Barrett, Hall, Loewy (1998) translated to quantum states)

The set of completely symmetric separable states on $\mathcal{B}(\mathbb{C}^d)^{\otimes 4}$ is the convex hull of 7 extreme points.

The separable region for $n = 4$

In the case $n = 4$ there are five Young diagrams:




Theorem (Barrett, Hall, Loewy (1998) translated to quantum states)

The set of completely symmetric separable states on $\mathcal{B}(\mathbb{C}^d)^{\otimes 4}$ is the convex hull of 7 extreme points. These extremal states are obtained by twirling and averaging 7 configurations of unit vectors in \mathbb{C}^d (with $d \geq 4$ to fit all of them). These configurations are given by the Gram matrices

The separable region for $n = 4$

In the case $n = 4$ there are five Young diagrams:




Theorem (Barrett, Hall, Loewy (1998) translated to quantum states)

The set of completely symmetric separable states on $\mathcal{B}(\mathbb{C}^d)^{\otimes 4}$ is the convex hull of 7 extreme points. These extremal states are obtained by twirling and averaging 7 configurations of unit vectors in \mathbb{C}^d (with $d \geq 4$ to fit all of them). These configurations are given by the Gram matrices

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

The separable region for $n = 4$

In the case $n = 4$ there are five Young diagrams:




Theorem (Barrett, Hall, Loewy (1998) translated to quantum states)

The set of completely symmetric separable states on $\mathcal{B}(\mathbb{C}^d)^{\otimes 4}$ is the convex hull of 7 extreme points. These extremal states are obtained by twirling and averaging 7 configurations of unit vectors in \mathbb{C}^d (with $d \geq 4$ to fit all of them). These configurations are given by the Gram matrices

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{pmatrix},$$

The separable region for $n = 4$

In the case $n = 4$ there are five Young diagrams:




Theorem (Barrett, Hall, Loewy (1998) translated to quantum states)

The set of completely symmetric separable states on $\mathcal{B}(\mathbb{C}^d)^{\otimes 4}$ is the convex hull of 7 extreme points. These extremal states are obtained by twirling and averaging 7 configurations of unit vectors in \mathbb{C}^d (with $d \geq 4$ to fit all of them). These configurations are given by the Gram matrices

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

The separable region for $n = 4$

In the case $n = 4$ there are five Young diagrams:



Theorem (Barrett, Hall, Loewy (1998) translated to quantum states)


The set of completely symmetric separable states on $\mathcal{B}(\mathbb{C}^d)^{\otimes 4}$ is the convex hull of 7 extreme points. These extremal states are obtained by twirling and averaging 7 configurations of unit vectors in \mathbb{C}^d (with $d \geq 4$ to fit all of them).

These configurations are given by the Gram matrices

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 \end{pmatrix}$$

The separable region for $n = 4$

In the case $n = 4$ there are five Young diagrams:



Theorem (Barrett, Hall, Loewy (1998) translated to quantum states)


The set of completely symmetric separable states on $\mathcal{B}(\mathbb{C}^d)^{\otimes 4}$ is the convex hull of 7 extreme points. These extremal states are obtained by twirling and averaging 7 configurations of unit vectors in \mathbb{C}^d (with $d \geq 4$ to fit all of them).

These configurations are given by the Gram matrices

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 \end{pmatrix}$$
$$\begin{pmatrix} 1 & -\frac{1}{2} & -\frac{1}{2} & 0 \\ -\frac{1}{2} & 1 & -\frac{1}{2} & 0 \\ -\frac{1}{2} & -\frac{1}{2} & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix},$$

The separable region for $n = 4$

In the case $n = 4$ there are five Young diagrams:



Theorem (Barrett, Hall, Loewy (1998) translated to quantum states)


The set of completely symmetric separable states on $\mathcal{B}(\mathbb{C}^d)^{\otimes 4}$ is the convex hull of 7 extreme points. These extremal states are obtained by twirling and averaging 7 configurations of unit vectors in \mathbb{C}^d (with $d \geq 4$ to fit all of them).

These configurations are given by the Gram matrices

$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 \end{pmatrix}$$
$$\begin{pmatrix} 1 & -\frac{1}{2} & -\frac{1}{2} & 0 \\ -\frac{1}{2} & 1 & -\frac{1}{2} & 0 \\ -\frac{1}{2} & -\frac{1}{2} & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & -\frac{1}{3} & -\frac{1}{3} & -\frac{1}{3} \\ -\frac{1}{3} & 1 & -\frac{1}{3} & -\frac{1}{3} \\ -\frac{1}{3} & -\frac{1}{3} & 1 & -\frac{1}{3} \\ -\frac{1}{3} & -\frac{1}{3} & -\frac{1}{3} & 1 \end{pmatrix},$$

The separable region for $n = 4$

In the case $n = 4$ there are five Young diagrams:



Theorem (Barrett, Hall, Loewy (1998) translated to quantum states)


The set of completely symmetric separable states on $\mathcal{B}(\mathbb{C}^d)^{\otimes 4}$ is the convex hull of 7 extreme points. These extremal states are obtained by twirling and averaging 7 configurations of unit vectors in \mathbb{C}^d (with $d \geq 4$ to fit all of them).

These configurations are given by the Gram matrices


$$\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & 1 & 0 & 0 \\ 1 & 1 & 0 & 0 \\ 0 & 0 & 1 & 1 \\ 0 & 0 & 1 & 1 \end{pmatrix}$$
$$\begin{pmatrix} 1 & -\frac{1}{2} & -\frac{1}{2} & 0 \\ -\frac{1}{2} & 1 & -\frac{1}{2} & 0 \\ -\frac{1}{2} & -\frac{1}{2} & 1 & 0 \\ 0 & 0 & 0 & 1 \end{pmatrix}, \quad \begin{pmatrix} 1 & -\frac{1}{3} & -\frac{1}{3} & -\frac{1}{3} \\ -\frac{1}{3} & 1 & -\frac{1}{3} & -\frac{1}{3} \\ -\frac{1}{3} & -\frac{1}{3} & 1 & -\frac{1}{3} \\ -\frac{1}{3} & -\frac{1}{3} & -\frac{1}{3} & 1 \end{pmatrix}, \quad \frac{1}{\sqrt{3}} \begin{pmatrix} \sqrt{3} & i & i & -i \\ -i & \sqrt{3} & i & i \\ -i & -i & \sqrt{3} & -i \\ i & -i & i & \sqrt{3} \end{pmatrix}.$$

Four qubits


Four qubits




Four qubits




Four qubits




Four qubits




Four qubits




Four qubits




Four qubits




Four qubits



Four qubits



Four qubits



Hope crashed at $n = 5$

Hope crashed at $n = 5$

Our hope was, to prove that for all $n \in \mathbb{N}$ the separable completely symmetric states would form a polytope.

Hope crashed at $n = 5$

Our hope was, to prove that for all $n \in \mathbb{N}$ the separable completely symmetric states would form a polytope.

However, this hope breaks down at $n = 5$:

Hope crashed at $n = 5$

Our hope was, to prove that for all $n \in \mathbb{N}$ the separable completely symmetric states would form a polytope.

However, this hope breaks down at $n = 5$:

Theorem (Barrett, Hall, Loewy (1999) translated)

The set of all completely symmetric separable states on $\mathcal{B}((\mathbb{C}^d)^{\otimes 5})$ has an infinite number of extremal points.

Hope crashed at $n = 5$

Our hope was, to prove that for all $n \in \mathbb{N}$ the separable completely symmetric states would form a polytope.

However, this hope breaks down at $n = 5$:


Theorem (Barrett, Hall, Loewy (1999) translated)

The set of all completely symmetric separable states on $\mathcal{B}((\mathbb{C}^d)^{\otimes 5})$ has an infinite number of extremal points.


In 1999 they showed that, already in the five qubit situation, the set of separable states on the center possesses a part that is **bulging outward**.

Proof that for $n = 5$ separable states do not form a polytope


Proof that for $n = 5$ separable states do not form a polytope




Proof that for $n = 5$ separables states do not form a polytope




Proof that for $n = 5$ separable states do not form a polytope




Proof that for $n = 5$ separable states do not form a polytope




Proof that for $n = 5$ separable states do not form a polytope




Proof that for $n = 5$ separable states do not form a polytope



Proof that for $n = 5$ separable states do not form a polytope




Proof that for $n = 5$ separable states do not form a polytope




Magnification

Magnification



Magnification



A measure for multiparticle entanglement

A measure for multiparticle entanglement

The maximal tensor norm:

A measure for multiparticle entanglement

The maximal tensor norm: Let $\mathcal{H} := (\mathbb{C}^d)^{\otimes n}$, and let V denote the set of linear functionals on $\mathcal{B}(\mathcal{H})$ of the form

$$x \mapsto \langle \psi_1 \otimes \dots \otimes \psi_n, x \vartheta_1 \otimes \dots \otimes \vartheta_n \rangle,$$

where $\psi_1, \psi_2, \dots, \psi_n, \vartheta_1, \vartheta_2, \dots, \vartheta_n$ are unit vectors in \mathbb{C}^d .

A measure for multiparticle entanglement

The maximal tensor norm: Let $\mathcal{H} := (\mathbb{C}^d)^{\otimes n}$, and let V denote the set of linear functionals on $\mathcal{B}(\mathcal{H})$ of the form

$$x \mapsto \langle \psi_1 \otimes \dots \otimes \psi_n, x \vartheta_1 \otimes \dots \otimes \vartheta_n \rangle,$$

where $\psi_1, \psi_2, \dots, \psi_n, \vartheta_1, \vartheta_2, \dots, \vartheta_n$ are unit vectors in \mathbb{C}^d .
Then a norm on the dual of $\mathcal{B}(\mathcal{H})$, is define by

$$\|\omega\|^V := \inf \left\{ \sum_{i=1}^k \lambda_i \mid \omega = \sum_{i=1}^k \lambda_i \nu_i, k \in \mathbb{N}, \lambda_1, \lambda_2, \dots, \lambda_k > 0, \nu_1, \nu_2, \dots, \nu_k \in V \right\};$$

A measure for multiparticle entanglement

The maximal tensor norm: Let $\mathcal{H} := (\mathbb{C}^d)^{\otimes n}$, and let V denote the set of linear functionals on $\mathcal{B}(\mathcal{H})$ of the form

$$x \mapsto \langle \psi_1 \otimes \dots \otimes \psi_n, x \vartheta_1 \otimes \dots \otimes \vartheta_n \rangle,$$

where $\psi_1, \psi_2, \dots, \psi_n, \vartheta_1, \vartheta_2, \dots, \vartheta_n$ are unit vectors in \mathbb{C}^d . Then a norm on the dual of $\mathcal{B}(\mathcal{H})$, is define by

$$\|\omega\|^V := \inf \left\{ \sum_{i=1}^k \lambda_i \mid \omega = \sum_{i=1}^k \lambda_i \nu_i, k \in \mathbb{N}, \lambda_1, \lambda_2, \dots, \lambda_k > 0, \nu_1, \nu_2, \dots, \nu_k \in V \right\};$$

When ρ is a state on $\mathcal{B}(\mathcal{H})$, we define its **entanglement** $E(\rho)$ by

$$E(\rho) := \|\rho\|^V.$$

Properties

Properties

The maximal tensor norm has all the required properties of an entanglement measure:

Properties

The maximal tensor norm has all the required properties of an entanglement measure:

- ▶ $E(\rho) \geq 1$ for all ρ ; $E(\rho) = 1$ iff ρ is separable;

Properties

The maximal tensor norm has all the required properties of an entanglement measure:

- ▶ $E(\rho) \geq 1$ for all ρ ; $E(\rho) = 1$ iff ρ is separable;
- ▶ $E((T_1 \otimes \dots \otimes T_n)\rho) \leq E(\rho)$ for all quantum operations T_1, T_2, \dots, T_n on $\mathcal{B}(\mathbb{C}^d)$;

Properties

The maximal tensor norm has all the required properties of an entanglement measure:

- ▶ $E(\rho) \geq 1$ for all ρ ; $E(\rho) = 1$ iff ρ is separable;
- ▶ $E((T_1 \otimes \dots \otimes T_n)\rho) \leq E(\rho)$ for all quantum operations T_1, T_2, \dots, T_n on $\mathcal{B}(\mathbb{C}^d)$;
- ▶ $E(\rho \otimes \vartheta) \leq E(\rho) \cdot E(\vartheta)$.

Properties

The maximal tensor norm has all the required properties of an entanglement measure:

- ▶ $E(\rho) \geq 1$ for all ρ ; $E(\rho) = 1$ iff ρ is separable;
- ▶ $E((T_1 \otimes \dots \otimes T_n)\rho) \leq E(\rho)$ for all quantum operations T_1, T_2, \dots, T_n on $\mathcal{B}(\mathbb{C}^d)$;
- ▶ $E(\rho \otimes \vartheta) \leq E(\rho) \cdot E(\vartheta)$.
(Here we would actually prefer equality!)

How entangled are the extremal completely symmetric states?

How entangled are the extremal completely symmetric states?

Theorem

Let $n, d \in \mathbb{N}$, and let Y denote an n -block Young frame with height $\leq d$. The entanglement of the completely symmetric state ρ_Y satisfies

$$\frac{n!}{d(Y) \cdot \text{Imm}_Y(G(\psi_{\max}))} \leq E(\rho_Y) \leq \frac{\sum_{\sigma \in S_n} |\chi_Y(\sigma)|}{\text{Imm}_Y(G(\psi_{\max}))},$$

where ψ_{\max} is that n -tuple of unit vectors in \mathbb{C}^d for which $\text{Imm}_Y(G(\psi))$ is maximal.

How entangled are the extremal completely symmetric states?

Theorem

Let $n, d \in \mathbb{N}$, and let Y denote an n -block Young frame with height $\leq d$. The entanglement of the completely symmetric state ρ_Y satisfies

$$\frac{n!}{d(Y) \cdot \text{Imm}_Y(G(\psi_{\max}))} \leq E(\rho_Y) \leq \frac{\sum_{\sigma \in S_n} |\chi_Y(\sigma)|}{\text{Imm}_Y(G(\psi_{\max}))},$$

where ψ_{\max} is that n -tuple of unit vectors in \mathbb{C}^d for which $\text{Imm}_Y(G(\psi))$ is maximal.

In particular, the antisymmetric state has entanglement

$$E(\rho_-) = n! .$$