

## **On charged fields with group symmetry and degeneracies of Verlinde's matrix $S^*$**

by

**Michael MÜGER<sup>1</sup>**

Dipartimento di Matematica, Università di Roma "Tor Vergata",  
Via della Ricerca Scientifica, 00133 Roma, Italy

Article received on 30 September 1997, revised 23 March 1998

---

**ABSTRACT.** – We consider the complete normal field net with compact symmetry group constructed by Doplicher and Roberts starting from a net of local observables in  $\geq 2 + 1$  spacetime dimensions and its set of localized (DHR) representations. We prove that the field net does not possess non-trivial DHR sectors, provided the observables have only finitely many sectors. Whereas the superselection structure in  $1 + 1$  dimensions typically does not arise from a group, the DR construction is applicable to 'degenerate sectors', the existence of which (in the rational case) is equivalent to non-invertibility of Verlinde's  $S$ -matrix. We prove Rehren's conjecture that the enlarged theory is non-degenerate, which implies that every degenerate theory is an 'orbifold' theory. Thus, the symmetry of a generic model 'factorizes' into a group part and a pure quantum part which still must be clarified. © Elsevier, Paris

**RÉSUMÉ.** – Nous étudions le réseau normal et complet d'algèbres des champs avec groupe de symétrie compact construit par Doplicher et Roberts à partir d'un réseau d'observables locales d'un espace temps de dimension supérieure ou égale à  $2 + 1$ , ainsi que l'ensemble des représentations localisés (DHR) qui lui est associé. Nous démontrons que le réseau

---

\* Partially supported by the Studienstiftung des deutschen Volkes and by CEE.

<sup>1</sup> E-mail: mueger@axp.mat.uniroma2.it.

des algèbres de champs ne possède pas de secteur DHR non-trivial, sous la condition que le réseau des algèbres d'observable n'ait qu'un nombre fini de secteur de superselections. Alors qu'en dimension  $1 + 1$  la structure des secteurs de superselection n'est pas déterminée par un groupe, la construction de D.R. s'applique aux 'secteurs dégénérés' dont l'existence, dans le cas rationnel, est équivalente à la non inversibilité de la matrice  $S$  de Verlinde. Nous démontrons aussi la conjecture de Rehren selon laquelle la théorie élargie est non-dégénérée, impliquant que toute théorie dégénérée est une théorie 'orbifold'. Aussi, la symétrie d'un modèle générique se décompose en une contribution venant d'un groupe et une autre purement quantique dont la structure nécessite encore d'être élucidée. © Elsevier, Paris

---

## 1. INTRODUCTION

A few years ago a long-standing problem in local quantum physics [28] (algebraic quantum field theory) was solved in [22], where the conjecture [10,13] was proved that the superselection structure of the local observables can always be described in terms of a compact group. This group (gauge group of the first kind) acts by automorphisms on a net of field algebras which generate the charged sectors from the vacuum and obey normal Bose and Fermi commutation relations. From the mathematical point of view this amounts to a new duality theory for compact groups [21] which considerably improves on the old Tannaka–Krein theory. These results rely on a remarkable chain of arguments [17–20] which we cannot review here. We refer to the first two sections of [22] for a relatively non-technical overview of the construction and restrict ourselves to a short introduction to the problem in order to set the stage for our considerations.

Our starting point is a net of local observables, i.e., an inclusion preserving map  $\mathcal{O} \mapsto \mathcal{A}(\mathcal{O})$  which assigns to each double cone  $\mathcal{O}$  (the set of these is denoted by  $\mathcal{K}$ ) in spacetime the algebra of observables measurable in  $\mathcal{O}$ . More specifically, identifying the abstract local algebras with their images in a faithful vacuum representation  $\pi_0$ , we assume the  $\mathcal{A}(\mathcal{O})$  to be von Neumann algebras acting on the Hilbert space  $\mathcal{H}_0$ . The  $C^*$ -algebra  $\mathcal{A}$  generated by all  $\mathcal{A}(\mathcal{O})$  is called the quasilocal algebra. As usual the property of Einstein causality (if  $\mathcal{O}_1, \mathcal{O}_2$  are mutually

spacelike double cones then  $\mathcal{A}(\mathcal{O}_1)$  and  $\mathcal{A}(\mathcal{O}_2)$  commute elementwise) is strengthened by requiring Haag duality

$$\mathcal{A}(\mathcal{O})' = \mathcal{A}(\mathcal{O}')^- \quad \forall \mathcal{O} \in \mathcal{K}, \quad (1.1)$$

where  $\mathcal{A}(\mathcal{O}')$  is the  $C^*$ -algebra generated by  $\mathcal{A}(\tilde{\mathcal{O}})$ ,  $\mathcal{K} \ni \tilde{\mathcal{O}} \subset \mathcal{O}'$ . Typically one requires Poincaré or conformal covariance but these properties will play no essential role for our considerations, apart from their being used to derive the Property B (cf. Section 2.1 below) which is needed for the analysis of the superselection structure.

We restrict our attention to superselection sectors which are localizable in arbitrary double cones, i.e., representations  $\pi$  of the quasilocal algebra  $\mathcal{A}$  satisfying the DHR criterion [12,14]:

$$\pi \upharpoonright \mathcal{A}(\mathcal{O}') \cong \pi_0 \upharpoonright \mathcal{A}(\mathcal{O}') \quad \forall \mathcal{O} \in \mathcal{K}. \quad (1.2)$$

These representations are called locally generated since they are indistinguishable from the vacuum when restricted to the spacelike complement of a double cone. Given a representation of this type, Haag duality implies [12] for any double cone  $\mathcal{O}$  the existence of a unital endomorphism of  $\mathcal{A}$  which is localized in  $\mathcal{O}$  (in the sense that  $\rho(A) = A \quad \forall A \in \mathcal{A}(\mathcal{O}')$ ) such that  $\pi \cong \pi_0 \circ \rho \equiv \rho$ . This is an important fact since endomorphisms can be composed, thereby defining a composition rule for this class of representations. Whereas (non-surjective) endomorphisms are not invertible, there are left inverses, i.e., (completely) positive linear maps  $\phi_\rho : \mathcal{A} \rightarrow \mathcal{A}$  such that

$$\phi_\rho(\rho(A)B\rho(C)) = A\phi_\rho(B)C,$$

in particular  $\phi_\rho \circ \rho = id$ . Localized endomorphisms obtained from DHR representations are transportable, i.e., given  $\rho$  there is an equivalent morphism localized in  $\mathcal{O}$  for every  $\mathcal{O} \in \mathcal{K}$ . Furthermore, given two localized endomorphisms, one can construct operators  $\varepsilon(\rho_1, \rho_2)$  which intertwine  $\rho_1\rho_2$  and  $\rho_2\rho_1$  and thereby formalize the notion of particle interchange (whence the name statistics operators). For  $\rho$  an irreducible morphism,  $\phi_\rho(\varepsilon(\rho, \rho)) = \lambda_\rho \mathbf{1}$  gives rise via polar decomposition  $\lambda_\rho = \omega_\rho/d_\rho$  to a phase and a positive number. From here on the analysis depends crucially on the number of spacetime dimensions. In  $\geq 2 + 1$  dimensions [12,14] the statistics operators  $\varepsilon(\rho_1, \rho_2)$  are uniquely defined and satisfy  $\varepsilon(\rho, \rho)^2 = \mathbf{1}$  such that one obtains, for each morphism  $\rho$ , a unitary representation of the permutation group in  $\mathcal{A}$  via  $\sigma_i \mapsto \rho^{i-1}(\varepsilon(\rho, \rho))$ . Furthermore, the statistics phase and

dimension satisfy  $\omega_\rho = \pm 1$  and  $d_\rho \in \mathbb{N} \cup \{\infty\}$ . The statistics phase  $\omega_\rho$  distinguishes representations with bosonic and fermionic character, and the statistical dimension  $d(\rho)$  measures the degree of parastatistics. Ignoring morphisms with infinite dimension, which are considered pathological, we denote by  $\Delta$  the semigroup of all transportable localized morphisms with finite statistics.

The analysis which was sketched above was motivated by the preliminary investigations conducted in [10]. There the starting point is a net of field algebras  $\mathcal{O} \mapsto \mathcal{F}(\mathcal{O})$  acted upon by a compact group  $G$  of inner symmetries (gauge group of the first kind):

$$\alpha_g(\mathcal{F}(\mathcal{O})) = \mathcal{F}(\mathcal{O}). \tag{1.3}$$

The field algebra acts irreducibly on a vacuum Hilbert space  $\mathcal{H}$  and the gauge group is unbroken, i.e., represented by unitary operators  $U(g)$  in a strongly continuous way:  $\alpha_g(F) = Ad U(g)(F)$ . (Compactness of  $G$  need in fact not be postulated, as it follows by [16, Theorem 3.1] if the field net satisfies the split property.)

The field net is supposed to fulfill Bose–Fermi commutation relations, i.e., any local operator decomposes into a bosonic and a fermionic part  $F = F_+ + F_-$  such that for spacelike separated  $F$  and  $G$  we have

$$[F_+, G_+] = [F_+, G_-] = [F_-, G_+] = [F_-, G_-] = 0. \tag{1.4}$$

The above decomposition is achieved by

$$F_\pm = \frac{1}{2}(F \pm \alpha_k(F)), \tag{1.5}$$

where  $k$  is an element of order 2 in the center of the group  $G$ .  $V \equiv U_k$  is the unitary operator which acts trivially on the space of bosonic vectors and like  $-1$  on the fermionic ones. To formulate this locality requirement in a way more convenient for later purposes we introduce the twist operation  $F^t = ZFZ^*$  where

$$Z = \frac{1 + iV}{1 - i} \quad (\Rightarrow Z^2 = V), \tag{1.6}$$

which leads to  $ZF_+Z^* = F_+$ ,  $ZF_-Z^* = iVF_-$ , implying  $[F, G^t] = 0$ . The (twisted) locality postulate (1.4) can now be stated simply as

$$\mathcal{F}(\mathcal{O})^t \subset \mathcal{F}(\mathcal{O}'). \tag{1.7}$$

In analogy to the bosonic case, this can be strengthened to twisted duality:

$$\mathcal{F}(\mathcal{O})' = \mathcal{F}(\mathcal{O}'). \tag{1.8}$$

The observables are now defined as the fixpoints under the action of  $G$ :

$$\mathcal{A}(\mathcal{O}) = \mathcal{F}(\mathcal{O})^G = \mathcal{F}(\mathcal{O}) \cap U(G)'. \tag{1.9}$$

The Hilbert space  $\mathcal{H}$  decomposes as follows:

$$\mathcal{H} = \bigoplus_{\xi \in \widehat{G}} \mathcal{H}_\xi \otimes \mathbb{C}^{d_\xi}, \tag{1.10}$$

where  $\xi$  runs through the equivalence classes of finite dimensional continuous unitary representations of  $G$  and  $d_\xi$  is the dimension of  $\xi$ . The observables and the group  $G$  act reducibly according to

$$\begin{aligned} A &= \bigoplus_{\xi \in \widehat{G}} \pi_\xi(A) \otimes \mathbf{1}, \\ U(g) &= \bigoplus_{\xi \in \widehat{G}} \mathbf{1}_{\mathcal{H}_\xi} \otimes U_\xi(g), \end{aligned} \tag{1.11}$$

where  $\pi_\xi$  and  $U_\xi$  are irreducible representations of  $\mathcal{A}$  and  $G$ , respectively. As a consequence of twisted duality for the fields, the restriction of the observables  $\mathcal{A}$  to a simple sector (subspace  $\mathcal{H}_\xi$  with  $d_\xi = 1$ ), in particular the vacuum sector, satisfies Haag duality. Since the unitary representation of the Poincaré group commutes with  $G$ , the restriction of  $\mathcal{A}$  to  $\mathcal{H}_0$  satisfies all requirements for a net of observables in the vacuum representation in the above sense. As shown in [10], the irreducible representations of  $\mathcal{A}$  in the charged sectors are globally inequivalent but strongly locally equivalent to each other (i.e.,  $\pi_1 \upharpoonright \mathcal{A}(\mathcal{O}') \cong \pi_2 \upharpoonright \mathcal{A}(\mathcal{O}')$ ), in particular they satisfy the DHR criterion. Obviously it is not necessarily true that the decomposition (1.11) contains all equivalence classes of DHR representations (take  $\mathcal{F} = \mathcal{A}$ ,  $\mathcal{H} = \mathcal{H}_0$ ,  $G = \{e\}$ ). This completeness is true, however, if the field net  $\mathcal{F}$  has trivial representation theory (equivalently ‘quasitrivial 1-cohomology’), see [43]. It was conjectured in [13] that every net of observables arises as a fixpoint net such that the representation of  $\mathcal{A}$  on  $\mathcal{H}$  contains all sectors, which furthermore means that the tensor category of DHR sectors with finite statistics is isomorphic to the representation category of a compact

group  $G$ . Under the restriction that all transportable localized morphisms are automorphisms, which is equivalent to  $G$  being abelian, this was proved in [11]. After the early works [13,41] the final proof in complete generality [17–22] turned out to be quite difficult, which is perhaps not too surprising in view of the non-triviality of the result.

In the next section we prove a few complementary results concerning the DR-construction in  $\geq 2 + 1$  dimensions. It is natural to conjecture that the DR field net does not possess localized superselection sectors provided it is complete, i.e., contains charged fields generating all DHR sectors (with finite statistics) of the observables. Whereas, at first sight, this may appear to be an obvious consequence of the uniqueness result [22] for the complete normal field net we have unfortunately been able to give a proof only for the case of a finite gauge group, i.e., for rational theories. Under the same assumption we show that the complete field net can also be obtained by applying the DR construction to an intermediate, i.e., incomplete field net. Whereas in higher dimensions the restriction to finite gauge groups is quite unsatisfactory, our results have a useful application to the low dimensional case to which we now turn.

In  $1 + 1$  dimensions there are in particular two interesting classes of models. The first consists of purely massive models, many of these being integrable. Concerning these it has been shown recently [37] that they do not have DHR sectors at all as long as one insists on the assumption of Haag duality. As to conformally covariant models, which constitute the other class of interest, the situation is quite different in that it has been shown [5] that positive-energy representations are necessarily of the DHR type due to local normality and compactness of the spacetime. It is particularly this class which we have in mind in our  $2d$  considerations, but the conformal covariance will play no role. Whereas in  $\geq 2 + 1$  dimensions one has  $\varepsilon(\rho_2, \rho_1)^* = \varepsilon(\rho_1, \rho_2)$ , in  $1 + 1$  dimensions these statistics operators are a priori different intertwiners between  $\rho_1\rho_2$  and  $\rho_2\rho_1$ . This phenomenon accounts for the occurrence of braid group statistics and provides the motivation for defining the *monodromy operators*:

$$\varepsilon_M(\rho_1, \rho_2) = \varepsilon(\rho_1, \rho_2) \varepsilon(\rho_2, \rho_1), \quad (1.12)$$

which measure the deviation from permutation group statistics. An irreducible morphism  $\rho$  is said to be degenerate if  $\varepsilon_M(\rho, \sigma) = \mathbf{1}$  for all  $\sigma$ . Given two irreducible morphisms  $\rho_i, \rho_j$  one obtains the  $\mathbb{C}$ -number

valued *statistics character* [39] via

$$Y_{ij}\mathbf{1} = d_i d_j \phi_j (\varepsilon_M(\rho_i, \rho_j)^*). \tag{1.13}$$

(Here  $\phi_j$  is the left inverse of  $\rho_j$  and the factor  $d_i d_j$  has been introduced for later convenience.) The numbers  $Y_{ij}$  depend only on the sectors, such that the matrix  $(Y_{ij})$  can be considered as indexed by the set of equivalence classes of irreducible sectors. The matrix  $Y$  satisfies the following identities:

$$Y_{0i} = Y_{i0} = d_i, \tag{1.14}$$

$$Y_{ij} = Y_{ji} = Y_{i\bar{j}}^* = Y_{\bar{i}j}, \tag{1.15}$$

$$Y_{ij} = \sum_k N_{ij}^k \frac{\omega_i \omega_j}{\omega_k} d_k, \tag{1.16}$$

$$\frac{1}{d_j} Y_{ij} Y_{kj} = \sum_m N_{ik}^m Y_{mj}. \tag{1.17}$$

Here  $[\rho_{\bar{i}}]$  is the conjugate morphism of  $[\rho_i]$  and  $N_{ij}^k \in \mathbb{N}_0$  is the multiplicity of  $[\rho_k]$  in the decomposition of  $[\rho_i \rho_j]$  into irreducible morphisms. The matrix of statistics characters is of particular interest if the theory is *rational*, i.e., has only a finite number of inequivalent irreducible representations. Then, as proved by Rehren [39], the matrix  $Y$  is invertible iff there is no degenerate morphism besides the trivial one which corresponds to the vacuum representation. In the non-degenerate case the number  $\sigma = \sum_i d_i^2 \omega_i^{-1}$  satisfies  $|\sigma|^2 = \sum_i d_i^2$  and the matrices

$$S = |\sigma|^{-1} Y, \quad T = \left( \frac{\sigma}{|\sigma|} \right)^{1/3} \text{Diag}(\omega_i) \tag{1.18}$$

are unitary and satisfy the relations

$$S^2 = (ST)^3 = C, \quad TC = CT, \tag{1.19}$$

where  $C_{ij} = \delta_{i,\bar{j}}$  is the charge conjugation matrix. That is,  $S$  and  $T$  constitute a representation of the modular group  $SL(2, \mathbb{Z})$ . Furthermore, the ‘fusion coefficients’  $N_{ij}^k$  are given by the Verlinde relation

$$N_{ij}^k = \sum_m \frac{S_{im} S_{jm} S_{km}^*}{S_{0m}}. \tag{1.20}$$

As was emphasized in [39], these relations hold independently of conformal covariance in every (non-degenerate) two dimensional theory with finitely many DHR sectors. This is remarkable, since Eq. (1.20) first appeared [47] in the context of conformal quantum field theory on the torus, where the  $S$ -matrix by definition has the additional property of describing the behavior of the conformal characters  $tr_{\pi_i} e^{-\tau L_0}$  under the inversion  $\tau \rightarrow -1/\tau$ .

Eqs. (1.19) and (1.20) do not hold if the matrix  $Y$  is not invertible, i.e., when there are degenerate sectors. One can show that the set of degenerate sectors is stable under composition and reduction into irreducibles (Lemma 3.4). It thus constitutes a closed subcategory of the category of DHR endomorphisms to which one can apply the DR construction of charged fields. In Section 4 we will prove Rehren's conjecture in [39] that the resulting 'field' net has no degenerate sectors. Furthermore, we will prove that the enlarged theory is rational, provided that the original one is. These results imply that the above Verlinde-type analysis is in fact applicable to  $\mathcal{F}$ .

## 2. ON THE RECONSTRUCTION OF FIELDS FROM OBSERVABLES

Our first aim in this section will be to prove the intuitively reasonable fact that a complete field net associated (in  $\geq 2 + 1$  dimensions) with a net of observables does not possess localized superselection sectors. For technical reasons we have been able to give a proof only for rational theories. This result, which may not be too useful in itself, will after some preparations be the basis of our proof of a conjecture by Rehren (Theorem 3.6). Furthermore, we show that the construction of the complete field net 'can be done in steps', that is, one also obtains the complete field net by applying the DR construction to an intermediate, thus incomplete, field net and its DHR sectors. For the sake of simplicity we defer the treatment of the general case for a while and begin with the purely bosonic case.

### 2.1. Absence of DHR sectors of the complete field net: Bose case

The superselection theory of a net of observables is called purely bosonic if all DHR sectors have statistics phase  $+1$ . In this case the charged fields which generate these sectors from the vacuum are local and the fields associated with different sectors can be chosen to be relatively



local. Then the Doplicher–Roberts construction [22] gives rise to a local field net  $\mathcal{F}$ , which in addition satisfies Haag duality. Thus it makes sense to consider the DHR sectors of  $\mathcal{F}$  and to apply the DR construction to these. (In analogy to [12,14] one requires  $\mathcal{F}$  to satisfy the technical ‘property B’ [12], which can be derived [2] from standard assumptions, in particular positive energy. Since a DR field net is Poincaré covariant with positive energy [22, Section 6], provided this is true for the vacuum sector and the DHR representations of the observables, we may take the property B for granted also for  $\mathcal{F}$ .)

We cite the following definitions from [22]:

**DEFINITION 2.1.** – *Given a net  $\mathcal{A}$  of observables and a vacuum representation  $\pi_0$ , a normal field system with gauge symmetry,  $\{\pi, \mathcal{F}, G\}$ , consists of a representation  $\pi$  of  $\mathcal{A}$  on a Hilbert space  $\mathcal{H}$  containing  $\pi_0$  as a subrepresentation on  $\mathcal{H}_0 \subset \mathcal{H}$ , a compact group  $G$  of unitaries on  $\mathcal{H}$  leaving  $\mathcal{H}_0$  pointwise fixed and a net  $\mathcal{O} \mapsto \mathcal{F}(\mathcal{O}) \subset \mathcal{B}(\mathcal{H})$  of von Neumann algebras such that:*

- ( $\alpha$ ) *the  $g \in G$  induce automorphisms  $\alpha_g$  of  $\mathcal{F}(\mathcal{O})$ ,  $\mathcal{O} \in \mathcal{K}$  with  $\pi(\mathcal{A}(\mathcal{O}))$  as fixed-point algebra,*
- ( $\beta$ ) *the field net  $\mathcal{F}$  is irreducible,*
- ( $\gamma$ )  *$\mathcal{H}_0$  is cyclic for  $\mathcal{F}(\mathcal{O}) \forall \mathcal{O} \in \mathcal{K}$ ,*
- ( $\delta$ ) *there is an element  $k$  in the center of  $G$  with  $k^2 = e$  such that the net  $\mathcal{F}$  obeys graded local commutativity for the  $\mathbb{Z}_2$ -grading defined by  $k$ , cf. (1.4) and (1.5).*

**DEFINITION 2.2.** – *A field system with gauge symmetry  $\{\pi, \mathcal{F}, G\}$  is complete if each equivalence class of irreducible representations of  $\mathcal{A}$  satisfying (1.2) and having finite statistics is realized as a subrepresentation of  $\pi$ , i.e.,  $\pi$  describes all relevant superselection sectors.*

For a given net of observables  $\mathcal{A}$  we denote by  $\Delta$  the set of all transportable localized morphisms with finite statistics. Let  $\Gamma$  be a closed semigroup of localized *bosonic* endomorphisms and let  $\mathcal{F}$  be the associated local field net. Now let  $\Sigma$  be a closed semigroup of localized endomorphisms of  $\mathcal{F}$ . After iterating the DR construction again we are faced with the following situation. There are three nets  $\mathcal{A}$ ,  $\mathcal{F}$ ,  $\tilde{\mathcal{F}}$  acting faithfully and irreducibly on the Hilbert spaces  $\mathcal{H}_0 \subset \mathcal{H} \subset \tilde{\mathcal{H}}$ , respectively, such that Haag duality holds (twisted duality in the case of  $\tilde{\mathcal{F}}$ ). The nets  $\tilde{\mathcal{F}}$  and  $\mathcal{F}$  are normal field nets with respect to the nets  $\mathcal{F}$  and  $\mathcal{A}$ , respectively, in the sense of Definition 2.1. Thus there are representations  $\pi$  of  $\mathcal{A}$  on  $\mathcal{H}$  and  $\tilde{\pi}$  of  $\mathcal{F}$  on  $\tilde{\mathcal{H}}$ , respectively, such

that  $\tilde{\pi} \circ \pi(\mathcal{A}) \subset \tilde{\pi}(\mathcal{F}) \subset \tilde{\mathcal{F}}$ . Furthermore, there are strongly compact groups  $G$  and  $\tilde{G}$  of unitaries on  $\mathcal{H}$  and  $\tilde{\mathcal{H}}$ , respectively, acting as local symmetries on  $\mathcal{F}$  and  $\tilde{\mathcal{F}}$ , respectively, such that  $\mathcal{F}(\mathcal{O})^G = \pi(\mathcal{A}(\mathcal{O}))$ ,  $\mathcal{O} \in \mathcal{K}$ , and  $\tilde{\mathcal{F}}(\mathcal{O})^{\tilde{G}} = \tilde{\pi}(\mathcal{F}(\mathcal{O}))$ ,  $\mathcal{O} \in \mathcal{K}$ . The following result is crucial:

**PROPOSITION 2.3.** – *Let the theory  $\mathcal{A}$  be rational (equivalently, let  $G$  be finite). Then the net  $\tilde{\mathcal{F}}$  is a normal field net with respect to the observables  $\mathcal{A}$ . In particular, there is a strongly compact group  $\overline{G}$  of unitaries on  $\tilde{\mathcal{H}}$  containing  $\tilde{G}$  as a closed normal subgroup.  $\overline{G}$  implements local symmetries of  $\tilde{\mathcal{F}}$  such that  $\tilde{\mathcal{F}}(\mathcal{O})^{\overline{G}} = \tilde{\pi} \circ \pi(\mathcal{A}(\mathcal{O}))$ .*

*Proof.* – Let  $\overline{G}$  be the group of unitaries on  $\tilde{\mathcal{H}}$  implementing local symmetries of  $\tilde{\mathcal{F}}$  which leave  $\mathcal{A}$  pointwise and the algebras  $\mathcal{F}(\mathcal{O})$ ,  $\mathcal{O} \in \mathcal{K}$  globally stable. Clearly,  $\overline{G}$  is strongly closed and contains  $\tilde{G}$  as a closed normal subgroup. We can now apply Proposition 3.1 of [7] to the effect that every element of  $G$  extends to a unitarily implemented local symmetry of  $\tilde{\mathcal{F}}$ , thus an element of  $\overline{G}$ , such that there is a short exact sequence

$$\mathbf{1} \rightarrow \tilde{G} \rightarrow \overline{G} \rightarrow G \rightarrow \mathbf{1}. \quad (2.1)$$

By assumption,  $\tilde{G}$  is known to be compact in the strong topology which, of course, coincides with the topology induced from  $\overline{G}$ . The group  $G$  being finite it is clearly compact with respect to any topology. Compactness of  $\tilde{G}$  and  $G$  imply compactness of  $\overline{G}$  (cf., e.g., [29, Theorem 5.25]).

It remains to prove the requirements  $(\beta)$ – $(\delta)$  of Definition 2.1. Now,  $(\beta)$  and  $(\delta)$  are automatically true by [22, Theorem 3.5]. Finally,  $(\gamma)$ , viz. the cyclicity of  $\mathcal{H}_0$  for  $\tilde{\mathcal{F}}(\mathcal{O})$ ,  $\mathcal{O} \in \mathcal{K}$  is also easy: in application to  $\mathcal{H}_0$ ,  $\tilde{\pi}(\mathcal{F}(\mathcal{O})) \subset \tilde{\mathcal{F}}(\mathcal{O})$  gives a dense subset of  $\mathcal{H}$ , the image of which under the action of the charged (with respect to  $\mathcal{F}$ ) fields in  $\mathcal{F}$  is dense in  $\tilde{\mathcal{H}}$ .  $\square$

*Remark.* – As to the general case of infinite  $G$  we note that,  $\tilde{G}$  being compact,  $\overline{G}$  is (locally) compact iff  $G = \overline{G}/\tilde{G}$  is (locally) compact in the quotient topology. It is easy to show that the identical map from  $G$  with the quotient topology to  $G$  with the strong topology induced from  $\mathcal{B}(\mathcal{H})$  is continuous. Since we know that  $G$  is compact with respect to the latter and since both topologies are Hausdorff,  $G$  is compact with respect to the former (and thus  $\overline{G}$  is compact) iff the identical map is open. This would follow from an open mapping theorem [29, Theorem 5.29] if we could prove that the  $G$  is locally compact and second countable with

the quotient topology. Clearly this idea can work only if the observables have at most countably many sectors. We hope to return to this problem in another paper.

We are now prepared to prove the absence of DHR sectors of the field net. Let  $\Gamma = \Delta$ , the set of all transportable localized morphisms of  $\mathcal{F}$  with finite statistics. Using Proposition 2.3 we easily prove the following:

**THEOREM 2.4.** – *The complete (local) field net  $\mathcal{F}$  associated with a purely bosonic rational theory has no DHR sectors with finite statistics.*

*Proof.* – Assuming the converse, the above proposition gives us a field net  $\tilde{\mathcal{F}}$  on a larger Hilbert space  $\tilde{\mathcal{H}}$ , which obviously is also complete, since the representation  $\pi$  of  $\mathcal{A}$  on  $\mathcal{H}$  is a subrepresentation of  $\tilde{\pi} \circ \pi$ . Thus, by [22, Theorem 3.5] both field systems are equivalent, that is, there is a unitary operator  $W : \mathcal{H} \rightarrow \tilde{\mathcal{H}}$  such that

$$W\pi(A) = \tilde{\pi} \circ \pi(A)W \quad \forall A \in \mathcal{A}$$

etc. In view of the decomposition

$$\pi = \bigoplus_{\xi \in \hat{G}} d_{\xi} \pi_{\xi}, \tag{2.2}$$

where the irreducible representations  $\pi_{\xi}$  are mutually inequivalent, and similarly for  $\tilde{\pi} \circ \pi$ ,  $\pi$  and  $\tilde{\pi}$  can be unitarily equivalent only if  $G = \tilde{G}$  and thus  $\mathcal{F} = \tilde{\mathcal{F}}$ .  $\square$

*Remark.* – After this paper was essentially completed I learned that this result (with the same restriction to finite groups) has been obtained about two years ago by R. Conti [8].

We have thus, in the purely bosonic case, reached our first goal. Before we turn to the general situation we show that the construction of the complete field net ‘can be done in steps’, that is, one also obtains the complete field net by applying the DR construction to an intermediate field net and its DHR sectors, again assuming that the intermediate net is local (this is not required for the complete field net).

## 2.2. Stepwise construction of the complete field net: Bose case

The following lemma is more or less obvious and is stated here since it does not appear explicitly in [20,22].

LEMMA 2.5. – Let  $\Gamma_1, \Gamma_2$  be subsemigroups of  $\Delta$  which are both closed under direct sums, subobjects and conjugates and let  $\mathcal{F}_i$ ,  $i = 1, 2$ , be the associated normal field nets on the Hilbert spaces  $\mathcal{H}_i$  with symmetry groups  $G_i$  and  $\pi_i$  the representations of  $\mathcal{A}$ . If  $\Gamma_1 \subset \Gamma_2$  then there is an isometry  $V : \mathcal{H}_1 \rightarrow \mathcal{H}_2$  such that

$$V\pi_1(A) = \pi_2(A)V, \quad A \in \mathcal{A}, \quad (2.3)$$

$$VG_1V^* = G_2E, \quad (2.4)$$

$$V\mathcal{F}_1V^* = (\mathcal{F}_2 \cap \{E\}')E, \quad (2.5)$$

where  $E = VV^*$ . Furthermore, there is a closed normal subgroup  $N$  of  $G_2$  such that  $E$  is the projection onto the subspace of  $N$ -invariant vectors in  $\mathcal{H}_2$  and  $\{\pi_1, G_1, \mathcal{F}_1\}$  is equivalent to  $\{\pi_2^N, G_2/N, \mathcal{F}_2^N\}$ .

*Proof.* – As usual, the field theory  $\mathcal{F}_2$  is constructed by applying [22, Corollary 6] to the quadruple  $(\mathcal{A}, \Delta_2, \varepsilon, \pi_0)$  and by defining  $\mathcal{F}(\mathcal{O})$  to be the von Neumann algebra on  $\mathcal{H}_2$  generated by the Hilbert spaces  $H_\rho$ ,  $\rho \in \Delta_2(\mathcal{O})$ . Let  $E$  be the projection  $[\mathcal{B}_1\mathcal{H}_0]$  where  $\mathcal{B}_1$  is the  $C^*$ -algebra generated by  $H_\rho$ ,  $\rho \in \Delta_1$ . Trivially,  $\mathcal{B}_1$  maps  $E\mathcal{H}_2$  into itself.  $\mathcal{B}_1$  is stable under  $G_2$  as each of the Hilbert spaces  $H_\rho$  is. This implies that  $G_2$  leaves  $E\mathcal{H}_2$  stable. Restricting  $\mathcal{B}_1$  and  $G_2$  to  $E\mathcal{H}_2$  one obtains the system  $(E\mathcal{H}_2, E\pi_2(\cdot)E, EU_2E, \rho \in \Delta_1 \rightarrow EH_\rho E)$  which satisfies a) to g) of [22, Corollary 6.2]. With the exception of g) all of these are trivially obtained as restrictions. Property g) follows by appealing to [19, Lemma 2.4]. We can thus conclude from the uniqueness result of [22, Corollary 6.2] that  $(E\mathcal{H}_2, E\pi_2E, EU_2E, \rho \in \Delta_1 \rightarrow H_\rho)$  is equivalent to the system  $(\mathcal{H}_1, \pi_1, U_1, \rho \in \Delta_1 \rightarrow H_\rho)$  obtained from the quadruple  $(\mathcal{A}, \Delta_2, \varepsilon, \pi_0)$ , that is, there is a unitary  $V$  from  $\mathcal{H}_1$  to  $E\mathcal{H}_2$  such that  $V\pi_1(A) = \pi_2(A)V$ ,  $V\mathcal{F}_1 = \mathcal{B}_1V$ ,  $VU_1 = U_2V$ . Interpreting  $V$  as an isometry mapping  $\mathcal{H}_1$  into  $\mathcal{H}_2$  we have (2.3)–(2.5). The rest follows from [22, Proposition 3.17].  $\square$

LEMMA 2.6. – Let  $\mathcal{O} \mapsto \mathcal{F}(\mathcal{O})$  be the field net associated to a subsemigroup  $\Gamma$  of  $\Delta$ , closed under direct sums, subobjects and conjugates. Then every localized endomorphism  $\eta \in \Delta$  of  $\mathcal{A}$  extends to an endomorphism  $\tilde{\eta}$  of  $\mathcal{F}$  commuting with the action of the gauge group. If  $\eta$  is localized in  $\mathcal{O}$  the same holds for  $\tilde{\eta}$ .

*Remark.* – This result is of interest only if  $\eta \notin \Gamma$ . Otherwise we already know that  $\eta$  extends to an inner endomorphism of  $\mathcal{F}$  by definition of the field algebra.

*Proof.* – By the preceding result we know that the field net  $\mathcal{F} = \mathcal{F}_\Gamma$  is equivalent to a subnet of the complete field net  $\overline{\mathcal{F}} = \mathcal{F}_\Delta$ . We identify  $\mathcal{F}$  with this subnet. By construction every localized endomorphism  $\eta \in \Delta(\mathcal{O})$  of  $\mathcal{A}$  extends to an inner endomorphism of  $\overline{\mathcal{F}}$ , i.e., there is a multiplet of isometries  $\psi_i \in \overline{\mathcal{F}}(\mathcal{O})$ ,  $i = 1, \dots, d$ , satisfying  $\sum_i \psi_i \psi_i^* = \mathbf{1}$ ,  $\psi_i^* \psi_j = \delta_{i,j} \mathbf{1}$  such that  $\hat{\eta} \circ \pi(A) = \pi \circ \eta(A)$  where

$$\hat{\eta}(\cdot) = \sum_i \psi_i \cdot \psi_i^*. \tag{2.6}$$

Since  $\hat{\eta}$  commutes with the action of  $G$ , it is easy to verify that  $\hat{\eta}$  leaves  $\mathcal{F} = \overline{\mathcal{F}}^N$  stable and thus restricts to an endomorphism of  $\mathcal{F}$  which extends  $\eta$ . This extension is not necessarily local, for  $\hat{\eta}(F) = -F$  if  $F$  is a fermionic operator localized spacelike to  $\mathcal{O}$  and  $\eta$  is a fermionic endomorphism. This defect is easily remedied by defining

$$\tilde{\eta} = \begin{cases} \hat{\eta} & \text{if } \omega(\eta) = 1, \\ Ad V \circ \hat{\eta} & \text{if } \omega(\eta) = -1. \end{cases} \tag{2.7}$$

Clearly,  $\tilde{\eta}$  has the desired localization properties and coincides with  $\eta$  on  $\mathcal{A}$ . Transportability of  $\tilde{\eta}$  is automatic as  $W \in (\eta, \eta')$  implies  $\pi(W) \in (\tilde{\eta}, \tilde{\eta}')$ . Finally the statistical dimensions of  $\eta$  and  $\tilde{\eta}$  coincide as is seen using, e.g., the arguments in [32].  $\square$

*Remark.* – The preceding lemmas do not depend on the restriction to bosonic families  $\Gamma$  of endomorphisms or on the finiteness of the gauge group.

LEMMA 2.7. – *Let  $\mathcal{A}$  be rational, let  $\Gamma$  be a semigroup of bosonic endomorphisms and let  $\mathcal{F}$  be the associated (incomplete) local field net. Let  $\Sigma$  be the semigroup of all localized endomorphisms of  $\mathcal{F}$ . Then the associated DR-field net  $\tilde{\mathcal{F}}$  is a complete field net with respect to  $\mathcal{A}$ .*

*Proof.* – Let  $\eta$  be a localized endomorphism of  $\mathcal{A}$ . By the preceding lemma, there is an extension (typically reducible) to a localized endomorphism  $\tilde{\eta}$  of  $\tilde{\mathcal{F}}$ . By Proposition 2.3,  $\tilde{\mathcal{F}}$  is a normal field net for  $\mathcal{A}$ . By completeness of  $\tilde{\mathcal{F}}$  with respect to endomorphisms of  $\mathcal{F}$ ,  $\tilde{\eta}$  is implemented by a Hilbert space in  $\tilde{\mathcal{F}}$  and there is a subspace  $\mathcal{H}_{\tilde{\eta}}$  of  $\mathcal{H}$  such that  $\tilde{\pi} \upharpoonright \mathcal{H}_{\tilde{\eta}} \cong \tilde{\eta}$  as a representation of  $\mathcal{F}$ . Restricting to  $\mathcal{A}$  and choosing an irreducible subspace  $\mathcal{H}_\eta$  we have  $\pi_\Sigma \upharpoonright \mathcal{H}_\eta \cong \pi_0 \circ \eta$ . Thus  $\tilde{\mathcal{F}}$  is a complete field net for  $\mathcal{A}$ .  $\square$

**THEOREM 2.8.** – *Let  $\mathcal{A}$  be a rational net of observables and let  $\Gamma$  be a bosonic subsemigroup of  $\Delta$  with the associated field net  $\mathcal{F}_\Gamma$ . Then the complete normal field net  $\mathcal{F}_{\Gamma, \Sigma}$  obtained from the net  $\mathcal{F}_\Gamma$  and its semigroup  $\Sigma$  of all localized endomorphisms is equivalent to the complete normal field net  $\mathcal{F}_\Delta$ . In particular the group  $\bar{G}$  obtained in Lemma 2.3 is isomorphic to the group belonging to  $\mathcal{F}_\Delta$ .*

*Proof.* – By Lemmas 2.3 and 2.7,  $\mathcal{F}_{\Gamma, \Sigma}$  is a complete normal field net for  $\mathcal{A}$ . The same trivially holding for  $\mathcal{F}_\Delta$ , we are done since two such nets are isomorphic by [22, Theorem 3.5].  $\square$

### 2.3. General case, including fermions

In the attempt to prove generalizations of Theorem 2.4 for theories possessing fermionic sectors and of Theorem 2.8 for fermionic intermediate nets  $\mathcal{F}$  we are faced with the problem that it is not entirely obvious what these generalizations should be. We would like to show the representation theory of a complete normal field net, which is now assumed to comprise Fermi fields, to be trivial in some sense. It is not clear a priori that the methods used in the purely bosonic case will lead to more than, at best, a partial solution. Yet we will adopt a conservative strategy and try to adapt the DHR/DR theory to  $\mathbb{Z}_2$ -graded nets. The fermionic version of Theorem 2.8 will vindicate this approach.

Clearly, the criterion (1.2) makes sense also for  $\mathbb{Z}_2$ -graded nets. Since things are complicated by the spacelike anticommutativity of fermionic operators, the assumption of twisted duality for  $\mathcal{F}$  is, however, not sufficient to deduce that representations satisfying (1.2) are equivalent to (equivalence classes) of transportable endomorphisms of  $\mathcal{F}$ . To make this clear, assume  $\pi$  satisfies (1.2), and let  $X^\mathcal{O} : \mathcal{H}_0 \rightarrow \mathcal{H}_\pi$  be such that  $X^\mathcal{O} A = \pi(A) X^\mathcal{O} \forall A \in \mathcal{F}(\mathcal{O}')$ . We would like to show that  $\rho(A) \equiv X^{\mathcal{O}*} \pi(A) X^\mathcal{O}$  maps  $\mathcal{F}(\mathcal{O}_1)$  into itself if  $\mathcal{O}_1 \supset \mathcal{O}$ . Now, let  $x \in \mathcal{F}(\mathcal{O}_1)$ ,  $y \in \mathcal{F}(\mathcal{O}_1)'$ , which implies  $xy = yx$ . We would like to apply  $\rho$  on both sides and use  $\rho(y) = y$  to conclude that

$$\rho(\mathcal{F}(\mathcal{O}_1)) \subset \mathcal{F}(\mathcal{O}_1)'' = \mathcal{F}(\mathcal{O}_1).$$

As it stands, this argument does not work, since  $\pi$  and thus  $\rho$  are defined only on the quasilocal algebra  $\mathcal{F}$ , but not on the operators  $VF_- \in \mathcal{F}'$  which result from the twisting operation. Assume, for a moment, that the representation  $\rho$  lifts to an endomorphism  $\hat{\rho}$  of the  $C^*$ -algebra  $\hat{\mathcal{F}}$  on  $\mathcal{H}$  generated by  $\mathcal{F}$  and the unitary  $V$ , such that  $\hat{\rho}(V) = V$  or, alternatively,

$\hat{\rho}(V) = -V$ . Using triviality of  $\rho$  in restriction to  $\mathcal{F}(\mathcal{O}'_1)$  we then obtain  $\rho(\mathcal{F}(\mathcal{O}'_1)') = \mathcal{F}(\mathcal{O}'_1)'$ , which justifies the above argument. Now, in order for  $\hat{\rho}(V) = \pm V$  to be consistent, we must have

$$\rho \circ \alpha_k(A) = \rho(VAV) = \hat{\rho}(V)\rho(A)\hat{\rho}(V) = V\rho(A)V = \alpha_k \circ \rho(A), \tag{2.8}$$

i.e.,  $\rho \circ \alpha_k = \alpha_k \circ \rho$ . In view of  $\rho(A) = X^{\mathcal{O}*}\pi(A)X^{\mathcal{O}}$  we can now claim:

LEMMA 2.9. – *There is a one-to-one correspondence between equivalence classes of:*

- (a) *Representations of  $\mathcal{F}$  which are, for every  $\mathcal{O} \in \mathcal{K}$ , unitarily equivalent to a representation  $\rho$  on  $\mathcal{H}_0$  such that  $\rho \upharpoonright \mathcal{F}(\mathcal{O}') = id$  and  $\rho \circ \alpha_k = \alpha_k \circ \rho$  (where  $Aut \mathcal{B}(\mathcal{H}_0) \ni \alpha_k \equiv Ad V$ );*
- (b) *Transportable localized endomorphisms of  $\mathcal{F}$  commuting with  $\alpha_k$ .*

Remark. – In (a) covariance of  $\pi$  with respect to  $\alpha_k$  is not enough. We need the fact that, upon transferring the representation to the vacuum Hilbert space via  $\rho(A) = X^{\mathcal{O}*}\pi(A)X^{\mathcal{O}}$ ,  $\alpha_k$  is implemented by the grading operator  $V$ .

Proof. – The direction (b)  $\Rightarrow$  (a) is trivial. As to the converse, by the above all that remains to prove is extendibility of  $\rho$  to  $\hat{\rho}$ . By the arguments in [46, p. 121] the  $C^*$ -crossed product (covariance algebra)  $\mathcal{F} \rtimes_{\alpha_k} \mathbb{Z}_2$  is simple such that the actions of  $\mathcal{F}$  and  $\mathbb{Z}_2$  on  $\mathcal{H}_0$  and  $\mathcal{H}_\pi$  via  $\pi_0 = id$ ,  $V$  and  $\pi$ ,  $V_\pi$  can be considered as faithful representations of the crossed product. Thus there is an isomorphism between  $C^*(\mathcal{F}, V)$  and  $C^*(\pi(\mathcal{F}), V_\pi)$  which maps  $F \in \mathcal{F}$  into  $\pi(F)$  and  $V$  into  $V_\pi$ .  $\square$

DEFINITION 2.10. – *DHR representations and transportable endomorphisms are called even iff they satisfy (a) and (b) of Lemma 2.9, respectively.*

We have thus singled out a class of representations which gives rise to localized endomorphisms of the field algebra  $\mathcal{F}$ . But this class is still too large in the sense that unitarily equivalent even representations need not be inner equivalent. Let  $\rho$  be an even endomorphism of  $\mathcal{F}$ , localized in  $\mathcal{O}$ . Then  $\sigma = Ad_{UV} \circ \rho$  with  $U \in \mathcal{F}_-(\mathcal{O})$  is even and equivalent to  $\rho$  as a representation, but  $(\rho, \sigma) \cap \mathcal{F} = \{0\}$ , which precludes an extension of the DHR analysis of permutation statistics etc. Furthermore,  $\rho$  and  $\sigma$ , although they are equivalent as representations of  $\mathcal{F}$ , restrict to inequivalent endomorphisms of  $\mathcal{F}_+$ . This observation leads us to confine our attention to the following class of representations.

DEFINITION 2.11. – *An even DHR representation of  $\mathcal{F}$  is called bosonic if it restricts to a bosonic DHR representation (in the conventional sense) of the even subnet  $\mathcal{F}_+$ .*

A better understanding of this class of representations is gained by the following lemma.

LEMMA 2.12. – *There is a one-to-one correspondence between the equivalence classes of bosonic even DHR representations of  $\mathcal{F}$  and bosonic DHR representations of  $\mathcal{F}_+$ ; that is, equivalent bosonic even DHR representations of  $\mathcal{F}$  restrict to equivalent bosonic DHR representations of  $\mathcal{F}_+$ . Conversely, every bosonic DHR representation of  $\mathcal{F}_+$  extends uniquely to a bosonic even DHR representation of  $\mathcal{F}$ .*

Remark. – It will become clear in Theorem 2.14 that nothing is lost by considering only representations which restrict to bosonic sectors of  $\mathcal{F}_+$ .

Proof. – Clearly, the restriction of a bosonic even DHR representation of  $\mathcal{F}$  to  $\mathcal{F}_+$  is a bosonic DHR representation. Let  $\rho, \sigma$  be irreducible even DHR morphisms of  $\mathcal{F}$ , localized in  $\mathcal{O}$ , and let  $T \in (\rho, \sigma)$ . Twisted duality implies  $T \in \mathcal{F}(\mathcal{O})'$ , i.e.,  $T = T_+ + T_- V$  where  $T_{\pm} \in \mathcal{F}_{\pm}$ . Now both sides of

$$\sigma(F) = T_+ \rho(F) T_+^* + T_- \alpha_k \circ \rho(F) T_-^* + T_+ \rho(F) V T_-^* + T_- V \rho(F) T_+^* \tag{2.9}$$

must commute with  $\alpha_k$ . The first two terms on the right hand side obviously having this property, we obtain

$$T_+ \rho(F) V T_-^* + T_- V \rho(F) T_+^* = 0 \quad \forall F \in \mathcal{F}.$$

For  $F = F^*$  this reduces to  $T_+ \rho(F) T_-^* = 0$ , which can be true only if  $T_+ = 0$  or  $T_- = 0$  since  $\rho$  is irreducible. The case  $T = T_- V$  is ruled out by the requirement that the restrictions of  $\rho$  and  $\sigma$  to  $\mathcal{F}_+$  are both bosonic. Thus we conclude that  $T \in \mathcal{F}_+(\mathcal{O})$  and the restrictions  $\rho_+$  and  $\sigma_+$  are equivalent.

As to the converse, a bosonic DHR representation  $\pi_+$  of  $\mathcal{F}_+$  gives rise to a local 1-cocycle [42,43] in  $\mathcal{F}_+$ , i.e., a mapping  $z: \Sigma_1 \rightarrow \mathcal{U}(\mathcal{F}_+)$  satisfying the cocycle identity

$$z(\partial_0 c) z(\partial_2 c) = z(\partial_1 c), \quad c \in \Sigma_2$$

and the locality condition  $z(b) \in \mathcal{F}_+(|b|)$ ,  $b \in \Sigma_1$ . This cocycle can be used as in [43,44] to extend  $\pi_+$  to a representation  $\pi$  of  $\mathcal{F}$  which has



all the desired properties. We omit the details. By this construction, the extensions of equivalent representations are equivalent, an intertwiner  $T \in (\rho, \sigma)$  lifting to  $\pi(T)$  on  $\mathcal{H}$ .  $\square$

**THEOREM 2.13.** – *Let  $\mathcal{F}$  be a complete normal field net associated to a rational net of observables. Then  $\mathcal{F}$  does not possess non-trivial bosonic even DHR representations with finite statistics. Equivalently, there are no non-trivial bosonic DHR representations of the even subalgebra  $\mathcal{F}_+$  with finite statistics.*

*Proof.* – Assume that  $\mathcal{F}$  has non-trivial bosonic even DHR representations; by the lemma this is equivalent to the existence of bosonic sectors of  $\mathcal{F}_+$ . For the latter the conventional DHR analysis goes through and gives rise to a semigroup  $\Sigma$  of endomorphisms of  $\mathcal{F}_+$  with permutation symmetry etc. These morphisms lift to  $\mathcal{F}$  and we can apply the DR construction to  $(\mathcal{F}, \Sigma)$ . Since all elements of  $\Sigma$  are bosonic, no bosonization in the sense of [22, (3.19)] is necessary. All this works irrespective of the fact that  $\mathcal{F}$  is not a local net since the fermionic fields are mere spectators. That the resulting field net again satisfies normal commutation relations is more or less evident since the ‘new’ fields are purely bosonic. Furthermore, Lemma 2.3 is still true when the ‘observable net’ is  $\mathbb{Z}_2$ -graded. Now the rest of the argument works just as in Theorem 2.4.  $\square$

*Remarks.* – 1. In the fermionic case, the even subnet  $\mathcal{F}_+$  has exactly one fermionic sector. This sector is simple and its square is equivalent to the identity, as follows from the fact that bosonic sectors of  $\mathcal{F}_+$  do not exist.

2. At this point one might be suspicious that there exist relevant DHR-like representations of  $\mathcal{F}$  which are not covered by this theorem. In particular the restriction to bosonic even DHR representations was made for reasons which may appear to be purely technical and physically weakly motivated. The next theorem shows that this is not the case.

**THEOREM 2.14.** – *Let  $\mathcal{A}$  be a rational net of observables, let  $\Gamma \subset \Delta$  be a subsemigroup of DHR morphisms containing not only bosonic sectors and let  $\mathcal{F}_\Gamma$  be the incomplete  $\mathbb{Z}_2$ -graded field net associated with  $(\mathcal{A}, \Gamma)$ . Then an application of the DR construction with respect to the bosonic even morphisms  $\Sigma$  of  $\mathcal{F}_\Gamma$ , as described above, leads to a field net  $\mathcal{F}_{\Gamma, \Sigma}$  which is equivalent to the complete normal field net  $\mathcal{F}_\Delta$ .*

*Proof.* – Since  $\mathcal{F}$  is assumed to contain fermions, every  $\mathcal{F}(\mathcal{O})$  contains unitaries which are odd under  $\alpha_k$ , giving rise to fermionic automorphisms of  $\mathcal{A}$ . By composition with one of these, every irreducible endomorphism

of  $\mathcal{A}$  can be made bosonic. It is thus clear that it suffices to extend  $\mathcal{F}$  by Bose fields which implement these bosonic sectors (more precisely, their extensions to  $\mathcal{F}$ ). The rest of the argument goes as in the preceding subsection.  $\square$

It is thus the existence of bosonic sectors of the even subnet which indicates that a fermionic field net is not complete, and only such sectors need to be considered when enlarging the field net in order to obtain the complete field net.

### 3. DEGENERATE SECTORS IN 1+1 DIMENSIONS

#### 3.1. General results on degenerate sectors

We begin with a few easy but crucial results on the set of degenerate DHR sectors. Let  $\mathcal{O} \mapsto \mathcal{A}(\mathcal{O})$  be a net of observables satisfying Haag duality on the line or in 1+1 dimensional Minkowski space. (For remarks on the duality assumption cf. the end of the next subsection.) As shown in [23], with each pair of localized endomorphisms there are associated two a priori different statistics operators  $\varepsilon(\rho, \eta), \varepsilon(\eta, \rho)^* \in (\rho\eta, \eta\rho)$ .

**DEFINITION 3.1** ([39]). – *Two DHR sectors have trivial monodromy iff the corresponding morphisms satisfy  $\varepsilon(\rho, \eta) = \varepsilon(\eta, \rho)^*$  or, equivalently,  $\varepsilon_M(\rho, \eta) = \mathbf{1}$  (this is independent of the choice of  $\rho, \eta$  within their equivalence classes). A DHR sector is degenerate iff it has trivial monodromy with all sectors (it suffices to consider the irreducible ones).*

A convenient criterion for trivality of the monodromy of two morphisms is given by the following

**LEMMA 3.2.** – *Let  $\rho, \eta$  be irreducible localized endomorphisms. Furthermore, let  $\eta_L, \eta_R$  be equivalent to  $\eta$  localized to the spacelike left and right of  $\rho$ , respectively, with the (unique up to a constant) intertwiner  $T \in (\eta_L, \eta_R)$ . Then trivality of the monodromy  $\varepsilon_M(\rho, \eta)$  is equivalent to  $\rho(T) = T$ .*

*Proof.* – Using the intertwiners  $T_{L/R} \in (\eta, \eta_{L/R})$ , the statistics operators are given by  $\varepsilon(\rho, \eta) = T_L^* \rho(T_L)$  and  $\varepsilon(\eta, \rho) = T_R^* \rho(T_R)$ . The monodromy operator is given by

$$\varepsilon_M(\rho, \eta) = T_L^* \rho(T_L T_R^*) T_R.$$

Thus,  $\varepsilon_M(\rho, \eta) = \mathbf{1}$  is equivalent to  $\rho(T_L T_R^*) = T_L T_R^*$ . The proof is completed by the observation that  $T_L T_R^*$  equals  $T^*$  up to a phase.  $\square$

At first sight one might be tempted to erroneously conclude from this lemma that there is no non-trivial braid statistics as follows: The above charge transporting intertwiner  $T$  commutes with  $\mathcal{A}(\mathcal{O})$  which, appealing to Haag duality, implies that it is contained in the algebra of the spacelike complement  $\mathcal{O}'$ . On this algebra every morphism localized in  $\mathcal{O}$  acts trivially, so that the lemma implies permutation group statistics. The mistake in this argument is, of course, that  $T$  is contained in the weakly closed algebra  $\mathcal{R}(\mathcal{O}') \equiv \mathcal{A}(\mathcal{O}')'' = \mathcal{A}(\mathcal{O})'$  but not necessarily in the  $C^*$ -subalgebra  $\mathcal{A}(\mathcal{O}')$  of the quasilocal algebra  $\mathcal{A}$ . It is only the latter on which  $\rho$  is known to act trivially.

LEMMA 3.3. – *A reducible DHR representation  $\pi$  is degenerate iff all irreducible subrepresentations are degenerate.*

*Proof.* – Let  $\rho$  be equivalent to  $\pi$  and localized in  $\mathcal{O}$ , decomposing into irreducibles according to  $\rho = \sum_{i \in I} V_i \rho_i(\cdot) V_i^*$ . That is, the  $\rho_i$  are localized in  $\mathcal{O}$  and  $V_i \in \mathcal{A}(\mathcal{O})$  with  $V_i^* V_j = \delta_{i,j} \mathbf{1}$  and  $\sum_i V_i V_i^* = \mathbf{1}$ . By Lemma 3.2,  $\pi$  is degenerate iff  $\rho(T) = T$  for every unitary intertwiner between (irreducible) morphisms  $\sigma, \sigma'$  which are localized in the two different connected components of  $\mathcal{O}'$ . Now,  $\rho(T) = \sum_i V_i \rho_i(T) V_i^*$  equals  $T$  iff ‘all matrix elements are equal’, i.e.,  $V_j^* T V_k = \delta_{j,k} \rho_j(T) \forall j, k \in I$ . But since  $T \in \mathcal{A}(\mathcal{O})'$  the left hand side equals  $T V_j^* V_k = T \delta_{j,k}$  which leads to the necessary and sufficient condition  $\rho_j(T) = T \forall j \in I$ , which in turn is equivalent to all  $\rho_i$  being degenerate.  $\square$

LEMMA 3.4. – *Let  $\Delta_D$  be the set of all degenerate morphisms with finite statistical dimension. Then  $(\Delta_D, \varepsilon)$  is a permutation symmetric, specially directed semigroup with subobjects and direct sums.*

*Proof.* – Let  $\rho_1, \rho_2$  be degenerate, i.e.,  $\varepsilon_M(\rho_i, \sigma) = \mathbf{1} \forall \sigma$ . Due to the identities [24]

$$\varepsilon(\rho_1 \rho_2, \sigma) = \varepsilon(\rho_1, \sigma) \rho_1(\varepsilon(\rho_2, \sigma)), \tag{3.1}$$

$$\varepsilon(\sigma, \rho_1 \rho_2) = \rho_1(\varepsilon(\sigma, \rho_2)) \varepsilon(\sigma, \rho_1) \tag{3.2}$$

we have

$$\begin{aligned} \varepsilon_M(\rho_1 \rho_2, \sigma) &= \varepsilon(\rho_1 \rho_2, \sigma) \varepsilon(\sigma, \rho_1 \rho_2) \\ &= \varepsilon(\rho_1, \sigma) \rho_1(\varepsilon(\rho_2, \sigma) \varepsilon(\sigma, \rho_2)) \varepsilon(\sigma, \rho_1) = \mathbf{1}. \end{aligned} \tag{3.3}$$

Thus  $\Delta_D$  is closed under composition. By the preceding lemma the direct sum of degenerate morphisms is degenerate, and every irreducible morphism contained in a degenerate one is again degenerate. That

$(\Delta_D, \varepsilon)$  is specially directed in the sense of [20, Section 5] follows as in [22, Lemma 3.7] from the fact that the degenerate sectors have permutation group statistics.  $\square$

### 3.2. Proof of a conjecture by Rehren

In Section 2 we proved that the DR field net corresponding to a rational theory  $\mathcal{A}$  in  $\geq 2 + 1$  dimensions does not have DHR sectors (with finite statistics). The dimensionality of spacetime entered in the arguments only insofar as it implies permutation group statistics. By the results of the preceding subsection it is now clear that we can proceed as in Section 2, restricting ourselves to the degenerate sectors. More concretely, we apply the spatial version [20, Corollary 6.2] of the construction of the crossed product to the quasilocal observable algebra and the semigroup  $\Delta_D$  of degenerate sectors with finite statistics. As the proofs in [22, Section 3] were given for  $\geq 2 + 1$  spacetime dimensions it seems advisable to reconsider them in order to be on the safe side, in particular as far as (twisted) duality for the field net is concerned.

**PROPOSITION 3.5.** – *Let  $\mathcal{F}$  be the spatial crossed product [20, Corollary 6.2] of  $\mathcal{A}$  by  $(\Delta, \varepsilon)$  where  $\Delta$  is as in Lemma 3.4, and let  $\mathcal{F}(\mathcal{O})$  be defined as in the proof of [22, Theorem 3.5]. Then  $\mathcal{O} \mapsto \mathcal{F}(\mathcal{O})$  is a normal field system with gauge symmetry and satisfies twisted duality.*

*Proof.* – The proof of existence in [22, Theorem 3.5] holds unchanged as it relies only on algebraic arguments independent of the spacetime dimension. The same holds for [22, Theorem 3.6] with the possible exception of the argument leading to twisted duality on p. 73. The latter boils down algebraically to the identity

$$\mathcal{F}(\mathcal{O})' \cap G' = \pi(\mathcal{A}(\mathcal{O}'))^-, \quad \mathcal{O} \in \mathcal{K}.$$

Finally, the proof of this formula in [22, Lemma 3.8] is easily verified to be correct in  $1 + 1$  dimensions, too, provided  $\mathcal{A} \upharpoonright \mathcal{H}_0$  satisfies duality. In our case this is true by assumption.  $\square$

*Remarks.* – 1. The reader who feels uneasy with these few remarks is encouraged to study the proofs of [22, Theorems 3.5, 3.6] himself, for it would make no sense to reproduce them here.

2. It may be confusing that in theories with group symmetry satisfying the split property for wedges (SPW), Haag duality for a field net  $\mathcal{F}$  and the  $G$ -fixpoint net  $\mathcal{A}$  (in the vacuum sector) are in fact incompatible [36].

The SPW has been verified for massive free scalar and Dirac fields and is probably true in all reasonable *massive* theories. On the other hand, a net of observables which satisfies Haag duality and the SPW does not admit DHR sectors anyway [37]. In view of this result, we implicitly assume in this section that the observables do *not* satisfy the SPW. The point is that one must be careful to distinguish between conformally covariant or at least massless theories, with which we are concerned here, and massive theories since the scenarios are quite different.

**THEOREM 3.6.** – *Let  $\Delta_D$  be the set of all degenerate morphisms with finite statistics, corresponding to only finitely many sectors. If  $\Delta_D$  is purely bosonic, the local field net  $\mathcal{F}$  constructed from  $\mathcal{A}$  and  $\Delta_D$  does not have degenerate sectors with finite statistics. If  $\Delta_D$  contains fermionic sectors, the normal field net  $\mathcal{F}$  does not have degenerate bosonic even sectors with finite statistics. Equivalently, the even subnet has no degenerate bosonic sectors with finite statistics.*

*Proof.* – The proofs of Theorems 2.4, 2.13 are valid also in the 2-dimensional situation since neither the argument of Lemma 2.3 on the extendibility of local symmetries nor the uniqueness result of [22, Theorem 3.5] require any modification.  $\square$

This result, which was conjectured by Rehren in [39], is quite interesting and potentially useful for the analysis of superselection structure in  $1 + 1$  dimensions. It seems worthwhile to restate it in the following form.

**COROLLARY 3.7.** – *Every degenerate quantum field theory in  $1 + 1$  dimensions (in the sense that there are degenerate sectors, which in the rational case is equivalent to non-invertibility of  $S$ ) arises as the fixpoint theory of a non-degenerate theory under the action of a compact group of inner symmetries. That is, all degenerate theories are orbifold theories in the sense of [9].*

Now we indicate how the preceding arguments have to be changed in the case of conformal theories. We first remark that everything we have said about theories in  $1 + 1$  dimension remains true for theories on the line. In the rest of this subsection we consider chiral theories on the circle [24], which are defined by associating a von Neumann algebra  $\mathcal{A}(I)$  to each interval  $I$  which is not dense in  $S^1$ . The ‘spacetime’ symmetry is given by the Möbius group  $PSU(1, 1) \cong PSL(2, \mathbb{R})$ . The set of intervals not being directed, the quasilocal algebra must be replaced by

the universal algebra of [24] which has a non-trivial center due to the non-simply connectedness of  $S^1$ . Triviality of the center of the  $C^*$ -algebra  $\mathcal{A}$  was, however, an essential requirement for the Doplicher–Roberts analysis, in particular [17]. One may try to eliminate this condition, but we prefer another approach. We begin by restricting the theory to the punctured circle, i.e., the line, for which one has the conventional quasilocal algebra  $\mathcal{A}$  which is simple. The following result shows that the restriction of generality which this step seems to imply—since Haag duality on  $\mathbb{R}$  is equivalent to strong additivity, which does not hold for all theories—is only apparent.

**PROPOSITION 3.8.** – *Given a chiral conformal precosheaf the localized endomorphisms of the algebra  $\mathcal{A}$  (corresponding to the punctured circle) and their statistics can be defined without assuming duality on the real line. In the case without fermionic degenerate sectors the DR construction can be applied to the (Möbius covariant) degenerate sectors with finite statistics and yields a conformal precosheaf  $\mathcal{F}$  on  $S^1$  which is Möbius covariant with positive energy and does not have degenerate sectors (with finite statistics).*

*Proof.* – Let  $\mathcal{A}$  be the quasilocal algebra obtained after removing a point at infinity. By the results of [24, Section 5] the semigroup  $\Delta \subset \text{End } \mathcal{A}$  of localized endomorphisms with braiding can be defined without assuming duality on the line. We can thus apply the DR construction to  $\mathcal{A}$  and  $\Delta_D$  and obtain a field net  $\mathbb{R} \supset I \mapsto \mathcal{F}(I)$ , but we clearly cannot hope to prove Haag duality on  $\mathbb{R}$ . Before we can prove duality on  $S^1$  (as it holds for the observables) we must define the local algebras for intervals which contain the point at infinity. Due the conformal spin-statistics theorem [27] a covariant bosonic degenerate sector, having statistics phase 1, is in fact covariant under the uncovered Möbius group  $PSL(2, \mathbb{R})$ , and consequentially also the extended theory  $\mathcal{F}$  is Möbius covariant with positive energy. This fact can be used to define the missing local algebras and to obtain a conformal precosheaf on  $S^1$ . Then the abstract results of [4,26] apply and yield Haag duality on  $S^1$ . The argument to the effect that the extended theory is non-degenerate works as above.  $\square$

### 3.3. Relating the superselection structures of $\mathcal{A}$ and $\mathcal{F}$

In the preceding subsection we have seen that whenever there are degenerate sectors one can construct an extended theory which is non-degenerate. The larger theory has a group symmetry such that the

original theory is reobtained by retaining only the invariant operators. Equivalently, all degenerate theories are orbifold theories. By this result, a general analysis of the superselection structure in  $1 + 1$  dimensions may begin by considering the non-degenerate case. It remains, however, to clarify the relation between the superselection structures of the degenerate theory and the extended theory. This will not be attempted here, but we will provide some results going in this direction.

**LEMMA 3.9.** – *All irreducible morphisms contained in the product of a degenerate morphism and a non-degenerate morphism are non-degenerate.*

*Proof.* – The fact that the composition of degenerate morphisms yields a sum of degenerate morphisms can be expressed in terms of the fusion coefficients  $N_{ij}^k$  as

$$i \text{ and } j \text{ degenerate, } k \text{ non-degenerate} \Rightarrow N_{ij}^k = 0. \quad (3.4)$$

By Frobenius reciprocity  $N_{ij}^k = N_{ik}^{\bar{j}}$  this implies

$$i \text{ and } j \text{ degenerate, } k \text{ non-degenerate} \Rightarrow N_{ik}^{\bar{j}} = 0. \quad (3.5)$$

(We have used the fact that the conjugate  $\bar{\rho}$  is degenerate iff  $\rho$  is degenerate.)  $\square$

*Remark.* – This simple fact may be interpreted by saying that the set of non-degenerate sectors is acted upon by the set of degenerate ones, i.e., a group dual  $\widehat{G}$ . If  $G$  is abelian,  $K = \widehat{G}$  is itself an abelian group and we are in the situation studied, e.g., in [25]. The result of the preceding subsection thus constitutes a mathematically rigorous though rather abstract solution of the field identification problem [45], cf. also the remarks in the concluding section.

Being able to apply the DR construction also in  $1 + 1$  dimensions provided we consider only semigroups of degenerate endomorphisms, we are led to reconsider Lemma 2.6 concerning the extension of localized endomorphisms of the observable algebra to the field net. The construction given in Section 2 cannot be used for the extension of non-degenerate morphisms  $\eta$  since we do not have a complete field net at our disposal. There are at least two approaches to the problem which do not rely on the existence of a complete field net. The first one [33, Proposition 3.9] uses the inclusion theory of von Neumann factors which,

however, we want to avoid in this work since a proof of factoriality of the local algebras exists only for conformally covariant QFTs but not for general theories. Another prescription was given by Rehren [40]. The claim of uniqueness made there has, however, to be made more precise. Furthermore, it is not completely trivial to establish the existence part. Fortunately, both of these questions can be clarified in a relatively straightforward manner by generalizing results by Doplicher and Roberts. In [20, Section 8] they considered a similar extension problem, namely the extension of automorphisms of  $\mathcal{A}$  to automorphisms of  $\mathcal{F}$  commuting with the gauge group. The application that these authors had in mind was the extension of spacetime symmetries to the field net [22, Section 6] under the provision that the endomorphisms implemented by the fields are covariant. For a morphism  $\rho \in \Delta$  the inner endomorphism of  $\mathcal{F}$  which extends  $\rho$  will also be denoted by  $\rho$ .

LEMMA 3.10. – *Let  $\mathcal{B}$  be the crossed product [20] of the  $C^*$ -algebra  $\mathcal{A}$  with center  $\mathbb{C}\mathbf{1}$  by the permutation symmetric, specially directed semigroup  $(\Delta, \varepsilon)$  of endomorphisms and let  $G$  be the corresponding gauge group. Let  $\Gamma$  be a semigroup of unital endomorphisms of  $\mathcal{A}$ . Then there is a one-to-one correspondence between actions of  $\Gamma$  on  $\mathcal{B}$  by unital endomorphisms  $\tilde{\eta}$  which extend  $\eta \in \Gamma$  and commute elementwise with  $G$  and mappings  $(\rho, \eta) \mapsto W_\rho(\eta)$  from  $\Delta \times \Gamma$  to unitaries of  $\mathcal{A}$  satisfying*

$$W_\rho(\eta) \in (\rho\eta, \eta\rho), \tag{3.6}$$

$$W_{\rho'}(\eta)TW_\rho(\eta)^* = \eta(T), \quad T \in (\rho, \rho'), \tag{3.7}$$

$$W_{\rho\rho'}(\eta) = W_\rho(\eta)\rho(W_{\rho'}(\eta)), \tag{3.8}$$

$$W_\rho(\eta\eta') = \eta(W_\rho(\eta'))W_\rho(\eta) \tag{3.9}$$

for all  $\eta, \eta' \in \Gamma, \rho, \rho' \in \Delta$ . The correspondence is determined by

$$\tilde{\eta}(\psi) = W_\rho(\eta)\psi, \quad \psi \in H_\rho, \rho \in \Delta, \eta \in \Gamma. \tag{3.10}$$

Furthermore, if a unitary  $S \in (\eta, \eta'), \eta, \eta' \in \Gamma$  satisfies

$$SW_\rho(\eta) = W_\rho(\eta')\rho(S) \quad \forall \rho \in \Delta, \tag{3.11}$$

then  $S \in (\tilde{\eta}, \tilde{\eta}')$ .

*Proof.* – An inspection of the proofs of [20, Theorem 8.2, Corollary 8.3], where groups of automorphisms were considered, makes plain that they are valid also for the case of semigroups of true endomorphisms and we refrain from repeating them. Besides  $\eta$  not necessarily



being onto, the only change occurred in (3.6) which replaces the property  $W_\rho \in (\rho, \beta\rho\beta^{-1})$  which does not make sense for a proper endomorphism  $\beta$ . Given  $\tilde{\eta}$  and setting

$$W_\rho = \sum_{i=1}^d \tilde{\eta}(\psi_i)\psi_i^*, \tag{3.12}$$

where  $\psi_i, i = 1, \dots, d$ , is a basis of  $H_\rho$ , it is clear that  $W_\rho$  satisfies (3.6). The other properties of the  $W$ 's are proved as in [20]. As to the converse direction, we are done provided we can show that [20, Theorem 8.1] concerning the extension of  $\eta$  to the cross product of  $\mathcal{A}$  by a single endomorphism  $\rho$  generalizes to the case of  $\eta$  an endomorphism. We give only that part of the argument which differs from the one in [20].

Therefore let  $\mathcal{A}$  and  $\rho$  satisfy the assumptions of [20, Theorem 8.1], let  $\eta$  be an injective unital endomorphism of  $\mathcal{A}$  and let  $W \in (\rho\eta, \eta\rho)$  satisfy [20, (8.1), (8.2)]. As in [20, Theorem 8.1] we consider the monomorphisms of  $\mathcal{A}$  and  $\mathcal{O}_d$  into  $\mathcal{A} \otimes_\mu \mathcal{O}_d$ , defined by  $\pi': A \mapsto \eta(A) \otimes_\mu \mathbf{1}$  and  $\zeta': \psi \mapsto W \otimes_\mu \psi, \psi \in H$ , respectively. The calculation leading to  $\zeta'(\psi)\pi'(A) = \pi' \circ \rho(A)\zeta'(\psi)$  is correct also for  $\eta$  a true endomorphism. Furthermore, with  $\zeta'_1 = \zeta' \upharpoonright \mathcal{O}_{SU(d)}$  we have  $\zeta'_1(\mathcal{O}_{SU(d)}) \in \eta(\mathcal{A})$  thanks to the conditions on  $W$  and the fact that  $\mathcal{O}_{SU(d)}$  is generated by the elements  $S$  and  $\theta$ , see [18]. Thus  $\eta\rho\eta^{-1} \circ \zeta'_1$  is well defined and equals  $\zeta'_1 \circ \sigma$ , where  $\sigma$  is the canonical endomorphism of  $\mathcal{O}_{SU(d)}$ . As in [20] we conclude that  $\zeta' \upharpoonright \mathcal{O}_{SU(d)} = \eta \circ \mu$ . Thus by the universality of  $\mathcal{A} \otimes_\mu \mathcal{O}_d$  there is an isomorphism between  $\mathcal{A} \otimes_\mu \mathcal{O}_d$  and the subalgebra generated by  $\pi'(\mathcal{A})$  and  $\zeta'(\psi)$ . Equivalently, there is an endomorphism  $\gamma$  of  $\mathcal{A} \otimes_\mu \mathcal{O}_d$  such that

$$\gamma(A \otimes_\mu \mathbf{1}) = \eta(A) \otimes_\mu \mathbf{1}, \quad A \in \mathcal{A}, \tag{3.13}$$

$$\gamma(\mathbf{1} \otimes_\mu \psi) = W \otimes_\mu \psi, \quad \psi \in H. \tag{3.14}$$

Now the rest of the proof goes exactly as in [20, Theorem 8.1], i.e., after factoring out the ideal  $J_\phi$  we obtain an endomorphism  $\tilde{\eta}$  of the crossed product  $\mathcal{B} = (\mathcal{A} \otimes_\mu \mathcal{O}_d) / J_\phi$  which commutes with the action of the gauge group  $G$ .

Now let  $S \in (\eta, \eta')$  satisfy (3.11). Then for  $\psi \in H_\rho$  we have

$$S\tilde{\eta}(\psi) = SW_\rho(\eta)\psi = W_\rho(\eta')\rho(S)\psi = W_\rho(\eta')\psi S = \tilde{\eta}'(\psi)S. \tag{3.15}$$

Since  $\tilde{\eta}, \tilde{\eta}'$  are determined by their action on the spaces  $H_\rho$  this implies  $S \in (\tilde{\eta}, \tilde{\eta}')$ .  $\square$

We are now ready to consider the wanted extensions of localized endomorphisms. Motivated by Lemma 2.6 where we had (in the case of bosonic  $\psi^\rho$ )

$$\widehat{\eta}(\psi^\rho) = \sum_i \psi_i^\eta \psi^\rho \psi_i^{\eta*} = \left( \sum_{i,j} \psi_i^\eta \psi_j^\rho \psi_i^{\eta*} \psi_j^{\rho*} \right) \psi^\rho = \varepsilon(\rho, \eta) \psi^\rho, \tag{3.16}$$

we appeal to the preceding lemma with  $W_\rho(\eta) = \varepsilon(\rho, \eta)$ .

**PROPOSITION 3.11.** – *Let  $\mathcal{O} \mapsto \mathcal{F}(\mathcal{O})$  be the field net obtained via the Doplicher–Roberts construction from the algebra  $\mathcal{A}$  of observables and a semigroup of degenerate morphisms, closed under direct sums, subobjects and conjugates. Then every localized (unital) endomorphism  $\eta$  of  $\mathcal{A}$  extends to a localized endomorphism  $\widetilde{\eta}$  of  $\mathcal{F}$  commuting with the action of the gauge group. If  $\eta$  is localized in  $\mathcal{O}$  the same holds for  $\widetilde{\eta}$ . Every  $S \in (\eta, \sigma)$  lifts to  $S \in (\widetilde{\eta}, \widetilde{\sigma})$ .*

*Proof.* – We set  $W_\rho(\eta) = \varepsilon(\rho, \eta)$  and verify the requirements (3.6)–(3.9). Obviously (3.6) is fulfilled by definition of the statistics operator. (3.8) and (3.9) follow from (3.1) and (3.2), respectively. Finally, (3.7) is just  $\varepsilon(\rho', \eta)T = \eta(T)\varepsilon(\rho, \eta)$  which holds for  $T \in (\rho, \rho')$ . The statement on the localizations follows from the fact that  $\varepsilon(\rho, \eta) = \mathbf{1}$  if  $\rho, \eta$  are spacelike localized, since  $\rho$  is degenerate. Finally, with  $S \in (\eta, \sigma)$  one has  $\varepsilon(\rho, \eta)S = \rho(S)\varepsilon(\rho, \sigma)$  such that the condition (3.11) is satisfied. Thus  $S \in (\widetilde{\eta}, \widetilde{\sigma})$ .  $\square$

*Remarks.* – 1. The above result is unaffected if the field net is fermionic. In this case the identity  $\widetilde{\eta} \circ \alpha_k = \alpha_k \circ \widetilde{\eta}$  where  $k \in G$  is the grading element (which distinguishes bosonic and fermionic fields) shows that  $\widetilde{\eta}$  leaves the statistics of fields invariant. In fact, this observation provided the motivation for introducing the notion of even DHR sectors in Section 2.3.

2. As already remarked, an alternative proof can be given using [33, Proposition 3.9]. In this approach the localization of the extended morphism in a double cone follows since the morphism  $\rho$  appearing there is the restriction to  $\mathcal{A}(\mathcal{O})$  of the canonical endomorphism  $\gamma$  from  $\mathcal{F}(\mathcal{O})$  into  $\mathcal{A}(\mathcal{O})$  for some  $\mathcal{O} \in \mathcal{K}$ . But in the situation at hand we have

$$\rho \cong \bigoplus_{\xi \in \Delta_{\text{deg}}} d\rho_\xi,$$

thus  $\rho$  is degenerate.

3. In principle, the construction of the field algebra works for every family of sectors with permutation group statistics which is closed under direct sums and subobjects. As emphasized by Rehren [40], the extension  $\tilde{\eta}$  is localized in a double cone only if the charged fields in  $\mathcal{F}$  correspond to *degenerate* sectors, for otherwise  $\varepsilon(\rho, \eta) = \mathbf{1}$  holds only if  $\rho$  is localized to the right of  $\eta$  (or left, if  $\varepsilon(\eta, \rho)^*$  is used).

In the special case where  $\eta$  is an automorphism, the extension  $\tilde{\eta}$  can be defined via [20, Theorem 8.2], using  $W_\rho(\eta)$  as above. Clearly,  $\tilde{\eta}$  is irreducible since it is an automorphism. In general, however, the extension  $\tilde{\eta}$  will not be irreducible. Rehren’s description [40] of the relative commutant can also be proved rigorously by adapting earlier results [19, Lemma 5.1].

LEMMA 3.12. – *The relative commutant  $\mathcal{F} \cap \tilde{\eta}(\mathcal{F})'$  is generated as a closed linear space by sets of the form  $(\rho\eta, \eta)H_\rho$ ,  $\rho \in \Delta$ .*

*Proof.* – By twisted duality, an operator in  $\tilde{\eta}(\mathcal{F})'$  is contained in  $\mathcal{F}(\mathcal{O})'$ , where  $\mathcal{O}$  is the localization region of  $\eta$ . Due to  $\mathcal{F} \cap \mathcal{F}' = \mathcal{F}_+$ , the selfintertwiners of  $\tilde{\eta}$  in  $\mathcal{F}$  are bosonic. Obviously,  $(\rho\eta, \eta)\psi$ ,  $\psi \in H_\rho$  is in  $\mathcal{F} \cap \eta(\mathcal{A})'$ . Now, just as  $\eta$ , so can  $\rho\eta$  be extended to an endomorphism  $\tilde{\rho\eta}$  of  $\mathcal{F}$  by the proposition. Furthermore,  $T \in (\rho\eta, \eta)$  lifts to an intertwiner between  $\tilde{\rho\eta}$  and  $\tilde{\eta}$ . Thus  $(\rho\eta, \eta)\psi^\rho$  is in  $\mathcal{F} \cap \tilde{\eta}(\mathcal{F})'$ . As to the converse,  $\tilde{\eta}(\mathcal{F})' \cap \mathcal{F}$  is globally stable under the action of  $G$  since  $\tilde{\eta}$  commutes with  $G$ . Thus  $\tilde{\eta}(\mathcal{F})' \cap \mathcal{F}$  is generated linearly by its irreducible tensors under  $G$ . If  $T_1, \dots, T_d$  is such a tensor from  $\mathcal{F} \cap \tilde{\eta}(\mathcal{F})'$ , then there is a multiplet  $\psi_i$ ,  $i = 1, \dots, d$ , of isometries in  $\mathcal{F}$  and transforming in the same way, since the field algebra has full Hilbert  $G$ -spectrum. With  $X = \sum_{i=1}^d T_i \psi_i^* \in \mathcal{F}^G$  we have  $T_i = X\psi_i$  and we must prove  $X \in (\rho\eta, \eta)$ . Now  $T_i F = F T_i$  for  $F \in \tilde{\eta}(\mathcal{F})$  implies

$$X\psi_i \tilde{\eta}(F) = X\rho(\tilde{\eta}(F))\psi_i = \tilde{\eta}(F)X\psi_i, \quad F \in \mathcal{F}, \quad i = 1, \dots, d. \quad (3.17)$$

Multiplying the second identity with  $\psi_i^*$  and summing over  $i$  we obtain  $X\rho(\tilde{\eta}(F)) = \tilde{\eta}(F)X$ ,  $F \in \mathcal{F}$  since  $\sum_i \psi_i \psi_i^* = \mathbf{1}$  by construction of the field algebra. Thus  $X \in (\rho\eta, \eta)$ .  $\square$

COROLLARY 3.13. –  *$\tilde{\eta}$  is irreducible iff the endomorphism  $\eta\tilde{\eta}$  of  $\mathcal{A}$  does not contain a nontrivial morphism  $\rho \in \Delta$ .*

*Proof.* – By the lemma, the existence of a morphism  $\rho \in \Delta$  with  $(\eta\rho, \eta) \neq \{0\}$  is necessary and sufficient for the non-triviality of  $\mathcal{F} \cap \tilde{\eta}(\mathcal{F})'$ . But by Frobenius reciprocity,  $\eta\rho \succ \eta$  is equivalent to  $\eta\tilde{\eta} \succ \tilde{\rho}$ .  $\square$

*Remark.* – The irreducible endomorphisms obtained by decomposing an extension  $\tilde{\eta}$  are even, provided we use bosonic isometric intertwiners. This can always be done as the relative commutant is contained in  $\mathcal{F}_+$  by Lemma 3.12.

In the above considerations we had to assume, for the technical reasons explained in Section 2, that there are only finitely many degenerate sectors. There was, however, no restriction on the number of non-degenerate sectors. We conclude with an important observation concerning rational theories.

**PROPOSITION 3.14.** – *Let  $\mathcal{A}$  have finitely many DHR sectors (degenerate and non-degenerate) with finite statistics. Then the extended theory  $\mathcal{F}$  has only finitely many sectors with finite statistics (all of which are non-degenerate by Theorem 3.6).*

*Proof.* – As a consequence of Proposition 3.11 we have  $\widetilde{\eta_1 \oplus \eta_2} \cong \tilde{\eta}_1 \oplus \tilde{\eta}_2$ , since the intertwiners  $V_i \in (\eta_i, \eta_1 \oplus \eta_2)$ ,  $i = 1, 2$ , are in  $(\tilde{\eta}_i, \tilde{\eta}_1 \oplus \tilde{\eta}_2)$ . Thus every irreducible  $\mathcal{F}$ -sector  $\beta$  which is contained in the extension  $\tilde{\eta}$  of an  $\mathcal{A}$ -sector  $\eta$  is already contained in the extension of an irreducible  $\eta'$ . As observed in [40] the statistical dimensions of  $\eta, \tilde{\eta}$  satisfy  $d_\eta = d_{\tilde{\eta}}$ . (This follows from the existence of a left inverse  $\tilde{\phi}_{\tilde{\eta}}$  which extends  $\phi_\eta$  and the fact that  $\varepsilon(\eta, \eta) = \varepsilon(\tilde{\eta}, \tilde{\eta})$  again by Proposition 3.11.) Thus  $\tilde{\eta}$  decomposes into finitely many irreducible  $\mathcal{F}$ -sectors. Since by assumption the number of irreducible  $\mathcal{A}$ -sectors is finite we obtain only finitely many irreducible  $\mathcal{F}$ -sectors by inducing from  $\mathcal{A}$ . The claim follows provided we can show that every irreducible  $\beta \in \Delta_{\mathcal{F}}$  is contained in some  $\tilde{\eta}$ . This is true by the observation after [3, Theorem 3.21], but we prefer to give a direct argument.

For  $\beta \in \Delta_{\mathcal{F}}(\mathcal{O})$  also  $\beta_g = \alpha_g \circ \beta \circ \alpha_g^{-1}$ ,  $g \in G$ , is localized in  $\mathcal{O}$  and transportability is easy. Let now  $\{V_g, g \in G\}$  be a multiplet of isometries in  $\mathcal{F}(\mathcal{O})$  satisfying  $V_g^* V_h = \delta_{g,h} \mathbf{1}$ ,  $\sum_g V_g V_g^* = \mathbf{1}$  and  $\alpha_h(V_g) = V_{hg}$ . Such a multiplet exists since the action of  $G$  on  $\mathcal{F}(\mathcal{O})$  by construction has full Hilbert spectrum. Then

$$\bar{\beta}(\cdot) := \sum_{g \in G} V_g \beta_g(\cdot) V_g^* \tag{3.18}$$

commutes with the action of  $G$  and thus restricts to a transportable endomorphism of  $\mathcal{A}$  which is localized in  $\mathcal{O}$ . Now  $\bar{\beta}$  can be considered as an extension to  $\mathcal{F}$ , commuting with  $G$ , of  $\bar{\beta} \upharpoonright \mathcal{A}$  and by Lemma 3.10 there exists a family  $\{W_\rho(\bar{\beta} \upharpoonright \mathcal{A}), \rho \in \Delta\}$  satisfying (3.6)–(3.8) and

the boundary condition  $W_\rho(\bar{\beta} \upharpoonright \mathcal{A}) = \mathbf{1}$  for  $\rho \in \Delta_{\mathcal{A}}(\mathcal{O}')$ . Since  $\rho$  is degenerate the unique solution to these equations is

$$W_\rho(\bar{\beta} \upharpoonright \mathcal{A}) = \varepsilon(\rho, \bar{\beta} \upharpoonright \mathcal{A}).$$

This implies  $\widetilde{\bar{\beta} \upharpoonright \mathcal{A}} = \bar{\beta}$  and in view of  $\beta = \beta_e \prec \bar{\beta}$ ,  $\beta$  is contained in the extension to  $\mathcal{F}$  of the  $\mathcal{A}$ -sector  $\bar{\beta} \upharpoonright \mathcal{A}$ .  $\square$

*Remark.* – The connection with the argument in [3, Theorem 3.21] is provided by the observation that the expression  $\gamma = \sum_g V_g \alpha_g(\cdot) V_g^*$  built with the above  $V$ 's is nothing but the extension [33, Corollary 3.3] to  $\mathcal{F}$  of a canonical endomorphism from  $\mathcal{F}(\mathcal{O})$  into  $\mathcal{A}(\mathcal{O})$ . Furthermore,  $\bar{\beta} \upharpoonright \mathcal{A}$  coincides with the canonical restriction  $\sigma_\beta := \gamma \circ \beta \upharpoonright \mathcal{A}$  of [33,3]. Relying on our version of the induction procedure (Lemma 3.10), the above proof is clearly independent of the inclusion theory of von Neumann factors used in [33,3] and thus more in the spirit of DHR/DR theory to which the assumption of local factoriality is alien. (The main application of our considerations being to conformal theories where local factoriality is automatic, this added generality admittedly is not very important.)

#### 4. A SUFFICIENT CRITERION FOR NON-DEGENERACY

The results of this section depend on an additional axiom, the split property.

**DEFINITION 4.1.** – *An inclusion  $A \subset B$  of von Neumann algebras is split [16] if there exists a type I factor  $N$  such that  $A \subset N \subset B$ . A net of algebras satisfies the ‘split property for double cones’ if the inclusion  $\mathcal{A}(\mathcal{O}_1) \subset \mathcal{A}(\mathcal{O}_2)$  is split whenever  $\mathcal{O}_1 \Subset \mathcal{O}_2$ , i.e., the closure of  $\mathcal{O}_1$  is contained in the interior of  $\mathcal{O}_2$ .*

The importance of this property derives from the fact [1,16] that it is equivalent to the following formulation: For each pair of double cones  $\mathcal{O}_1 \Subset \mathcal{O}_2$  the algebra  $\mathcal{A}(\mathcal{O}_1) \vee \mathcal{A}(\mathcal{O}_2)'$  is unitarily equivalent to the tensor product  $\mathcal{A}(\mathcal{O}_1) \otimes \mathcal{A}(\mathcal{O}_2)'$ . It is believed that this form of the split property is satisfied in all reasonable quantum field theories.

In the rest of this section we will give a sufficient criterion for the absence of degenerate sectors. The following result is independent of the number of spacetime dimensions.

**PROPOSITION 4.2.** – *Let  $\mathcal{O} \mapsto \mathcal{A}(\mathcal{O})$  be a net of observables fulfilling Haag duality and the split property for double cones. Let  $\rho$  be an*

endomorphism of the quasilocal algebra  $\mathcal{A}$  which is localized in  $\mathcal{O}$  and acts identically on the relative commutant  $\mathcal{A}(\widehat{\mathcal{O}}) \cap \mathcal{A}(\mathcal{O})'$  whenever  $\widehat{\mathcal{O}} \ni \mathcal{O}$ . Then  $\rho$  is an inner endomorphism of  $\mathcal{A}$ , i.e., a direct sum of copies of the identity morphism.

*Proof.* – Choose double cones  $\mathcal{O}_1, \mathcal{O}_2$  fulfilling  $\mathcal{O} \Subset \mathcal{O}_1 \Subset \mathcal{O}_2 \Subset \widehat{\mathcal{O}}$ . Thanks to the split property there exist type I factors  $M_1, M_2$  such that

$$\mathcal{A}(\mathcal{O}) \subset \mathcal{A}(\mathcal{O}_1) \subset M_1 \subset \mathcal{A}(\mathcal{O}_2) \subset M_2 \subset \mathcal{A}(\widehat{\mathcal{O}}). \quad (4.1)$$

We first show  $\rho(M_1) \subset M_1$ . If  $A \in M_1$  we have  $\rho(A) \in \mathcal{A}(\mathcal{O}_2)$ . Due to  $\mathcal{A}(\mathcal{O}) \subset M_1$  and the premises,  $\rho$  acts trivially on  $M_1' \cap \mathcal{A}(\widehat{\mathcal{O}})$ . Thus

$$\rho(A) \in (M_1' \cap \mathcal{A}(\widehat{\mathcal{O}}))' \cap M_2 \subset (M_1' \cap M_2)' \cap M_2 = M_1. \quad (4.2)$$

The last identity follows from  $M_1, M_2$  being type I factors. Thus  $\rho$  restricts to an endomorphism of  $M_1$ . Now every endomorphism of a type I factor is inner [31, Corollary 3.8], i.e., there is a (possibly infinite) family of isometries  $V_i \in M_1$ ,  $i \in I$ , with  $V_i^* V_j = \delta_{i,j}$ ,  $\sum_{i \in I} V_i V_i^* = \mathbf{1}$  such that

$$\rho \upharpoonright M_1 = \eta(A) \equiv \sum_{i \in I} V_i \cdot V_i^*. \quad (4.3)$$

(The sums over  $I$  are understood in the strong sense.) Now by (4.3) and the premises,  $\eta \upharpoonright \mathcal{A}(\mathcal{O})' \cap M_1 = id$ , implying

$$V_i \in (\mathcal{A}(\mathcal{O})' \cap M_1)' \cap M_1 = \mathcal{A}(\mathcal{O}) \quad \forall i \in I.$$

Therefore  $\rho = \eta$  on  $\mathcal{A}(\mathcal{O}_1)$  and  $\rho = \eta = id$  on  $\mathcal{A}(\mathcal{O}')$ . In order to prove  $\rho = \eta$  on all of  $\mathcal{A}$  it suffices to show  $\rho(A) = \eta(A) \quad \forall A \in \mathcal{A}(\mathcal{O}_2)$  where we may, of course, assume  $\mathcal{O}_2 \supset \mathcal{O}_1$ . For the moment we take for granted that

$$\mathcal{A}(\mathcal{O}_2) = \mathcal{A}(\mathcal{O}_1) \vee (\mathcal{A}(\mathcal{O}_2) \wedge \mathcal{A}(\mathcal{O})'). \quad (4.4)$$

Having just proved  $\rho \upharpoonright \mathcal{A}(\mathcal{O}_1) = \eta$  and remarking that  $\rho \upharpoonright \mathcal{A}(\mathcal{O}_2) \wedge \mathcal{A}(\mathcal{O})' = id = \eta$  is true by assumption, we conclude by local normality that  $\rho \upharpoonright \mathcal{A}(\mathcal{O}_2) = \eta$ . In order to prove (4.4), apply the split property to the inclusion  $\mathcal{O} \Subset \mathcal{O}_1$ . Under the split isomorphism we have

$$\begin{aligned} \mathcal{A}(\mathcal{O}) &\cong \mathcal{A}(\mathcal{O}) \otimes \mathbf{1}, \\ \mathcal{A}(\mathcal{O}_i) &\cong \mathcal{B}(\mathcal{H}_0) \otimes \mathcal{A}(\mathcal{O}_i), \quad i = 1, 2. \end{aligned} \quad (4.5)$$

Thus  $\mathcal{A}(\mathcal{O})' \cong \mathcal{A}(\mathcal{O})' \otimes B(\mathcal{H}_0)$  and  $\mathcal{A}(\mathcal{O}_2) \wedge \mathcal{A}(\mathcal{O})' \cong \mathcal{A}(\mathcal{O})' \otimes \mathcal{A}(\mathcal{O}_2)$ , from which (4.4) follows at once.  $\square$

*Remark.* – The first part of the proof is essentially identical to [15, Proposition 2.3]. There it was stated only for automorphisms but the possibility of the above extension was remarked. In [15] the  $C^*$ -version of the time-slice axiom was used to conclude  $\rho = \eta$  on  $\mathcal{A}$ . We have dispensed with this assumption by requiring trivality of  $\rho$  on the relative commutant  $\mathcal{A}(\hat{\mathcal{O}}) \cap \mathcal{A}(\mathcal{O})'$  for all  $\hat{\mathcal{O}} \ni \mathcal{O}$ . For our purposes this will be sufficient.

We are now in a position to state our criterion for the absence of degenerate sectors in  $1 + 1$  dimensions:

**COROLLARY 4.3.** – *Assume in addition to the conditions of the proposition that for each pair  $\mathcal{O} \in \hat{\mathcal{O}}$  the algebra  $\mathcal{A}(\hat{\mathcal{O}}) \cap \mathcal{A}(\mathcal{O})'$  is generated by the charge transporters from  $\mathcal{O}_L$  to  $\mathcal{O}_R$  (and vice versa). Here  $\mathcal{O}_L, \mathcal{O}_R$  are the connected components of  $\hat{\mathcal{O}} \cap \mathcal{O}'$ , see the figure below. Then there are no degenerate sectors. More precisely, every degenerate endomorphism is inner in the above sense.*

*Proof.* – Due to Lemma 3.2, a degenerate morphism localized in  $\mathcal{O}$  acts trivially on the charge transporters between  $\mathcal{O}_L$  to  $\mathcal{O}_R$ . As these are weakly dense in  $\mathcal{A}(\hat{\mathcal{O}}) \cap \mathcal{A}(\mathcal{O})'$  by assumption and due to local normality, the morphism acts trivially on the relative commutant. This is true for every  $\hat{\mathcal{O}} \ni \mathcal{O}$ . The statement now follows from Proposition 4.2.  $\square$

*Remarks.* – 1. In [37] we show that a much further-reaching result can be proved if one requires the split property not only for double cones but also for wedge regions. This property can, however, hold only in massive quantum field theories.

2. Admittedly the condition on the relative commutant seems difficult to verify. One may perhaps hope that something can be said in the case of rational theories, which have finitely many sectors.

3. It is likely that the condition on the relative commutant made in the corollary is also necessary. The argument goes as follows. If there are degenerate sectors then there is a field net  $\mathcal{F}$  with group symmetry such that the net  $\mathcal{A}$  is the restriction of the invariant subnet to the vacuum sector, cf. the next section. Assuming that the field net also satisfies the split property, one can define localized implementers of the gauge group as in [6]. In particular, for every inclusion  $\Lambda = (\mathcal{O} \in \hat{\mathcal{O}})$  and every  $x \in U(G)'' \cap U(G)'$  one obtains an operator  $U_\Lambda(x) \in \mathcal{A}(\hat{\mathcal{O}}) \cap \mathcal{A}(\mathcal{O})'$ . We




Fig. 1. Relative commutant of double cones.

see no reason why  $U_A(x)$  should be contained in the algebra generated by  $\mathcal{A}(O_L)$ ,  $\mathcal{A}(O_R)$  and the charge transporters.

## 5. SUMMARY AND OUTLOOK

In this work we have proved two intuitively reasonable properties of the Doplicher–Roberts construction: The (essentially unique) complete field net which describes all DHR sectors has itself no localized sectors, and also it can be obtained from an intermediate, thus incomplete, field net by an application of the DR construction. Unfortunately, we have been able give a proof only under the quite restrictive assumption that there are only finitely many sectors, which is equivalent to finiteness of the gauge group  $G$ . We emphasize that the problem consists in proving compactness of  $\overline{G}$  in Proposition 2.3. If this proposition can be generalized the rest of the arguments goes through unchanged. In any case, the complete (with respect to the DHR sectors) field net may still have non-trivial representations with the weaker Buchholz–Fredenhagen localization property.

In a sense, the situation in low dimensions is quite similar. The degenerate sectors may be considered ‘better localized’ than generic DHR sectors insofar as they arise from *local* fields, in contrast to what is to be expected in the general case. Non-local charged fields played a role, e.g., in [36] where, however, the underlying quantum symmetry was spontaneously broken. As we intend to show elsewhere, the symmetry breakdown encountered there is generic in massive models. As was mentioned above, the peculiar nature of the superselection structure of massive models manifests itself also in an analysis which starts from the observables [37]. For this reason, the considerations in Sections 3



and 4 were aimed primarily at conformally covariant theories in  $1 + 1$  dimensions.

Turning to a brief discussion of open problems, the most obvious one is relaxing the rationality assumption on the superselection structure of the observable net in the proof of Proposition 2.3, which is the basis of most of our results. In trying so it is not inconceivable that one may find counterexamples, but the author is convinced that this cannot happen for theories with countably many sectors.

The hope expressed in Remark 2 after Corollary 4.3 has already been vindicated by reducing the relative commutant property needed there to a simple numerical identity which can be proved for large classes of models, cf. [38].

In Section 3.2 and Proposition 3.14 we have proven that every rational QFT in  $1 + 1$  dimensions can be extended to a rational non-degenerate one (on a bigger Hilbert space). To the new theory  $\mathcal{F}$  Rehren's analysis [39] applies and proves that the category of DHR sectors is a modular category in the sense of Turaev. Since the identification of the degenerate sectors with a group dual [17–20] has a categorical analogue [21] it is very natural to conjecture that there exists an abstract version of this construction for braided  $C^*$ -tensor categories. We conclude this paper with the following

*CONJECTURE 5.1. – Given a braided  $C^*$ -tensor category with dual objects, direct sums and subobjects (thus automatically a ribbon category by [34, Section 4]) one can construct a non-degenerate category of the same type using methods from [21]. If the original category is rational, the same is true for the new one which thus is modular.*

Work on this conjecture is in progress.

## ACKNOWLEDGEMENT

I am greatly indebted to K.-H. Rehren for proposing the subject of this work and for many useful discussions. This paper would not have been possible without his earlier work in [39,40]. I thank R. Longo and J. Roberts for drawing my attention to a serious gap in the first version of this paper. I am grateful to them as well as to R. Conti, D. Guido and R. Verch for useful discussions. Again, many thanks are due to P. Croome for a very thorough proofreading of an earlier version.

## REFERENCES

- [1] C. D'ANTONI and R. LONGO, Interpolation by type I factors and the flip automorphism, *J. Funct. Anal.* 51 (1983) 361–371.
- [2] C. D'ANTONI, Technical properties of the quasilocal algebra, in: D. Kastler, ed., *The Algebraic Theory of Superselection Sectors. Introduction and Recent Results*, World Scientific, 1990.
- [3] J. BÖCKENHAUER and D.E. EVANS, Modular invariants, graphs and  $\alpha$ -induction for nets of subfactors I, *Commun. Math. Phys.* 197 (1998) 361–386.
- [4] R. BRUNETTI, D. GUIDO and R. LONGO, Modular structure and duality in QFT, *Commun. Math. Phys.* 156 (1993) 201–219.
- [5] D. BUCHHOLZ, G. MACK and I. TODOROV, The current algebra on the circle as a germ of local field theories, *Nucl. Phys. B (Proc. Suppl.)* 5B (1988) 20–56.
- [6] D. BUCHHOLZ, S. DOPLICHER and R. LONGO, On Noether's theorem in quantum field theory, *Ann. Phys.* 170 (1986) 1–17.
- [7] D. BUCHHOLZ, S. DOPLICHER, R. LONGO and J. ROBERTS, Extensions of automorphisms and gauge symmetries, *Commun. Math. Phys.* 155 (1993) 123–134.
- [8] R. CONTI, unpublished.
- [9] R. DIJKGRAAF, C. VAFA, E. VERLINDE and H. VERLINDE, The operator algebra of orbifold models, *Commun. Math. Phys.* 123 (1989) 485–527.
- [10] S. DOPLICHER, R. HAAG and J.E. ROBERTS, Fields, observables and gauge transformations I, *Commun. Math. Phys.* 13 (1969) 1–23.
- [11] S. DOPLICHER, R. HAAG and J.E. ROBERTS, Fields, observables and gauge transformations II, *Commun. Math. Phys.* 15 (1969) 173–200.
- [12] S. DOPLICHER, R. HAAG and J.E. ROBERTS, Local observables and particle statistics I, *Commun. Math. Phys.* 23 (1971) 199–230.
- [13] S. DOPLICHER and J.E. ROBERTS, Fields, statistics and non-abelian gauge groups, *Commun. Math. Phys.* 28 (1972) 331–348.
- [14] S. DOPLICHER, R. HAAG and J.E. ROBERTS, Local observables and particle statistics II, *Commun. Math. Phys.* 35 (1974) 49–85.
- [15] S. DOPLICHER, Local aspects of superselection rules, *Commun. Math. Phys.* 85 (1982) 73–86.
- [16] S. DOPLICHER and R. LONGO, Standard and split inclusions of von Neumann algebras, *Invent. Math.* 75 (1984) 493–536.
- [17] S. DOPLICHER and J.E. ROBERTS, A remark on compact automorphism groups of  $C^*$ -algebras, *J. Funct. Anal.* 66 (1986) 67–72.
- [18] S. DOPLICHER and J.E. ROBERTS, Duals of compact Lie groups realized in the Cuntz algebras and their actions on  $C^*$ -algebras, *J. Funct. Anal.* 74 (1987) 96–120.
- [19] S. DOPLICHER and J.E. ROBERTS, Compact group actions on  $C^*$ -algebras, *J. Oper. Theor.* 19 (1988) 283–305.
- [20] S. DOPLICHER and J.E. ROBERTS, Endomorphisms of  $C^*$ -algebras, cross products and duality for compact groups, *Ann. Math.* 130 (1989) 75–119.
- [21] S. DOPLICHER and J.E. ROBERTS, A new duality theory for compact groups, *Invent. Math.* 98 (1989) 157–218.

- [22] S. DOPLICHER and J.E. ROBERTS, Why there is a field algebra with a compact gauge group describing the superselection structure in particle physics, *Commun. Math. Phys.* 131 (1990) 51–107.
- [23] K. FREDENHAGEN, K.-H. REHREN and B. SCHROER, Superselection sectors with braid group statistics and exchange algebras I, General theory, *Commun. Math. Phys.* 125 (1989) 201–226.
- [24] K. FREDENHAGEN, K.-H. REHREN and B. SCHROER, Superselection sectors with braid group statistics and exchange algebras II. Geometric aspects and conformal covariance, *Rev. Math. Phys.* (Special Issue) (1992) 113–157.
- [25] J. FUCHS, A.N. SCHELLEKENS and C. SCHWEIGERT, A matrix  $S$  for all simple current extensions, *Nucl. Phys. B* 473 (1996) 323–366.
- [26] F. GABBIANI and J. FRÖHLICH, Operator algebras and conformal field theory, *Commun. Math. Phys.* 155 (1993) 569–640.
- [27] D. GUIDO and R. LONGO, The conformal spin-statistics theorem, *Commun. Math. Phys.* 181 (1996) 11–35.
- [28] R. HAAG, *Local Quantum Physics*, 2nd Ed., Springer, 1996.
- [29] E. HEWITT and K.A. ROSS, *Abstract Harmonic Analysis I*, 2nd Ed., Springer, 1979.
- [30] D. KASTLER (ed.), *The Algebraic Theory of Superselection Sectors. Introduction and Recent Results*, World Scientific, 1990.
- [31] R. LONGO, Simple injective subfactors, *Adv. Math.* 63 (1987) 152–171.
- [32] R. LONGO, Index of subfactors and statistics of quantum fields I, *Commun. Math. Phys.* 126 (1989) 217–247.
- [33] R. LONGO and K.-H. REHREN, Nets of subfactors, *Rev. Math. Phys.* 7 (1995) 567–597.
- [34] R. LONGO and J.E. ROBERTS, A theory of dimension, *K-Theory* 11 (1997) 103–159.
- [35] M. LÜSCHER and G. MACK, Global conformal invariance in QFT, *Commun. Math. Phys.* 41 (1975) 203–234.
- [36] M. MÜGER, Quantum double actions on operator algebras and orbifold quantum field theories, *Commun. Math. Phys.* 191 (1998) 137–181.
- [37] M. MÜGER, Superselection structure of massive quantum field theories in  $1 + 1$  dimensions, *Rev. Math. Phys.* 10 (1998) 1147–1170.
- [38] M. MÜGER, On the algebraic structure of rational conformal theories, in preparation.
- [39] K.-H. REHREN, Braid group statistics and their superselection rules, in: D. Kastler, ed., *The Algebraic Theory of Superselection Sectors. Introduction and Recent Results*, World Scientific, 1990.
- [40] K.-H. REHREN, Markov traces as characters for local algebras, *Nucl. Phys. B* (Proc. Suppl.) 18B (1990) 259–268.
- [41] J.E. ROBERTS, Cross products of von Neumann algebras by group duals, *Sympos. Math.* 20 (1976) 335–363.
- [42] J.E. ROBERTS, Local cohomology and superselection structure, *Commun. Math. Phys.* 51 (1976) 107–119.
- [43] J.E. ROBERTS, Net cohomology and its applications to field theory, in: L. Streit, ed., *Quantum Fields, Particles, Processes*, Springer, 1980.
- [44] J.E. ROBERTS, Lectures on algebraic quantum field theory, in: D. Kastler, ed., *The Algebraic Theory of Superselection Sectors. Introduction and Recent Results*, World Scientific, 1990.

- [45] A.N. SCHELLEKENS and S. YANKIELOWICZ, Simple currents, modular invariants, and fixed points, *Int. J. Mod. Phys. A* 5 (1990) 2903–2952.
- [46] S.J. SUMMERS, Normal product states for Fermions and twisted duality for CCR- and CAR-type algebras with application to the Yukawa<sub>2</sub> quantum field model, *Commun. Math. Phys.* 86 (1982) 111–141.
- [47] E. VERLINDE, Fusion rules and modular transformations in 2D conformal field theory, *Nucl. Phys. B* 300 (1988) 360–376.

*Note added in proof.* The work announced as [38] has been merged into a joint paper with Y. Kawahigashi and R. Longo, Multiinterval subfactors and modularity of representations in conformal field theory (math.OA/9903104). The abstract category theoretic approach which was conjectured to exist at the end of the paper has been developed in M. Müger: Galois theory for braided tensor categories and the modular closure (math.CT/9812040), to appear in *Adv. Math.*