

The Arithmetical Complexity of Dimension and Randomness

John M. Hitchcock^{*} Jack H. Lutz[†] Sebastiaan A. Terwijn[‡]

Abstract

Constructive dimension and constructive strong dimension are effectivizations of the Hausdorff and packing dimensions, respectively. Each infinite binary sequence A is assigned a dimension $\dim(A) \in [0, 1]$ and a strong dimension $\text{Dim}(A) \in [0, 1]$.

Let DIM^α and $\text{DIM}_{\text{str}}^\alpha$ be the classes of all sequences of dimension α and of strong dimension α , respectively. We show that DIM^0 is properly Π_2^0 , and that for all Δ_2^0 -computable $\alpha \in (0, 1]$, DIM^α is properly Π_3^0 .

To classify the strong dimension classes, we use a more powerful effective Borel hierarchy where a co-enumerable predicate is used rather than an enumerable predicate in the definition of the Σ_1^0 level. For all Δ_2^0 -computable $\alpha \in [0, 1)$, we show that $\text{DIM}_{\text{str}}^\alpha$ is properly in the Π_3^0 level of this hierarchy. We show that $\text{DIM}_{\text{str}}^1$ is properly in the Π_2^0 level of this hierarchy.

We also prove that the class of Schnorr random sequences and the class of computably random sequences are properly Π_3^0 .

Keywords: arithmetical hierarchy, Wadge reductions, constructive dimension, Schnorr randomness, computable randomness

1 Introduction

Hausdorff dimension – the most extensively studied fractal dimension – has recently been effectivized at several levels of complexity, yielding applications to a variety of topics in theoretical computer science, including data compression, polynomial-time degrees, approximate optimization, feasible prediction, circuit-size complexity, Kolmogorov complexity, and randomness [14, 15, 3, 1, 7, 5, 8, 17]. The most fundamental of these effectivizations is *constructive dimension*, which is closely related to Kolmogorov complexity and algorithmic randomness. Every subset \mathcal{X} of \mathbf{C} , the *Cantor space* of all infinite binary sequences, is assigned a constructive dimension $\text{cdim}(\mathcal{X}) \in [0, 1]$. Informally, this dimension is determined by the maximum rate of growth that a lower semicomputable martingale can achieve on all sequences in \mathcal{X} .

Just as Martin-Löf [16] used constructive measure to define the randomness of individual sequences, Lutz [15] used constructive dimension to define the dimensions of individual sequences.

^{*}Department of Computer Science, University of Wyoming, Laramie, WY 82071, USA. jhitchco@cs.uwyo.edu. This author's research was supported in part by National Science Foundation grant 9988483.

[†]Department of Computer Science, Iowa State University, Ames, IA 50011, USA. lutz@cs.iastate.edu. This author's research was supported in part by National Science Foundation grants 9988483 and 0344187.

[‡]Technische Universität Wien, Wiedner Hauptstrasse 8-10, A-1040 Vienna, Austria. terwijn@logic.at. Supported by the Austrian Research Fund (Lise Meitner grant M699-N05). Part of this author's research was done while visiting the second author at Caltech in January 2002.

Each sequence $A \in \mathbf{C}$ is assigned a *dimension* $\dim(A) \in [0, 1]$ by $\dim(A) = \text{cdim}(\{A\})$. Every Martin-Löf random sequence has dimension 1, but there are nonrandom sequences with dimension 1. For every real number $\alpha \in [0, 1]$, there is a sequence with dimension α .

It is useful to understand the arithmetical complexity of a class of sequences. For example, knowing that RAND , the class of Martin-Löf random sequences, is a Σ_2^0 -class allows the application of Kreisel's Basis Lemma [12, 18] to give a short proof [26] that

$$\text{RAND} \cap \Delta_2^0 \neq \emptyset. \quad (1.1)$$

For any $\alpha \in [0, 1]$, let

$$\text{DIM}^\alpha = \{A \in \mathbf{C} \mid \dim(A) = \alpha\}.$$

Lutz [15] showed that

$$\text{DIM}^\alpha \cap \Delta_2^0 \neq \emptyset \quad (1.2)$$

for any Δ_2^0 -computable $\alpha \in [0, 1]$. As these dimension classes do not appear to be Σ_2^0 , Lutz was unable to apply the Basis Lemma to them, so he used different techniques to prove (1.2).

We investigate the complexities of these dimension classes in terms of the arithmetical hierarchy of subsets of \mathbf{C} . We show that DIM^0 is properly Π_2^0 , and for all Δ_2^0 -computable $\alpha \in (0, 1]$ we show that DIM^α is properly Π_3^0 . Therefore, the proof for (1.1) using Kreisel's Basis Lemma cannot be used directly to establish (1.2). (See however the comments made after Corollary 4.11.)

More recently, packing dimension, another important fractal dimension, has also been effectivized by Athreya, Hitchcock, Lutz, and Mayordomo [2]. At the constructive level, this is used in an analogous way to define the *strong dimension* $\text{Dim}(A) \in [0, 1]$ for every sequence A . For any $\alpha \in [0, 1]$, let

$$\text{DIM}_{\text{str}}^\alpha = \{A \in \mathbf{C} \mid \text{Dim}(A) = \alpha\}.$$

To classify these strong dimension classes, we use a more powerful effective Borel hierarchy where a co-enumerable predicate is used rather than an enumerable predicate in the definition of the Σ_1^0 level. We show that $\text{DIM}_{\text{str}}^1$ is properly in the Π_2^0 level of this stronger hierarchy. For all Δ_2^0 -computable $\alpha \in [0, 1]$, we show that $\text{DIM}_{\text{str}}^\alpha$ is properly in the Π_3^0 level of this hierarchy.

Our techniques for classifying the dimension and strong dimension classes include Baire category, Wadge reductions, and Kolmogorov complexity. In Section 4.3 we point out that ad hoc methods are sometimes necessary.

Section 2 gives an overview of the randomness and dimension notions used in this paper. In Section 3 we introduce the stronger effective Borel hierarchy that we use for the strong dimension classes. Section 4 presents the classification of DIM^α and $\text{DIM}_{\text{str}}^\alpha$.

We conclude the paper with Section 5 on effective randomness classes. We restate a result of Schnorr [20] concerning computable null sets of exponential order in terms of *computable dimension* and point out a relationship with *Church randomness*. We prove that the class of *Schnorr random* sequences and that the class of *computably random* sequences are properly Π_3^0 .

2 Background on Randomness and Dimension

This section provides an overview of the notions of randomness and dimension used in this paper. We write $\{0, 1\}^*$ for the set of all finite binary *strings* and \mathbf{C} for the *Cantor space* of all infinite binary *sequences*. In the standard way, a sequence $A \in \mathbf{C}$ can be identified with the subset of

$\{0, 1\}^*$ or \mathbb{N} for which it is the characteristic sequence, or with a real number in the unit interval. The length of a string $w \in \{0, 1\}^*$ is $|w|$. The string consisting of the first n bits of $x \in \{0, 1\}^* \cup \mathbf{C}$ is denoted by $x \upharpoonright n$. We write $w \sqsubseteq x$ if w is a prefix of x .

2.1 Martin-Löf Randomness

Martin-Löf [16] introduced the notion of a *constructive null set*. A set is constructively null if it can be covered by a uniform sequence of c.e. open sets that are shrinking in size. That is, $\mathcal{A} \subseteq \mathbf{C}$ is constructively null if $\mathcal{A} \subseteq \bigcap_i \mathcal{U}_i$, where $\{\mathcal{U}_i\}_{i \in \mathbb{N}}$ is uniformly c.e. such that $\mu(\mathcal{U}_i) \leq 2^{-i}$. The sequence $\{\mathcal{U}_i\}_{i \in \mathbb{N}}$ is called a *Martin-Löf test*. An individual sequence $A \in \mathbf{C}$ is *Martin-Löf random* if $\{A\}$ is not constructively null. The Martin-Löf random sequences play an important role in algorithmic information theory, see e.g. Li and Vitányi [13].

Schnorr [20], following Ville [27], characterized constructive null sets in terms of martingales. A function $d : \{0, 1\}^* \rightarrow [0, \infty)$ is a *martingale* if for every $w \in \{0, 1\}^*$, d satisfies the averaging condition

$$2d(w) = d(w0) + d(w1),$$

and d is a *supermartingale* if it satisfies

$$2d(w) \geq d(w0) + d(w1).$$

The *success set* of d is

$$S^\infty[d] = \left\{ A \in \mathbf{C} \mid \limsup_{n \rightarrow \infty} d(A \upharpoonright n) = \infty \right\},$$

i.e., it is the set of all sequences on which d has unbounded value. We say that d succeeds on a class $\mathcal{A} \subseteq \mathbf{C}$ if $\mathcal{A} \subseteq S^\infty[d]$.

Ville [27] proved that a set $\mathcal{A} \subseteq \mathbf{C}$ has Lebesgue measure 0 if and only if there is a martingale d that succeeds on \mathcal{A} . Schnorr [20] showed that \mathcal{A} is constructively null if and only if d can be chosen to be lower semicomputable, that is, if d can be computably approximated from below. We call such a d *constructive*.

Martin-Löf [16] proved that there is a universal constructive null set. That is, he proved that there is a Martin-Löf test $\{\mathcal{U}_i\}_i$ such that for every other test $\{\mathcal{V}_i\}$ it holds that $\bigcap_i \mathcal{V}_i \subseteq \bigcap_i \mathcal{U}_i$. By Schnorr's analysis this implies that there is also a universal constructive supermartingale \mathbf{d} . That is, for any constructive supermartingale d' there is a $c > 0$ such that $\mathbf{d}(w) \geq cd'(w)$ for all $w \in \{0, 1\}^*$. We will use this universal supermartingale in section 4. We denote the complement of $S^\infty[\mathbf{d}]$ by **RAND**, so that **RAND** consists of all the Martin-Löf random sequences.

2.2 Schnorr Randomness

Schnorr [20] criticized the notion of constructive null for an actual lack of constructiveness, and introduced the more constructive notion of a *Schnorr null set*, which is defined by requiring that the measure of the levels \mathcal{U}_i in a Martin-Löf test be computably approximable to within any given precision. It is easy to see that this is equivalent to the following: \mathcal{A} is Schnorr null if $\mathcal{A} \subseteq \bigcap_i \mathcal{U}_i$, where $\{\mathcal{U}_i\}_{i \in \mathbb{N}}$ is uniformly c.e. such that $\mu(\mathcal{U}_i) = 2^{-i}$. The sequence $\{\mathcal{U}_i\}_{i \in \mathbb{N}}$ is called a *Schnorr test*.

Following Schnorr [20], we call an unbounded nondecreasing function $h : \{0, 1\}^* \rightarrow \{0, 1\}^*$ an *order*. (N.B. An “Ordnungsfunktion” in Schnorr’s terminology is always computable, whereas we

prefer to leave the complexity of orders unspecified in general.) For any order h and martingale d , we define the *order h success set of d* as

$$S^h[d] = \left\{ A \in \mathbf{C} \mid \limsup_{n \rightarrow \infty} \frac{d(A \upharpoonright n)}{h(n)} \geq 1 \right\}.$$

Schnorr pointed out that the rate of success of a constructive martingale d can be so slow that it cannot be computably detected. Thus rather than working with constructive null sets of the form $S^\infty[d]$ with d constructive, he worked with null sets of the form $S^h[d]$, where both d and h are computable. He proved that a set \mathcal{A} is Schnorr null if and only if it is included in a null set of the form $S^h[d]$, with d and h computable.

A sequence $A \in \mathbf{C}$ is *Schnorr random* if $\{A\}$ is not Schnorr null. This is related the notion of computable randomness. A sequence A is *computably random* if for every computable martingale d , $A \notin S^\infty[d]$.

We write $\text{RAND}_{\text{Schnorr}}$ for the class of all Schnorr random sequences and $\text{RAND}_{\text{comp}}$ for the class of all computably random sequences. By definition we have that

$$\text{RAND} \subseteq \text{RAND}_{\text{comp}} \subseteq \text{RAND}_{\text{Schnorr}}.$$

The first inclusion was proved strict by Schnorr [20] and the second inclusion was proved strict by Wang [28].

2.3 Constructive Dimension

Hausdorff [6] introduced the concept of null covers that “succeed exponentially fast” to define what is now commonly called Hausdorff dimension, the most widely used dimension in fractal geometry. Basically, this notion allows one to discern structure in classes of measure zero, and to calibrate them. As for constructive measure, already Schnorr (see Theorem 5.1) drew special attention to null sets of “exponential order”, although he did not make an explicit connection to Hausdorff dimension.

Lutz [14, 15] gave a characterization of Hausdorff dimension in terms of *gales*, which are a generalization of martingales. Let $s \in [0, \infty)$. An s -*gale* is a function $d : \{0, 1\}^* \rightarrow [0, \infty)$ that satisfies the averaging condition

$$2^s d(w) = d(w0) + d(w1) \tag{2.1}$$

for every $w \in \{0, 1\}^*$. Similarly, d is an s -*supergale* if (2.1) holds with \geq instead of equality. The success set $S^\infty[d]$ is defined exactly as was done for martingales above. Lutz showed that for any class $\mathcal{A} \subseteq \mathbf{C}$, the Hausdorff dimension of \mathcal{A} is

$$\dim_{\text{H}}(\mathcal{A}) = \inf \left\{ s \mid \begin{array}{l} \text{there exists an } s\text{-gale} \\ d \text{ for which } \mathcal{A} \subseteq S^\infty[d] \end{array} \right\}. \tag{2.2}$$

Lutz [15] effectivized this characterization to define the constructive dimensions of sets and sequences. An s -(super)gale is called *constructive* if it is lower semicomputable. The *constructive dimension* of a class $\mathcal{A} \subseteq \mathbf{C}$ is

$$\text{cdim}(\mathcal{A}) = \inf \left\{ s \mid \begin{array}{l} \text{there exists a constructive } s\text{-gale} \\ d \text{ for which } \mathcal{A} \subseteq S^\infty[d] \end{array} \right\} \tag{2.3}$$

and the constructive dimension of an individual sequence $A \in \mathbf{C}$ is

$$\dim(A) = \text{cdim}(\{A\}).$$

(Supergales can be equivalently used in place of gales in both (2.2) and (2.3) [14, 9, 4].)

Constructive dimension has some remarkable properties. For example, Lutz [15] showed that for any class \mathcal{A} ,

$$\text{cdim}(\mathcal{A}) = \sup_{A \in \mathcal{A}} \dim(A). \quad (2.4)$$

Also, Mayordomo [17] established a strong connection with *Kolmogorov complexity*: for any $A \in \mathbf{C}$,

$$\dim(A) = \liminf_{n \rightarrow \infty} \frac{K(A \upharpoonright n)}{n}, \quad (2.5)$$

where $K(A \upharpoonright n)$ is the size of the smallest program that causes a fixed universal self-delimiting Turing machine to output the first n bits of A . (For comments on the relation of this result to earlier results, see the report [24] by Staiger and section 6 of [15]. For more details on Kolmogorov complexity, we refer to [13].)

One can also characterize constructive dimension using the Schnorr null sets (see Section 2.2) of exponential order. The following proposition was observed by several authors, including those of [1, 25].

Proposition 2.1. *Let \mathbf{d} be the universal constructive supermartingale. For any $\mathcal{A} \subseteq \mathbf{C}$,*

$$\text{cdim}(\mathcal{A}) = \inf \{s \in \mathbb{Q} : \mathcal{A} \subseteq S_{2^{(1-s)n}}[\mathbf{d}]\}.$$

2.4 Constructive Strong Dimension

More recently, Athreya, Hitchcock, Lutz, and Mayordomo [2] also characterized *packing dimension*, another important fractal dimension, in terms of gales. For this, the notion of *strong success* of an s -gale d was introduced. The *strong success set* of d is

$$S_{\text{str}}^\infty[d] = \left\{ A \in \mathbf{C} \mid \liminf_{n \rightarrow \infty} d(A \upharpoonright n) = \infty \right\}.$$

Analogously to what was done for Hausdorff dimension, packing dimension can be characterized using strong success sets of gales. Effectivizing this in the same way leads to the definition of the *constructive strong dimension* of a class $\mathcal{A} \subseteq \mathbf{C}$ as

$$\text{cDim}(\mathcal{A}) = \inf \left\{ s \mid \begin{array}{l} \text{there exists a constructive } s\text{-gale } d \\ \text{for which } \mathcal{A} \subseteq S_{\text{str}}^\infty[d] \end{array} \right\}.$$

The constructive strong dimension of a sequence $A \in \mathbf{C}$ is

$$\text{Dim}(A) = \text{cDim}(\{A\}).$$

A pointwise stability property analogous to (2.4) also holds for strong dimension, as well as a Kolmogorov complexity characterization [2]:

$$\text{Dim}(A) = \limsup_{n \rightarrow \infty} \frac{K(A \upharpoonright n)}{n} \quad (2.6)$$

for any $A \in \mathbf{C}$.

3 Borel Hierarchies

Σ_n^0 and Π_n^0 denote the levels of the Borel hierarchy for subsets of Cantor space. The levels of the arithmetical hierarchy (the corresponding effective hierarchy for sets of reals) are denoted by Σ_n^0 and Π_n^0 .

We will also make use of the following more general hierarchy definition.

Definition. Let \mathcal{P} be a class of predicates, let $n \geq 1$, and let $\mathcal{X} \subseteq \mathbf{C}$.

- $\mathcal{X} \in \Sigma_n^0[\mathcal{P}]$ if for some predicate $P \in \mathcal{P}$,

$$A \in \mathcal{X} \iff (\exists k_n)(\forall k_{n-1}) \cdots (Qk_1)P(k_n, \dots, k_2, A \upharpoonright k_1),$$

where $Q = \exists$ if n is odd and $Q = \forall$ if n is even.

- $\mathcal{X} \in \Pi_n^0[\mathcal{P}]$ if for some predicate $P \in \mathcal{P}$,

$$A \in \mathcal{X} \iff (\forall k_n)(\exists k_{n-1}) \cdots (Qk_1)P(k_n, \dots, k_2, A \upharpoonright k_1),$$

where $Q = \forall$ if n is odd and $Q = \exists$ if n is even.

If we take \mathcal{P} to be Δ_1^0 (decidable), then the above definition is equivalent to the standard arithmetical hierarchy of reals, that is

$$\Sigma_n^0 = \Sigma_n^0[\Delta_1^0]$$

and

$$\Pi_n^0 = \Pi_n^0[\Delta_1^0]$$

hold for all n . Also, if ALL is the class of all predicates, then we obtain the classical Borel hierarchy:

$$\Sigma_n^0 = \Sigma_n^0[\text{ALL}]$$

and

$$\Pi_n^0 = \Pi_n^0[\text{ALL}].$$

In this paper, we will also be interested in the cases where \mathcal{P} is Σ_1^0 (computably enumerable) or Π_1^0 (co-c.e.). In some cases, the classes in the generalized hierarchy using these sets of predicates are no different than the standard arithmetical hierarchy classes. If n is odd, then $\Sigma_n^0 = \Sigma_n^0[\Sigma_1^0]$ as the existential quantifier in the Σ_1^0 predicate can be absorbed into the last quantifier in the definition of $\Sigma_n^0[\Delta_1^0] = \Sigma_n^0$. Analogously, $\Pi_n^0 = \Pi_n^0[\Pi_1^0]$ for odd n , and for even n we have $\Sigma_n^0 = \Sigma_n^0[\Pi_1^0]$ and $\Pi_n^0 = \Pi_n^0[\Sigma_1^0]$. On the other hand, using the complementary set of predicates defines an effective hierarchy that is distinct from and interleaved with the arithmetical hierarchy.

Proposition 3.1. 1. If n is odd, then

$$\Sigma_n^0 \subsetneq \Sigma_n^0[\Pi_1^0] \subsetneq \Sigma_{n+1}^0$$

and

$$\Pi_n^0 \subsetneq \Pi_n^0[\Sigma_1^0] \subsetneq \Pi_{n+1}^0.$$

2. If n is even, then

$$\Sigma_n^0 \subsetneq \Sigma_n^0[\Sigma_1^0] \subsetneq \Sigma_{n+1}^0$$

and

$$\Pi_n^0 \subsetneq \Pi_n^0[\Pi_1^0] \subsetneq \Pi_{n+1}^0.$$

Proof. We only show $\Sigma_n^0 \subsetneq \Sigma_n^0[\Pi_1^0] \subsetneq \Sigma_{n+1}^0$ for odd n ; the arguments for the other statements are analogous.

The inclusion $\Sigma_n^0 \subseteq \Sigma_n^0[\Pi_1^0]$ is obvious. To show that it is proper, let P be a predicate that is complete for the class of Π_n^0 predicates. Then there is a decidable predicate R such that

$$P(n) \iff (\forall k_n)(\exists k_{n-1}) \cdots (\forall k_1) R(n, k_n, \dots, k_1).$$

Define $\mathcal{X} \subseteq \mathbf{C}$ as

$$\mathcal{X} = \bigcup_{n \in P} 0^n 1 \mathbf{C}.$$

Then $\mathcal{X} \in \Sigma_n^0[\Pi_1^0]$ as we have

$$\begin{aligned} S \in \mathcal{X} &\iff (\exists n) P(n) \text{ and } 0^n 1 \subseteq S \\ &\iff (\exists n)(\forall k_n)(\exists k_{n-1}) \cdots (\forall k_1) R(n, k_n, \dots, k_1) \text{ and } 0^n 1 \subseteq S \\ &\iff (\exists n)(\forall k_n)(\exists k_{n-1}) \cdots (\exists k_2) T(n, k_n, \dots, k_3, S \upharpoonright k_2), \end{aligned}$$

where T is the Π_1^0 predicate defined by

$$T(n, k_n, \dots, k_3, w) \iff (\forall k_1) R(n, k_n, \dots, k_3, |w|, k_1) \text{ and } 0^n 1 \subseteq w.$$

Now suppose that $\mathcal{X} \in \Sigma_n^0$. Then for some decidable predicate U ,

$$S \in \mathcal{X} \iff (\exists k_n)(\forall k_{n-1}) \cdots (\exists k_1) U(k_n, \dots, k_2, S \upharpoonright k_1).$$

We then have

$$\begin{aligned} n \in P &\iff 0^n 1 \mathbf{C} \subseteq \mathcal{X} \\ &\iff 0^n 10^\infty \in \mathcal{X} \\ &\iff (\exists k_n)(\forall k_{n-1}) \cdots (\exists k_1) U(k_n, \dots, k_2, 0^n 10^\infty \upharpoonright k_1), \end{aligned}$$

so P is a Σ_n^0 predicate, which contradicts its Π_n^0 -completeness. Therefore $\mathcal{X} \notin \Sigma_n^0$ and we have established $\Sigma_n^0 \subsetneq \Sigma_n^0[\Pi_1^0]$.

The inclusion $\Sigma_n^0[\Pi_1^0] \subseteq \Sigma_{n+1}^0$ is immediate from the definitions using $\Sigma_{n+1}^0 = \Sigma_{n+1}^0[\Delta_1^0]$. That it is proper follows from the facts $\Sigma_{n+1}^0 - \Sigma_n^0 \neq \emptyset$ and $\Sigma_n^0[\Pi_1^0] \subseteq \Sigma_n^0$. \square

The next proposition shows that there are no unexpected inclusions:

Proposition 3.2. 1. If n is odd, then

$$\Sigma_n^0 \not\subseteq \Pi_n^0[\Sigma_1^0] \not\subseteq \Sigma_{n+1}^0$$

and

$$\Pi_n^0 \not\subseteq \Sigma_n^0[\Pi_1^0] \not\subseteq \Pi_{n+1}^0.$$

2. If n is even, then

$$\Sigma_n^0 \not\subseteq \Pi_n^0[\Pi_1^0] \not\subseteq \Sigma_{n+1}^0$$

and

$$\Pi_n^0 \not\subseteq \Sigma_n^0[\Sigma_1^0] \not\subseteq \Pi_{n+1}^0.$$

Proof. The noninclusions on the left side all follow from Borel considerations. E.g. for the noninclusion $\Pi_n^0 \not\subseteq \Sigma_n^0[\Pi_1^0]$ take any Π_n^0 -class that is not in Σ_n^0 .

The noninclusions on the right side can all be proved by direct diagonalization. As an example we prove that $\Sigma_1^0[\Pi_1^0] \not\subseteq \Pi_2^0$. The proof easily generalizes to the higher levels. The proof is a fairly straightforward diagonalization against all possible Π_2^0 -definitions, although the details are a bit cumbersome. Let R_i , $i \in \omega$, be a computable list of all partial computable predicates. We define a class $\mathcal{X} \in \Sigma_1^0[\Pi_1^0]$ such that for all i there is $X \sqsupset 0^i 1$ such that

$$X \in (\mathcal{X} \setminus \mathcal{Y}) \cup (\mathcal{Y} \setminus \mathcal{X}), \quad (3.1)$$

where $\mathcal{Y} = \{X : \forall n \exists m R_i(n, X \upharpoonright m)\}$. So the definition of \mathcal{X} in the interval above the string $0^i 1$ will make sure that \mathcal{X} is not Π_2^0 -defined by R_i .

For the definition of \mathcal{X} we will need a uniform sequence of Π_1^0 -sets of strings P_n^i . We start by defining $\text{dom}(P_n^i)$ for each i and n such that

- $\text{dom}(P_n^i)$ is a computable subset of $\{\sigma 1 : \sigma \sqsupseteq 0^i 1\}$,
- $\text{dom}(P_n^i)$ is dense above the string $0^i 1$, i.e. for all $\tau \sqsupseteq 0^i 1$ there is $\sigma \sqsupseteq \tau$ with $\sigma \in \text{dom}(P_n^i)$,
- $\text{dom}(P_n^i) \cap \text{dom}(P_m^i) = \emptyset$ if $n \neq m$.

Then we define P_n^i by

- for every $\sigma 1 \in \text{dom}(P_n^i)$,

$$\sigma 1 \notin P_n^i \iff (\exists \tau \sqsupseteq \sigma 1)(\exists m) [\tau \text{ is of the form } \sigma 1 0^k \wedge R_i(n, \tau \upharpoonright m)]. \quad (3.2)$$

It is easy to see that such a uniform sequence of P_n^i 's exists. Now \mathcal{X} is defined as

$$\mathcal{X} = \{X : (\exists i)(\exists n)(\exists \sigma \in P_n^i) [\sigma \sqsubset X]\}.$$

The idea is that to show that R_i does not give a Π_2^0 -definition of \mathcal{X} , we challenge it by choosing a string $\sigma 1 \in P_n^i$ and extend it by 0's. Now if R_i responds by providing us with an extension τ as in (3.2), we take this τ as an initial segment of our set X , which means that there is an m such that $R_i(n, X \upharpoonright m)$, so that the condition $\forall n \exists m R_i(n, X \upharpoonright m)$ is verified for n . But by definition of P_n^i , as soon as the witness m is found, the string $\sigma 1$ falls out of the set P_n^i , so we have managed to keep X outside of \mathcal{X} while at the same time obtaining a piece of evidence that $\forall n \exists m R_i(n, X \upharpoonright m)$. If on the other hand R_i does not respond this means that $\sigma 1 \in P_n^i$, so X will be in \mathcal{X} , but *no* extension Y of $\sigma 1$ will satisfy $\forall n \exists m R_i(n, Y \upharpoonright m)$. So in both cases X is a counterexample showing that R_i does not define \mathcal{X} .

We now give the formal construction. Fix i . We construct X as in (3.1) by a finite extension construction. Let $X_0 = 0^i 1$. At stage s of the construction we are given X_s , no initial segment of which is in any P_n^i , and such that $(\forall n \leq s)(\exists m \leq |X_s|) [R_i(n, X_s \upharpoonright m)]$. Choose $\sigma 1 \in \text{dom}(P_{s+1}^i)$ such that $\sigma 1 \sqsubset X_s$.

Case I. There exists a τ as in (3.2), with $n = s + 1$. Define $X_{s+1} = \tau$ and go to the next stage of the construction.

Case II. There does not exist such a τ . Then define $X = X_s \hat{\ } 0^\omega$ and end the construction.

To verify that $X = \bigcup_s X_s$ thus constructed satisfies (3.1), note that if there is a stage where Case II obtains, then the string $\sigma 1$ chosen at that stage is in P_s^i and proves that $X \in \mathcal{X}$, whereas no extension Y of $\sigma 1$ satisfies $\exists m R_i(n, Y \upharpoonright m)$. So in this case we are done. If on the other hand at every stage of the construction Case I obtains, then for every $\sigma 1$ chosen at any stage s we have $\sigma 1 \notin P_s^i$, and hence $X \notin \mathcal{X}$ since apart from the 1's in $\sigma 1$ the string X only contains 0's, and $\text{dom}(P_n^i) \subseteq \{\sigma 1 : \sigma \sqsupseteq 0^i 1\}$. But also at every stage s a new witness m is found such that $R_i(s, X \upharpoonright m)$, hence $\forall s \exists m R_i(s, X \upharpoonright m)$, and again X satisfies (3.1). \square

Intuitively, the classes $\Sigma_1^0[\Pi_1^0]$, $\Pi_1^0[\Sigma_1^0]$, $\Sigma_2^0[\Sigma_1^0]$, $\Pi_2^0[\Pi_1^0], \dots$ are slightly more powerful than their respective counterparts in the arithmetical hierarchy because they use one additional quantifier that is limited to the predicate. We now give a simple example of a class that is best classified in this hierarchy: the class of 1-generic sequences.

Proposition 3.3. *The class of all 1-generic sequences is $\Pi_2^0[\Pi_1^0]$ but not Σ_3^0 . It is also not Σ_2^0 .*

Proof. Recall that a sequence $X \in \mathbf{C}$ is 1-generic (see e.g. Jockusch [10]) if

$$(\forall e)(\exists \sigma \sqsubset X) [\{e\}^\sigma(e) \downarrow \vee (\forall \tau \sqsupset \sigma) [\{e\}^\tau(e) \uparrow]]$$

From this definition it is immediate that the class $\mathcal{G} = \{X \mid X \text{ is 1-generic}\}$ is in $\Pi_2^0[\Pi_1^0]$. To show that \mathcal{G} is not Σ_3^0 , suppose that it is. Then there is a uniform sequence of Σ_1^0 -classes $\mathcal{O}_{n,m}$ such that $\mathcal{G} = \bigcup_n \bigcap_m \mathcal{O}_{n,m}$. Without loss of generality $\mathcal{O}_{n,m} \supseteq \mathcal{O}_{n,m+1}$ for all n, m . Now \mathcal{G} is comeager, so there is n such that $\bigcap_m \mathcal{O}_{n,m}$ is not nowhere dense, hence dense in some interval C_σ . Then every $\mathcal{O}_{n,m}$, $m \in \mathbb{N}$, is dense in C_σ . Now it is easy to construct, using a computable finite extension construction, a computable sequence (starting with σ) in $\bigcap_m \mathcal{O}_{n,m}$, contradicting that 1-generic sets are noncomputable.

That the 1-generic sets are not Σ_2^0 follows quickly from Lemma 4.3 below, noting again that the 1-generic sets are a comeager class. \square

Staiger has pointed out to us that the class $\Pi_1^0[\Sigma_1^0]$ already occurred under a different guise in [23] where it was called \mathfrak{P} , and several presentations were proven to be equivalent to it. The following definitions are contained in [22]. Let W be any set of initial segments. Define

$$\begin{aligned} \lim W &= \{A \in 2^\omega : \forall \sigma \sqsubset A (\sigma \in W)\}, \\ W^\sigma &= \{A \in 2^\omega : \forall^\infty \sigma \sqsubset A (\sigma \in W)\}. \end{aligned}$$

Staiger proved that the classes in $\Pi_1^0[\Sigma_1^0]$ are those of the form $\lim W$, for $W \in \Sigma_1^0$, and the classes in $\Sigma_2^0[\Sigma_1^0]$ are those of the form W^σ , for $W \in \Sigma_1^0$.

4 Classification of DIM^α and $\text{DIM}_{\text{str}}^\alpha$

In this section we investigate the arithmetical complexity of the following dimension and strong dimension classes.

$$\begin{aligned} \text{DIM}^\alpha &= \{A \in \mathbf{C} \mid \dim(A) = \alpha\} \\ \text{DIM}^{\leq \alpha} &= \{A \in \mathbf{C} \mid \dim(A) \leq \alpha\} \\ \text{DIM}^{\geq \alpha} &= \{A \in \mathbf{C} \mid \dim(A) \geq \alpha\} \\ \text{DIM}_{\text{str}}^\alpha &= \{A \in \mathbf{C} \mid \text{Dim}(A) = \alpha\} \\ \text{DIM}_{\text{str}}^{\leq \alpha} &= \{A \in \mathbf{C} \mid \text{Dim}(A) \leq \alpha\} \\ \text{DIM}_{\text{str}}^{\geq \alpha} &= \{A \in \mathbf{C} \mid \text{Dim}(A) \geq \alpha\} \end{aligned}$$

Let $\alpha \in [0, 1]$ be Δ_2^0 -computable. For any such α , it is well known that there is a computable function $\hat{\alpha} : \mathbb{N} \rightarrow \mathbb{Q}$ such that $\lim_{n \rightarrow \infty} \hat{\alpha}(n) = \alpha$. Using (2.5), we have

$$\begin{aligned} \dim(X) \leq \alpha &\iff \liminf_{n \rightarrow \infty} \frac{K(X \upharpoonright n)}{n} \leq \alpha \\ &\iff (\forall k)(\forall N)(\exists n \geq N) K(X \upharpoonright n) < (\hat{\alpha}(n) + 1/k)n, \end{aligned}$$

so $\text{DIM}^{\leq \alpha}$ is a Π_2^0 -class. Also,

$$\begin{aligned} \dim(X) \geq \alpha &\iff \liminf_{n \rightarrow \infty} \frac{K(X \upharpoonright n)}{n} \geq \alpha \\ &\iff (\forall k)(\exists N)(\forall n \geq N) K(X \upharpoonright n) > (\hat{\alpha}(N) - 1/k)n, \end{aligned}$$

so $\text{DIM}^{\geq \alpha}$ is a Π_3^0 -class. Therefore we have the following.

Proposition 4.1. 1. The class DIM^0 is Π_2^0 .

2. For all Δ_2^0 -computable $\alpha \in (0, 1]$, DIM^α is a Π_3^0 -class.

3. For arbitrary $\alpha \in (0, 1]$, DIM^α is a $\mathbf{\Pi}_3^0$ -class.

The situation is slightly more complicated for strong dimension. By (2.6), we have

$$\begin{aligned} \text{Dim}(X) \leq \alpha &\iff \limsup_{n \rightarrow \infty} \frac{K(X \upharpoonright n)}{n} \leq \alpha \\ &\iff (\forall k)(\exists N)(\forall n \geq N) K(X \upharpoonright n) < (\hat{\alpha}(N) + 1/k)n \\ &\iff (\forall k)(\exists N)(\forall n \geq N)(\exists \langle \pi, t \rangle) |\pi| < (\hat{\alpha}(N) + 1/k)n \\ &\quad \text{and } U(\pi) = X \upharpoonright n \text{ in } \leq t \text{ computation steps,} \end{aligned}$$

where U is the fixed universal self-delimiting Turing machine used to define K . From this it is clear that $\text{DIM}_{\text{str}}^{\leq \alpha} \in \Pi_4^0$. However, the “ $(\exists \langle \pi, t \rangle)$ ” quantifier is local to the defining predicate, so we have $\text{DIM}_{\text{str}}^{\leq \alpha} \in \mathbf{\Pi}_3^0$, and in fact, it is a $\Pi_3^0[\Sigma_1^0]$ -class. Also,

$$\begin{aligned} \text{Dim}(X) \geq \alpha &\iff \limsup_{n \rightarrow \infty} \frac{K(X \upharpoonright n)}{n} \geq \alpha \\ &\iff (\forall k)(\forall N)(\exists n \geq N) K(X \upharpoonright n) > (\hat{\alpha}(n) - 1/k)n, \end{aligned}$$

so $\text{DIM}_{\text{str}}^{\geq \alpha}$ is a $\Pi_2^0[\Pi_1^0]$ -class. This establishes the following analogue of Proposition 4.1.

Proposition 4.2. 1. The class $\text{DIM}_{\text{str}}^1$ is $\Pi_2^0[\Pi_1^0]$.

2. For all Δ_2^0 -computable $\alpha \in [0, 1)$, $\text{DIM}_{\text{str}}^\alpha$ is a $\Pi_3^0[\Sigma_1^0]$ -class.

3. For arbitrary $\alpha \in [0, 1)$, $\text{DIM}_{\text{str}}^\alpha$ is a Π_3^0 -class.

In the remainder of this section we prove that the classifications in Propositions 4.1 and 4.2 cannot be improved in their respective hierarchies.

4.1 Category Methods

Recall that a class \mathcal{X} is *meager* if it is included in a countable union of nowhere dense subsets of \mathbf{C} , and *comeager* if its complement $\overline{\mathcal{X}}$ is meager. The following lemma (implicit in Rogers [19, p341]) will be useful.

Lemma 4.3. If $\mathcal{X} \in \Sigma_2^0$ and $\overline{\mathcal{X}}$ is dense then \mathcal{X} is meager.

Proof. Suppose that $\mathcal{X} = \bigcup_n \mathcal{X}_n$, \mathcal{X}_n closed. Since $\overline{\mathcal{X}}$ is dense, \mathcal{X}_n contains no basic open set, hence \mathcal{X}_n is nondense (i.e. its closure contains no basic open set), and \mathcal{X} is a countable union of nondense sets. \square

As a warm-up we give a short proof of Shoenfield's result that the class of computable sequences is not a Π_3^0 -class.

Theorem 4.4. (Shoenfield [19, p344]) The class Δ_1^0 of computable sets is a $\Sigma_2^0[\Sigma_1^0]$ -class, but it is not a Π_3^0 -class. It is also not a Π_2^0 -class.

Proof. Clearly $\Delta_1^0 \in \Sigma_2^0[\Sigma_1^0]$. Suppose for a contradiction that $\overline{\Delta_1^0}$ is Σ_3^0 . Then there is a uniform sequence of Σ_1^0 -classes $\mathcal{O}_{n,m}$ such that $\overline{\Delta_1^0} = \bigcup_n \bigcap_m \mathcal{O}_{n,m}$. Without loss of generality $\mathcal{O}_{n,m} \supseteq \mathcal{O}_{n,m+1}$ for all n, m . Now Δ_1^0 is meager because it is countable, so $\overline{\Delta_1^0}$ is comeager, so there is an n such that $\bigcap_m \mathcal{O}_{n,m}$ is not nowhere dense, hence dense in some interval C_σ . Then every $\mathcal{O}_{n,m}$, $m \in \mathbb{N}$, is dense in C_σ . Now it is easy to construct a computable sequence (starting with σ) in $\bigcap_m \mathcal{O}_{n,m}$, contradicting that $\bigcap_m \mathcal{O}_{n,m} \subseteq \overline{\Delta_1^0}$.

That Δ_1^0 is not Π_2^0 follows from Lemma 4.3, since $\overline{\Delta_1^0}$ is comeager. \square

The class RAND of Martin-Löf random sets can easily be classified with category methods.

Theorem 4.5. (folk) RAND is a Σ_2^0 -class, but it is not a Π_2^0 -class.

Proof. This is analogous to the proof in Rogers [19, p 341] that $\{X : X \text{ finite}\}$ is a Σ_2^0 -class but not a Π_2^0 -class. Both RAND and its complement are dense, so by Lemma 4.3, RAND is meager. If RAND were a Π_2^0 -class, then again using Lemma 4.3, its complement would also be meager. This contradicts the fact that \mathbf{C} is not meager. \square

As DIM^0 and $\text{DIM}_{\text{str}}^1$ are dense Π_2^0 -classes that have dense complements, an argument similar to the one used for Theorem 4.5 shows that they are not Σ_2^0 -classes.

Theorem 4.6. The classes DIM^0 and $\text{DIM}_{\text{str}}^1$ are not Σ_2^0 -classes.

We now develop category methods for the other DIM^α classes. For every rational s , define the computable order $h_s(n) = 2^{(1-s)n}$. Let \mathbf{d} be the optimal constructive supermartingale.

Lemma 4.7. *For every rational $s \in (0, 1)$, $S^{h_s}[\mathbf{d}]$ is a comeager Π_2^0 -class.*

Proof. Notice that $\overline{S^{h_s}[\mathbf{d}]} \in \Sigma_2^0$ and $S^{h_s}[\mathbf{d}]$ is dense. Now apply Lemma 4.3. \square

Lemma 4.8. *For all $\alpha \in (0, 1]$, DIM^α is meager.*

Proof. Let $s < \alpha$ be rational. Lutz [15] showed that $\mathbf{d}^{(s)}(w) = 2^{(s-1)|w|}\mathbf{d}(w)$ is an optimal constructive s -supergale. It follows that for any $A \in \mathbf{C}$, $A \in S^{h_s}[\mathbf{d}] \Rightarrow \dim(S) < \alpha$. Therefore $\text{DIM}^\alpha \subseteq \overline{S^{h_s}}$, so DIM^α is meager by Lemma 4.7. \square

Proposition 4.9. *For all $\alpha \in (0, 1]$, DIM^α is not a Π_2^0 -class.*

Proof. If $\text{DIM}^\alpha \in \Pi_2^0$, then Lemma 4.3 implies that DIM^α is comeager, contradicting Lemma 4.8. \square

To strengthen Proposition 4.9 to show that DIM^α is not Σ_3^0 , we now turn to Wadge reductions.

4.2 Wadge Reductions

Let $\mathcal{A}, \mathcal{B} \subseteq \mathbf{C}$. A *Wadge reduction* of \mathcal{A} to \mathcal{B} is a function $f : \mathbf{C} \rightarrow \mathbf{C}$ that is continuous and satisfies $\mathcal{A} = f^{-1}(\mathcal{B})$, i.e., $X \in \mathcal{A} \iff f(X) \in \mathcal{B}$. We say that \mathcal{B} is *Wadge complete* for a class Γ of subsets of \mathbf{C} if $\mathcal{B} \in \Gamma$ and every $\mathcal{A} \in \Gamma$ Wadge reduces to \mathcal{B} . As the classes of the Borel hierarchy are closed under Wadge reductions, Wadge completeness can be used to properly identify the location of a subset of \mathbf{C} in the hierarchy.

We now prove that DIM^1 is Wadge complete for Π_3^0 . We will then give Wadge reductions from it to DIM^α for the other values of α .

Theorem 4.10. *DIM^1 is Wadge complete for Π_3^0 . Therefore DIM^1 is not a Σ_3^0 -class, and in particular it is not a Σ_3^0 -class.*

Proof. One could prove this by reducing a known Π_3^0 -complete class to DIM^1 , e.g. the class of sets that have a limiting frequency of 1's that is 0 (this class was proved to be Π_3^0 -complete by Ki and Linton [11]), but it is just as easy to build a direct reduction from an arbitrary Π_3^0 -class.

Let \mathbf{d} be the universal constructive supermartingale. Note that we have (cf. Proposition 2.1)

$$S^{2^n}[\mathbf{d}] \subsetneq \dots \subsetneq S^{2^{\frac{1}{k}n}}[\mathbf{d}] \subsetneq S^{2^{\frac{1}{k+1}n}}[\mathbf{d}] \subsetneq \dots \subsetneq \text{DIM}^1.$$

Let $\bigcup_k \bigcap_s \mathcal{O}_{k,s}$ be a Σ_3^0 -class. Without loss of generality $\mathcal{O}_{k,s} \supseteq \mathcal{O}_{k,s+1}$ for all k, s . We define a continuous function $f : \mathbf{C} \rightarrow \mathbf{C}$ such that

$$\forall k \left(X \in \bigcap_s \mathcal{O}_{k,s} \iff f(X) \in S^{2^{\frac{1}{k}n}}[\mathbf{d}] \right) \tag{4.1}$$

so that we have

$$\begin{aligned} X \notin \bigcup_k \bigcap_s \mathcal{O}_{k,s} &\iff \forall k \left(f(X) \notin S^{2^{\frac{1}{k}n}}[\mathbf{d}] \right) \\ &\iff f(X) \in \text{DIM}^1. \end{aligned}$$

The image $Y = f(X)$ is defined in stages, $Y = \bigcup_s Y_s$, such that every initial segment of X defines an initial segment of Y .

At stage 0 we define Y_0 to be the empty sequence.

At stage $s > 0$ we consider $X \upharpoonright s$, and for each k we define $t_{k,s}$ to be the largest stage $t \leq s$ such that $X \upharpoonright s \in \mathcal{O}_{k,t}$. (Let $t_{k,s} = 0$ if such a t does not exist.) Define k to be *expansionary* at stage s if $t_{k,s-1} < t_{k,s}$. Now we let $k(s) = \min\{k : k \text{ is expansionary at } s\}$. There are two substages.

Substage (a). First consider all strings σ extending Y_{s-1} of minimal length with $\mathbf{d}(\sigma) \geq 2^{\frac{1}{k(s)}|\sigma|}$, and take the leftmost one of these σ 's. Such σ 's exist because $S^{2^{\frac{1}{k(s)}n}}[\mathbf{d}]$ is dense. If $k(s)$ does not exist, let $\sigma = Y_{s-1}$.

Substage (b). Next consider all extensions $\tau \sqsupseteq \sigma$ of minimal length such that $\mathbf{d}(\tau \upharpoonright i) \leq \mathbf{d}(\tau \upharpoonright (i-1))$ for every $|\sigma| < i < |\tau|$, and $\mathbf{d}(\tau) \leq |\tau|$. Clearly such τ exist, by direct diagonalization against \mathbf{d} . Define Y_s to be the leftmost of these τ . This concludes the construction.

So Y_s is defined by first building a piece of evidence σ that \mathbf{d} achieves growth rate $2^{\frac{1}{k(s)}n}$ on Y and then slowing down the growth rate of \mathbf{d} to the order n . Note that f is continuous. If $X \in \bigcup_k \bigcap_s \mathcal{O}_{k,s}$, then for the minimal k such that $X \in \bigcap_s \mathcal{O}_{k,s}$, infinitely many pieces of evidence σ witness that \mathbf{d} achieves growth rate $2^{\frac{1}{k}n}$ on Y , so $Y \notin \text{DIM}^1$. On the other hand, if $X \notin \bigcup_k \bigcap_s \mathcal{O}_{k,s}$ then for every k only finitely often $\mathbf{d}(Y_s) \geq 2^{\frac{1}{k}|Y_s|}$ because in substage (a) the extension σ is chosen to be of minimal length, so $Y \notin S_{h_k}[\mathbf{d}]$. Hence $Y \in \text{DIM}^1$. \square

As RAND is a Σ_2^0 -class, we have the following corollary (which can also be proved by a direct construction).

Corollary 4.11. (Lutz [15]) *RAND is a proper subset of DIM^1 .*

In order to establish the existence of Δ_2^0 -computable sequences of any Δ_2^0 -computable dimension $\alpha \in [0, 1]$, Lutz [15] defined a *dilution function* $g_\alpha : \mathbf{C} \rightarrow \mathbf{C}$ that is computable and satisfies $\dim(g_\alpha(X)) = \alpha \cdot \dim(X)$ for all $X \in \mathbf{C}$. Applying this to any Δ_2^0 -computable Martin-Löf random sequence (which must have dimension 1) establishes the existence theorem. (We note that $g_\alpha(X)$ has the same Turing degree as X . Since by the Low Basis Theorem of Jockusch and Soare [18, Theorem V.5.32] there are Martin-Löf random sets of low degree, we immediately obtain that there are low sets of any Δ_2^0 -computable dimension α .) As g_α is continuous, it is a Wadge reduction from DIM^1 to DIM^α if $\alpha > 0$. Combining this with the previous theorem, we have that DIM^α is Wadge complete for Π_3^0 for all Δ_2^0 -computable $\alpha \in (0, 1)$. We now give a similar dilution construction that will allow us to prove this for *arbitrary* $\alpha \in (0, 1)$.

Let $X \in \mathbf{C}$ and let $\alpha \in (0, 1)$. Write $X = x_1 x_2 x_3 \dots$ where $|x_n| = 2n - 1$ for all n , noting that $|x_1 \cdots x_n| = n^2$. For each n , let

$$k_n = \left\lceil n \frac{1 - \alpha}{\alpha} \right\rceil$$

and $y_n = 0^{k_n}$. We then define

$$f_\alpha(X) = x_1 y_1 x_2 y_2 \cdots x_n y_n \cdots$$

Observe that f_α is a continuous function mapping \mathbf{C} to \mathbf{C} . We now show that it modifies the dimension of X in a controlled manner.

Lemma 4.12. *For any $X \in \mathbf{C}$ and $\alpha \in (0, 1)$,*

$$\dim(f_\alpha(X)) = \alpha \cdot \dim(X)$$

and

$$\text{Dim}(f_\alpha(X)) = \alpha \cdot \text{Dim}(X).$$

Proof. The proof uses (2.5) and (2.6), the Kolmogorov complexity characterizations of dimension and strong dimension.

Let $w \sqsubseteq f_\alpha(X)$. For some n ,

$$w = x_1 y_1 \cdots x_{n-1} y_{n-1} v,$$

where $v \sqsubseteq x_n y_n$. Then

$$\begin{aligned} K(w) &\leq K(x_1 \cdots x_{n-1}) + K(v) \\ &\quad + K(k_1) + \cdots + K(k_{n-1}) + O(1) \\ &\leq K(x_1 \cdots x_{n-1}) + O(n \log n). \end{aligned}$$

Because

$$|w| \geq |x_1 y_1 \cdots x_{n-1} y_{n-1}| \geq \frac{(n-1)^2}{\alpha},$$

we have

$$\frac{K(w)}{|w|} \leq \frac{\alpha \cdot K(x_1 \cdots x_{n-1})}{|x_1 \cdots x_{n-1}|} + \frac{O(n \log n)}{(n-1)^2},$$

It follows that

$$\begin{aligned} \dim(f_\alpha(X)) &\leq \alpha \liminf_{n \rightarrow \infty} \frac{K(x_1 \cdots x_{n-1})}{|x_1 \cdots x_{n-1}|} \\ &= \alpha \liminf_{n \rightarrow \infty} \frac{K(x \upharpoonright n)}{n} \\ &= \alpha \cdot \dim(X), \end{aligned}$$

where the first equality holds because the block x_n is short relative to $x_1 \cdots x_{n-1}$. Similarly, $\text{Dim}(f_\alpha(X)) \leq \alpha \cdot \text{Dim}(X)$.

For the other inequality, we have

$$\begin{aligned} K(x_1 \cdots x_{n-1}) &\leq K(w) + K(k_1) + \cdots + K(k_{n-1}) \\ &\quad + O(1) \\ &\leq K(w) + O(n \log n) \end{aligned}$$

and

$$|w| \leq |x_1 y_1 \cdots x_n y_n| \leq \frac{n^2}{\alpha} + n \leq \frac{(n+1)^2}{\alpha},$$

so

$$\begin{aligned} \frac{K(w)}{|w|} &\geq \alpha \frac{K(x_1 \cdots x_{n-1}) - O(n \log n)}{(n+1)^2} \\ &= \alpha \frac{K(x_1 \cdots x_{n-1})}{|x_1 \cdots x_{n-1}|} \frac{(n-1)^2}{(n+1)^2} - \frac{O(n \log n)}{(n+1)^2}. \end{aligned}$$

Therefore

$$\begin{aligned}
\dim(f_\alpha(X)) &\geq \alpha \liminf_{n \rightarrow \infty} \frac{K(x_1 \cdots x_{n-1})}{|x_1 \cdots x_{n-1}|} \\
&= \alpha \liminf_{n \rightarrow \infty} \frac{K(x \upharpoonright n)}{n} \\
&= \alpha \cdot \dim(X),
\end{aligned}$$

and analogously, $\text{Dim}(f_\alpha(X)) \geq \alpha \cdot \text{Dim}(X)$. \square

The function f_α establishes the completeness of DIM^α .

Theorem 4.13. *For all $\alpha \in (0, 1)$, DIM^α is Wadge complete for $\mathbf{\Pi}_3^0$. Therefore it is not a $\mathbf{\Sigma}_3^0$ -class, and in particular it is not a Σ_3^0 -class.*

Proof. By Lemma 4.12, f_α is a Wadge reduction from DIM^1 to DIM^α . Therefore DIM^α is Wadge complete for $\mathbf{\Pi}_3^0$ by composing f_α with the reduction from Theorem 4.10. \square

As g_α is also a Wadge reduction from $\text{DIM}_{\text{str}}^1$ to $\text{DIM}_{\text{str}}^\alpha$, we have from Theorem 4.6 that $\text{DIM}_{\text{str}}^\alpha$ is not a $\mathbf{\Sigma}_2^0$ -class for all $\alpha \in (0, 1)$. We now prove that $\text{DIM}_{\text{str}}^\alpha$ is not even $\mathbf{\Sigma}_3^0$ for all $\alpha \in [0, 1)$.

Theorem 4.14. *For all $\alpha \in [0, 1)$, $\text{DIM}_{\text{str}}^\alpha$ is Wadge complete for $\mathbf{\Pi}_3^0$. Therefore $\text{DIM}_{\text{str}}^\alpha$ is not a $\mathbf{\Sigma}_3^0$ -class, and in particular it is not a $\Sigma_3^0[\Pi_1^0]$ -class.*

Proof. The proof is similar to that of Theorem 4.10, but uses (2.6), the Kolmogorov complexity characterization of strong dimension. Let $\mathcal{C} = \bigcup_k \bigcap_s \mathcal{O}_{k,s}$ be a $\mathbf{\Sigma}_3^0$ -class and without loss of generality assume that $\mathcal{O}_{k,s} \supseteq \mathcal{O}_{k,s+1}$ for all k, s .

Let $\alpha \in (0, 1)$. (We will discuss the simpler case $\alpha = 0$ later.) We define a continuous function $f : \mathbf{C} \rightarrow \mathbf{C}$ in stages that will Wadge reduce \mathcal{C} to $\overline{\text{DIM}_{\text{str}}^\alpha}$. The image $Y = f(X)$ will be the unique sequence extending Y_s for all s . At stage 0 we define Y_0 to be the empty sequence.

At stage $s > 0$ we consider $X \upharpoonright s$, and define $k(s)$ as in the proof of Theorem 4.10. There are three substages.

Substage (a). First consider all strings ρ extending Y_{s-1} of minimal length with $K(\rho) \geq \alpha|\rho|$, and take the leftmost one of these ρ 's.

Substage (b). Next consider all strings σ extending ρ of minimal length with $K(\sigma) \geq (\alpha + \frac{1}{k(s)})|\sigma|$, and take the leftmost one of these σ 's. If $k(s)$ does not exist, let $\sigma = \rho$.

Substage (c). Extend σ with a block of 0's to obtain $Y_s = \sigma 0^{|\sigma|^2 - |\sigma|}$.

That is, to define Y_s , we first select ρ to increase the Kolmogorov complexity rate to α . This ensures that Y will have strong dimension at least α . We then construct a piece of evidence σ that Y has strong dimension at least $\alpha + \frac{1}{k(s)}$. We finish Y_s with a long block of 0's to bring the Kolmogorov complexity down to a near-zero rate, so that the next stage will work properly.

If $X \in \mathcal{C}$, then for the minimal k such that $X \in \bigcap_s \mathcal{O}_{k,s}$, infinitely many prefixes $\sigma \sqsubseteq Y$ satisfy $K(\sigma) \geq (\alpha + \frac{1}{k})|\sigma|$. Therefore $\text{Dim}(Y) \geq \alpha + \frac{1}{k}$, so $Y \notin \text{DIM}_{\text{str}}^\alpha$.

Now let $X \notin \mathcal{C}$. Let $\alpha' > \alpha$ be arbitrary, and choose k so that $\frac{1}{k} < \alpha' - \alpha$. Because $X \notin \mathcal{C}$, we have $k(s) > k$ for all sufficiently large s . Let s_0 be large enough to ensure $k(s) > s$ and $K(Y_{s-1}) \leq \sqrt{|Y_{s-1}|} + O(1) < \alpha|Y_{s-1}|$ hold for all $s \geq s_0$. Suppose that

$$K(w) \geq \alpha'|w|. \quad (4.2)$$

holds for some w with $Y_{s-1} \sqsubseteq w \sqsubseteq Y_s$ for some stage $s \geq s_0$. We then have that ρ is a proper extension of Y_{s-1} . By choice of ρ and σ and the fact that $\alpha' > \alpha + \frac{1}{k} > \alpha + \frac{1}{k(s)}$, we must have $w = \rho$ or $\sigma \sqsubseteq w$. We analyze these two cases separately.

- (i) $w = \rho$: Let ρ' be the string obtained from ρ by removing the last bit. Then $K(\rho) \leq K(\rho') + O(1)$. By choice of ρ , we have $K(\rho') < \alpha|\rho'|$. We also have $K(\rho) \geq (\alpha')|\rho|$ by (4.2). Putting these three statements together yields

$$\alpha'|\rho| < \alpha(|\rho| - 1) + O(1),$$

which is a contradiction if $|\rho| = |w|$ is sufficiently large.

- (ii) $\sigma \sqsubseteq w$: Obtain σ' from σ by removing the last bit of σ . Then we have

$$\begin{aligned} K(w) &\leq K(\sigma') + K(|w| - |\sigma|) + O(1) \\ &\leq K(\sigma') + \log(|w| - |\sigma|) + O(1) \\ &\leq K(\sigma') + 2\log|\sigma| + O(1). \end{aligned}$$

By choice of σ , $K(\sigma') < (\alpha + \frac{1}{k(s)})|\sigma'|$. These two facts together with (4.2) tell us that

$$\alpha'|w| < \left(\alpha + \frac{1}{k(s)} \right) (|\sigma| - 1) + 2\log|\sigma| + O(1),$$

which is a contradiction for large $|w|$ because $|w| \geq |\sigma|$ and $\alpha' > \alpha + \frac{1}{k(s)}$.

Therefore, for all sufficiently long $w \sqsubseteq Y$, (4.2) does not hold. It follows that $\text{Dim}(Y) \leq \alpha$. On the other hand, there are infinitely many $\rho \sqsubseteq Y$ with $K(\rho) \geq \alpha|\rho|$, so $\text{Dim}(Y) \geq \alpha$. Therefore $Y \in \text{DIM}_{\text{str}}^\alpha$.

This shows that f is a Wadge reduction from \mathcal{C} to $\overline{\text{DIM}_{\text{str}}^\alpha}$. As \mathcal{C} is an arbitrary Σ_3^0 -class, this shows that $\text{DIM}_{\text{str}}^\alpha$ is Wadge complete for Π_3^0 .

The proof for the case $\alpha = 0$ is similar, but simpler as substage (a) is omitted in the construction. \square

4.3 Ad Hoc Methods

When classifying classes in the arithmetical hierarchy of reals there are several methods one can use. As we have seen, category methods are sometimes useful up to the second level, Wadge reductions are useful if the classification in the effective (lightface) hierarchy coincides with that in the classical (boldface) hierarchy, and sometimes (as in Proposition 3.1 and Example 3.3) one just needs something else. In particular when the level of the class in the effective hierarchy is not the same as the level in the classical hierarchy one often needs to resort to ad hoc arguments. One might think that the notion of effective Wadge reduction, or recursive functional, would be the proper notion to use in classifying classes of reals in the effective hierarchy. However, this notion is rarely useful for the following reason. Let \mathcal{X} be a class without computable elements, such as the class of Martin-Löf random sets or the class of 1-generic sets. Then \mathcal{X} cannot be proven to be complete for any level of the effective hierarchy by a recursive Wadge reduction f . For if X is recursive, then so is $f(X)$, so we can never have $X \in \mathbf{C} \iff f(X) \in \mathcal{X}$. So we see that “easy”

classes like \mathbf{C} that contain recursive elements cannot be reduced in such a way to many “difficult” classes, which renders the notion rather useless.

We have left open the question whether $\text{DIM}_{\text{str}}^1$ is not in Π_2^0 , and whether $\text{DIM}_{\text{str}}^\alpha$ is not in Π_3^0 for any Δ_2^0 -computable $\alpha \in [0, 1)$. We have no answer to the second question, but we provide an answer to the first in the next theorem. We make use of the following lemma.

Lemma 4.15. *If $\mathcal{X} \in \Pi_2^0$ is dense then there is a computable $X \in \mathcal{X}$.*

Proof. This is an easy finite extension argument. Suppose that $\mathcal{X} = \{X : (\forall m)(\exists k)R^X(m, k) \downarrow = 1\} \in \Pi_2^0$ is dense. (Here R is a computable predicate. Note that R does not have to be defined with oracles X that are not in \mathcal{X} .) Given any initial segment τ such that

$$(\forall n < m)(\exists k)R^\tau(m, k) \downarrow = 1,$$

we show how to compute an extension $\sigma \sqsupset \tau$ such that

$$(\exists k)R^\sigma(m, k) \downarrow = 1. \quad (4.3)$$

Because \mathcal{X} is dense, there are $X \sqsupset \tau$ and k such that $R^X(m, k) \downarrow = 1$. Let u be the use of this computation, i.e. the part of the oracle X used in it. Now define $\sigma = \max\{X \upharpoonright u, \tau\}$. Then $\sigma \sqsupseteq \tau$ satisfies (4.3).

Now it is clear that for every m we can compute appropriate extensions σ_m such that $X = \bigcup_m \sigma_m$ is computable and $(\forall m)(\exists k)R^{\sigma_m}(m, k) \downarrow = 1$, so that $X \in \mathcal{X}$. \square

Theorem 4.16. *$\text{DIM}_{\text{str}}^1$ is not a Π_2^0 -class. Hence it is properly $\Pi_2^0[\Pi_1^0]$.*

Proof. Suppose that $\text{DIM}_{\text{str}}^1$ is Π_2^0 . Then, since clearly $\text{DIM}_{\text{str}}^1$ is dense, by Lemma 4.15 it contains a computable real, contradicting that every computable real has strong dimension 0. \square

We conclude this section by summarizing its main results in the following table.

	DIM^α	$\text{DIM}_{\text{str}}^\alpha$
$\alpha = 0$	$\Pi_2^0 - \Sigma_2^0$	$\Pi_3^0[\Sigma_1^0] - \Sigma_3^0$
$\alpha \in (0, 1) \cap \Delta_2^0$	$\Pi_3^0 - \Sigma_3^0$	$\Pi_3^0[\Sigma_1^0] - \Sigma_3^0$
$\alpha = 1$	$\Pi_3^0 - \Sigma_3^0$	$\Pi_2^0[\Pi_1^0] - (\Sigma_2^0 \cup \Pi_2^0)$
arbitrary $\alpha \in (0, 1)$	$\Pi_3^0 - \Sigma_3^0$	$\Pi_3^0 - \Sigma_3^0$

Question 4.17. *Is it the case that $\text{DIM}_{\text{str}}^\alpha$ is not in Π_3^0 for any Δ_2^0 -computable $\alpha \in [0, 1)$?*

5 Effective Randomness Classes

We begin this section by pointing out some relationships between computable dimension, Church randomness, and Schnorr randomness.

Analogously to what was done for the constructive case, the *computable dimension* of a class $\mathcal{A} \subseteq \mathbf{C}$ is defined as

$$\dim_{\text{comp}}(\mathcal{A}) = \inf \left\{ s \mid \begin{array}{l} \text{there exists a computable} \\ s\text{-gale } d \text{ for which } \mathcal{A} \subseteq S^\infty[d] \end{array} \right\}.$$

A *selection rule* is a function $\varphi : \{0, 1\}^* \rightarrow \{0, 1\}$. With every selection rule φ we associate a function $\Phi : \{0, 1\}^* \rightarrow \{0, 1\}^*$ defined by $\Phi(\lambda) = \lambda$ and

$$\Phi(wi) = \begin{cases} \Phi(w)i & \text{if } \varphi(w) = 1, \\ \Phi(w) & \text{if } \varphi(w) = 0. \end{cases}$$

A set A is called *Church random* if every substring of χ_A (the characteristic string of A) defined by a computable selection rule is stochastic, i.e., satisfies the law of large numbers. Consider the following property of selection rules:

$$\inf_{w \in \{0, 1\}^*} \frac{|\Phi(w)|}{|w|} > 0. \quad (5.1)$$

A *computable null set of exponential order* is a set of the form $S^{a^n}[d]$, where d is a computable martingale and $a > 1$. It is easy to check that a set is not in any computable null set of exponential order if and only if $\{A\}$ has computable dimension 1. With this observation, we can restate a result of Schnorr as follows.

Theorem 5.1. (Schnorr [20, Satz 17.8]) *$\{A\}$ has computable dimension 1 if and only if every substring of χ_A selected by a computable selection rule with the property (5.1) is stochastic.*

In particular every Church random sequence is not in any null set of the form $S^{a^n}[d]$ where d is computable. In the words of Schnorr [21], “Church random sequences approximate the behavior of Schnorr random sequences.”

Proposition 5.2. *There are sequences with computable dimension 1 that are not Church random.*

Proof. Let R be computably random, and let $D = \{2^n \mid n \in \mathbb{N}\}$ be an exponentially sparse decidable domain. Then $A = R - D$ has computable dimension 1, but D can be computably selected, so A is not Church random. \square

We now classify the Schnorr random sequences in the arithmetical hierarchy.

Theorem 5.3. $\text{RAND}_{\text{Schnorr}}$ *is a Π_3^0 -class, but not a Σ_3^0 -class.*

Proof. First note that $\text{RAND}_{\text{Schnorr}} \in \Pi_3^0$: $X \in \text{RAND}_{\text{Schnorr}}$ if and only if for every pair of codes e and f , either the e -th partial computable function φ_e is not a computable order (i.e. is not total or decreases at some point), φ_f is not a computable martingale (i.e. is not total or violates the martingale property at some point), or $X \notin S^{\varphi_e}[\varphi_f]$, and that every one of these options is Σ_2^0 .

The rest of the proof resembles that of Theorem 4.10. Fix a (non-computable) sequence of computable martingales $\{d_k\}_{k \in \mathbb{N}}$ and a sequence of computable orders $\{h_k\}_{k \in \mathbb{N}}$ such that

- (i) $X \in \text{RAND}_{\text{Schnorr}} \iff \forall k (X \notin S^{h_k}[d_k]).$
- (ii) $S^{h_k}[d_k] - S^{\min\{h_j : j < k\}}[\sum_{j < k} d_j]$ is dense for every k .

The d_k can be defined by taking appropriate sums of computable martingales so that for any computable martingale d , there is some d_k such that $d_k(w) \geq d(w)$ for all w . For the h_k one can take any family of computable orders such that every computable order h dominates some h_k . (Of course the d_k and h_k cannot be uniformly computable families, but that is of no concern to us.)

Let $\bigcup_k \bigcap_s \mathcal{O}_{k,s}$ be a Σ_3^0 -class. We define a continuous function $f : \mathbf{C} \rightarrow \mathbf{C}$ such that

$$\forall k \left(X \in \bigcap_s \mathcal{O}_{k,s} \iff f(X) \in S^{h_k}[d_k] \right) \quad (5.2)$$

so that by (i) we have $X \notin \bigcup_k \bigcap_s \mathcal{O}_{k,s} \iff f(X) \in \text{RAND}_{\text{Schnorr}}$.

As in the proof of Theorem 4.10 we define the image $Y = f(X)$ in stages. Every time we find a new piece of evidence that $X \in \bigcap_s \mathcal{O}_{k,s}$, at stage s say, we build a piece of evidence that $Y \in S^{h_k}[d_k]$ by choosing an appropriate finite extension at stage s . Such an extension can be found by (ii). The rest of the proof is identical to that of Theorem 4.10. \square

With only some obvious changes one can also prove the following theorem.

Theorem 5.4. $\text{RAND}_{\text{comp}}$ is a Π_3^0 -class, but not a Σ_3^0 -class.

Proof. Note that X is computably random if and only if for every e , φ_e is not a computable martingale or $X \notin S^\infty[\varphi_e]$, so the class is Π_3^0 . That it is properly Π_3^0 is actually easier than the proof of Theorem 5.3 since we only need the sequence $\{d_k\}$ and not the $\{h_k\}$. \square

In contrast to the universal constructive supermartingale \mathbf{d} satisfying $\text{RAND} = \overline{S^\infty[\mathbf{d}]}$, Theorem 5.4 implies that, even from a noncomputable standpoint, $\text{RAND}_{\text{comp}}$ has no such universal object. That is, $\text{RAND}_{\text{comp}} \neq \overline{S^\infty[d]}$ for any (arbitrarily noncomputable) supermartingale d , as otherwise $\text{RAND}_{\text{comp}}$ would be a Σ_2^0 -class.

In this paper we have considered only the extension of the arithmetical hierarchy of reals by adding one local quantifier. We end by remarking that one can add of course more local quantifiers. The classes thus obtained also have natural inhabitants. To give an example, again from the theory of randomness, recall that a set A is n -random if it is Martin-Löf random relative to $\emptyset^{(n-1)}$. So it is 1-random if it is Martin-Löf random, 2-random if it is Martin-Löf random relative to K , etc. Now the class of n -random sets is Σ_2^0 for every n , and in fact one can check that it is $\Sigma_2^0[\Sigma_{n-1}^0]$.

Acknowledgment. The second and third authors thank Alekos Kechris for a helpful discussion.

References

- [1] K. Ambos-Spies, W. Merkle, J. Reimann, and F. Stephan. Hausdorff dimension in exponential time. In *Proceedings of the 16th IEEE Conference on Computational Complexity*, pages 210–217. IEEE Computer Society, 2001.
- [2] K. B. Athreya, J. M. Hitchcock, J. H. Lutz, and E. Mayordomo. Effective strong dimension in algorithmic information and computational complexity. In *Proceedings of the 21st Annual Symposium on Theoretical Aspects of Computer Science*, pages 632–643. Springer-Verlag, 2004.
- [3] J. J. Dai, J. I. Lathrop, J. H. Lutz, and E. Mayordomo. Finite-state dimension. *Theoretical Computer Science*, 310(1–3):1–33, 2004.
- [4] S. A. Fenner. Gales and supergales are equivalent for defining constructive Hausdorff dimension. Technical Report cs.CC/0208044, Computing Research Repository, 2002.

- [5] L. Fortnow and J. H. Lutz. Prediction and dimension. *Journal of Computer and System Sciences*. To appear.
- [6] F. Hausdorff. Dimension und äußeres Maß. *Mathematische Annalen*, 79:157–179, 1919.
- [7] J. M. Hitchcock. MAX3SAT is exponentially hard to approximate if NP has positive dimension. *Theoretical Computer Science*, 289(1):861–869, 2002.
- [8] J. M. Hitchcock. Fractal dimension and logarithmic loss unpredictability. *Theoretical Computer Science*, 304(1–3):431–441, 2003.
- [9] J. M. Hitchcock. Gales suffice for constructive dimension. *Information Processing Letters*, 86(1):9–12, 2003.
- [10] C. G. Jockusch. Degrees of generic sets. In *Recursion Theory: its Generalizations and Applications*, volume 45 of *London Mathematical Society Lecture Notes Series*, pages 110–139. Cambridge University Press, 1980.
- [11] H. Ki and T. Linton. Normal numbers and subsets of \mathbb{N} with given densities. *Fundamenta Mathematicae*, 144:163–179, 1994.
- [12] G. Kreisel. Note on arithmetical models for consistent formulae of the predicate calculus. *Fundamenta Mathematicae*, 37:265–285, 1950.
- [13] M. Li and P. M. B. Vitányi. *An Introduction to Kolmogorov Complexity and its Applications*. Springer-Verlag, Berlin, 1997. Second Edition.
- [14] J. H. Lutz. Dimension in complexity classes. *SIAM Journal on Computing*, 32(5):1236–1259, 2003.
- [15] J. H. Lutz. The dimensions of individual strings and sequences. *Information and Computation*, 187(1):49–79, 2003.
- [16] P. Martin-Löf. The definition of random sequences. *Information and Control*, 9:602–619, 1966.
- [17] E. Mayordomo. A Kolmogorov complexity characterization of constructive Hausdorff dimension. *Information Processing Letters*, 84(1):1–3, 2002.
- [18] P. Odifreddi. *Classical recursion theory*, volume 125 of *Studies in Logic and the Foundations of Mathematics*. North-Holland, 1989.
- [19] H. Rogers, Jr. *Theory of Recursive Functions and Effective Computability*. McGraw - Hill, New York, N.Y., 1967.
- [20] C. P. Schnorr. Zufälligkeit und Wahrscheinlichkeit. *Lecture Notes in Mathematics*, 218, 1971.
- [21] C. P. Schnorr. A survey of the theory of random sequences. In R. E. Butts and J. Hintikka, editors, *Basic Problems in Methodology and Linguistics*, pages 193–211. D. Reidel, 1977.
- [22] L. Staiger. Recursive automata on infinite words. In *Proceedings of the 10th Annual Symposium on Theoretical Aspects of Computer Science*, pages 629–639. Springer-Verlag, 1993.

- [23] L. Staiger. On the power of reading the whole infinite input tape. In C. S. Calude and Gh. Paun, editors, *Finite Versus Infinite: Contributions to an Eternal Dilemma*, pages 335–348. Springer-Verlag, 2000.
- [24] L. Staiger. Constructive dimension equals Kolmogorov complexity. Technical Report CDMTCS-210, University of Auckland, January 2003.
- [25] S. A. Terwijn. Complexity and randomness. Technical Report CDMTCS-212, University of Auckland, March 2003. Notes for a course given at the University of Auckland.
- [26] M. van Lambalgen. *Random Sequences*. PhD thesis, Department of Mathematics, University of Amsterdam, 1987.
- [27] J. Ville. *Étude Critique de la Notion de Collectif*. Gauthier–Villars, Paris, 1939.
- [28] Y. Wang. *Randomness and Complexity*. PhD thesis, Department of Mathematics, University of Heidelberg, 1996.