

CHARACTERS AND GROWTH OF ADMISSIBLE REPRESENTATIONS OF REDUCTIVE p -ADIC GROUPS

RALF MEYER AND MAARTEN SOLLEVELD

ABSTRACT. We use coefficient systems on the affine Bruhat–Tits building to study admissible representations of reductive p -adic groups in characteristic not equal to p . We show that the character function is locally constant and provide explicit neighbourhoods of constancy. We estimate the growth of the subspaces of invariants for compact open subgroups.

MATHEMATICS SUBJECT CLASSIFICATION (2010). 20G25, 20E42

CONTENTS

1. Introduction	1
2. The structure of reductive algebraic groups	3
3. Some Bruhat–Tits theory	6
4. Fixed points in the building	9
5. The groups $U_\Omega^{(e)}$	14
6. Characters of admissible representations	19
7. The local constancy of characters	23
8. A bound for the dimension of V^K	28
9. Conclusion	33
References	34

1. INTRODUCTION

Let \mathbb{F} be a non-Archimedean local field, possibly of nonzero characteristic, and let G be a reductive algebraic group over \mathbb{F} , briefly called a reductive p -adic group. Let π be an admissible representation of G on a complex vector space V . Since V^K has finite dimension for every compact open subgroup $K \subseteq G$, the operator $\pi(f)$ has finite rank for all test functions f . The resulting distribution $\theta_\pi(f) := \text{tr}(\pi(f), V)$ is called the *character* of π . Since V usually has infinite dimension, the operators $\pi(g)$ need not be trace-class for $g \in G$. Nevertheless, Harish-Chandra could show that the character is described by a locally integrable function:

Theorem 1.1 (Harish-Chandra). *Let $\pi: G \rightarrow \text{Aut}(V)$ be an admissible representation of a reductive p -adic group.*

- (a) *The operator $\pi(g)$ has a well-defined trace $\text{tr}_\pi(g)$ when g belongs to the set G_{rss} of regular semisimple elements.*
- (b) *The function $\text{tr}_\pi: G_{\text{rss}} \rightarrow \mathbb{C}$ is locally constant.*
- (c) *The function tr_π , extended by 0 on $G \setminus G_{\text{rss}}$, is locally integrable with respect to the Haar measure μ on G , and for any test function f ,*

$$\theta_\pi(f) = \int_G f(g) \text{tr}_\pi(g) d\mu(g).$$

Supported by the German Research Foundation (Deutsche Forschungsgemeinschaft (DFG)) through the Institutional Strategy of the University of Göttingen.

(d) Let $D(g)$ for $g \in G_{\text{rss}}$ be the determinant of $\text{Ad}(g) - 1$ acting on $\text{Lie}(G) / \text{Lie}(T)$ for a maximal torus T in G containing g . The function $G \ni g \mapsto |D(g)|^{1/2} \text{tr}_\pi(g)$ is locally bounded.

The original proof of this deep theorem is distributed over various papers of Harish-Chandra collected in [7]. A complete account of it can be found in [8]. The proofs of (c) and (d) use the exponential mapping for G , which only works well if the characteristic of \mathbb{F} is zero. It is reasonable to expect that (c) and (d) are valid in non-zero characteristic as well, but the authors are not aware of a proof. According to [24, paragraph E.4.4] Harish-Chandra's proof of (a) and (b) remains valid if one replaces \mathbb{C} by an algebraically closed field of characteristic unequal to p .

In this article we generalise part of Theorem 1.1 to representations on modules over unital rings in which p is invertible. In this purely algebraic setting, we can only define the character as a function where it is locally constant. To prove (a) and (b), we describe explicit neighbourhoods on which tr_π is constant. In characteristic 0, similar results are due to Adler and Korman [1].

Parts (c) and (d) seem specific to real or complex representations because they involve analysis. Unfortunately, our methods are insufficient to (re)prove them, as we discuss in the last section.

As a substitute we estimate the dimension of invariant subspaces V^K for certain compact open subgroups K in G . The authors have not found growth estimates for these dimensions in the literature. Since V^K is the range of an idempotent $\langle K \rangle$ in the Hecke algebra associated to K , we get

$$\dim V^K = \frac{1}{|K|} \int_K \text{tr}_\pi(g) d\mu(g).$$

But the estimate in (d) is not strong enough to control these integrals.

Our methods are of a geometric nature and involve the affine building of G . Thus we will make extensive use of Bruhat–Tits theory, including some hard parts. At the same time, we use only little representation theory. Both of our main results use the resolutions constructed by Schneider and Stuhler [18]. These resolutions are based on a family of compact open subgroups $U_x^{(e)}$ for $e \in \mathbb{N}$, indexed by vertices of the affine Bruhat–Tits building. These generate subgroups $U_\sigma^{(e)}$ indexed by polysimplices in the building. The invariant subspaces $V^{U_\sigma^{(e)}}$ in an admissible representation V form a locally finite-dimensional coefficient system on the building. It is shown in [11] that this coefficient system is acyclic on any convex subcomplex of the building. In particular, it provides a resolution of V of finite type.

Here we need acyclicity also for finite subcomplexes of the building because this provides chain complexes of finite-dimensional vector spaces, which are used in [11] to express the character of V as a sum over contributions of polysimplices in the building. We use this formula to find for each regular semisimple element γ and each vertex x in the building a number r such that the character is constant on $U_x^{(r)}\gamma$; the constant r depends on the distance between x and a subset of the building corresponding to the maximal torus containing γ , on the (ir)regularity of γ , and on the level of the representation V , that is, on the smallest $e \in \mathbb{N}$ such that V is generated by the $U_y^{(e)}$ -invariants for all vertices y .

Along the way, we also prove some auxiliary results that may be useful in other contexts. We prove that the parabolic subgroup contracted by an element of a reductive p -adic group is indeed parabolic and, in particular, algebraic (Proposition 2.3). We describe which points in the building are fixed by a semisimple element in Section 4. We establish that the level of representations is preserved by Jacquet induction and restriction (Proposition 5.8). The relationship between character function and distribution is made precise in an algebraic setting in Section 6.

2. THE STRUCTURE OF REDUCTIVE ALGEBRAIC GROUPS

We fix our notation and recall some general facts from the theory of linear algebraic groups. Nothing in this section is new and most of it can be found in several textbooks, for example [20].

Let \mathcal{G} be a linear algebraic group defined over a field \mathbb{F} . The collections of characters and cocharacters of \mathcal{G} are denoted by $X^*(\mathcal{G})$ and $X_*(\mathcal{G})$, respectively. Let $G := \mathcal{G}(\mathbb{F})$ be its group of \mathbb{F} -rational points. By definition, an algebraic (co)character of G is a (co)character of \mathcal{G} that is defined over \mathbb{F} . The corresponding sets are denoted by $X^*(G)$ and $X_*(G)$. Let $\mathcal{Z}(\mathcal{G})$ be the centre of \mathcal{G} and let $\mathcal{Z}_c(\mathcal{G})$ be the maximal connected algebraic subgroup of $\mathcal{Z}(\mathcal{G})$. We denote the centraliser in \mathcal{G} of an element $g \in G$ by $\mathcal{Z}_{\mathcal{G}}(g)$.

We will assume throughout that \mathcal{G} is connected and reductive. An algebraic subgroup \mathcal{P} of \mathcal{G} is *parabolic* if \mathcal{G}/\mathcal{P} is a complete algebraic variety. We denote the unipotent radical of \mathcal{P} by $\mathcal{R}_u(\mathcal{P})$. A *Levi factor* of \mathcal{P} is a reductive subgroup \mathcal{M} such that $\mathcal{P} = \mathcal{M} \ltimes \mathcal{R}_u(\mathcal{P})$.

We write $Z(G)$, $Z_c(G)$, P , $R_u(P)$, and M for the groups of \mathbb{F} -points of $\mathcal{Z}(\mathcal{G})$, $\mathcal{Z}_c(\mathcal{G})$, \mathcal{P} , $\mathcal{R}_u(\mathcal{P})$, and \mathcal{M} , respectively. We denote the space of \mathbb{F} -points of the Lie algebra of \mathcal{G} by $\text{Lie}_{\mathbb{F}}(\mathcal{G})$.

We say that an algebraic torus \mathcal{T} *splits over \mathbb{F}* if $\mathcal{T}(\mathbb{F}) \cong (\mathbb{F}^\times)^{\dim \mathcal{T}}$ as \mathbb{F} -groups. We say that G *splits* (over \mathbb{F}) if there is a maximal torus \mathcal{T} of \mathcal{G} that splits over \mathbb{F} .

Proposition 2.1. *There is a finite Galois extension of \mathbb{F} over which \mathcal{G} splits.*

Proof. For tori this was first proven by Ono [14, Proposition 1.2.1]. This implies the result for general reductive groups. \square

Let \mathcal{S} be maximal among the tori in \mathcal{G} that split over \mathbb{F} and let $S := \mathcal{S}(\mathbb{F})$. We call S a *maximal split torus* in G . Notice that every algebraic (co)character of \mathcal{S} is defined over \mathbb{F} , as S is split. Let $\Phi = \Phi(\mathcal{G}, \mathcal{S}) \subset X^*(\mathcal{S})$ be the root system of \mathcal{G} with respect to \mathcal{S} , and let $\Phi^\vee \subset X_*(\mathcal{S})$ be the dual root system. Let $\mathcal{Z}_{\mathcal{G}}(\mathcal{S})$ and $N_{\mathcal{G}}(\mathcal{S})$ denote the centraliser and the normaliser of \mathcal{S} in \mathcal{G} and let $Z_G(S)$ and $N_G(S)$ be their groups of \mathbb{F} -points. The *Weyl group* of Φ is

$$W(\Phi) := N_G(S) / Z_G(S).$$

The root system Φ need not be reduced if G is not split. The corresponding reduced root system is

$$(1) \quad \Phi^{\text{red}} := \{\alpha \in \Phi(\mathcal{G}, \mathcal{S}) : \alpha/2 \notin \Phi(\mathcal{G}, \mathcal{S})\}.$$

For every root $\alpha \in \Phi(\mathcal{G}, \mathcal{S})$ there is a unipotent algebraic subgroup $\mathcal{U}_\alpha \subset \mathcal{G}$ with group of \mathbb{F} -points U_α , characterised by the following two conditions:

- $\mathcal{Z}_{\mathcal{G}}(\mathcal{S})$ normalises \mathcal{U}_α ,
- $\text{Lie}_{\mathbb{F}}(\mathcal{U}_\alpha)$ is the sum of the S -weight spaces for α and 2α , with respect to the adjoint action of S on $\text{Lie}_{\mathbb{F}}(\mathcal{G})$.

If $\alpha, 2\alpha \in \Phi$ then $U_{2\alpha} \subsetneq U_\alpha$, and it is convenient to write $U_{2\alpha} = \{1\}$ if $\alpha \in \Phi$ but $2\alpha \notin \Phi$. The groups $U_\alpha/U_{2\alpha}$ and $U_{2\alpha}$ are naturally endowed with the structure of an \mathbb{F} -vector space and are isomorphic to their respective Lie algebras. The subset $\bigcup_{\alpha \in \Phi^{\text{red}}} U_\alpha \cup Z_G(S)$ generates the group G .

Let Φ^+ be a system of positive roots in Φ and let $\Delta \subseteq \Phi^{\text{red}}$ be the corresponding basis. Any subset $D \subseteq \Delta$ is a basis of a root system $\Phi_D := \mathbb{Z}D \cap \Phi$. The algebraic subgroup \mathcal{P}_D of \mathcal{G} generated by $\mathcal{Z}_{\mathcal{G}}(\mathcal{S})$ and the \mathcal{U}_α with $\alpha \in \Phi_D \cup \Phi^+$ is parabolic. Its unipotent radical is generated by the \mathcal{U}_α with $\alpha \in \Phi^+ \setminus \Phi_D^+$. The group \mathcal{M}_D that is generated by $\bigcup_{\alpha \in \Phi_D} \mathcal{U}_\alpha \cup \mathcal{Z}_{\mathcal{G}}(\mathcal{S})$ is a Levi subgroup of \mathcal{P}_D . Moreover,

$\mathcal{M}_D = \mathcal{Z}_{\mathcal{G}}(\mathcal{S}_D)$, where \mathcal{S}_D is the connected component of

$$\bigcap_{\alpha \in \Phi_D} \ker \alpha \subseteq \mathcal{Z}(\mathcal{M}_D).$$

We note that \mathcal{P}_{\emptyset} is a Borel subgroup of \mathcal{G} , that $\mathcal{P}_{\Delta} = \mathcal{M}_{\Delta} = \mathcal{G}$, and that $\mathcal{S}_{\Delta}(\mathbb{F})$ is the unique maximal split torus of $\mathcal{Z}(\mathcal{G})$.

Definition 2.2. Groups of the form \mathcal{P}_D are called *standard parabolic* (with respect to \mathcal{S} and Φ^+).

Every parabolic subgroup of \mathcal{G} is conjugate to exactly one standard parabolic subgroup. Let $\Phi^- := -\Phi^+$ be the set of negative roots and let $\bar{\mathcal{P}}_D$ be subgroup of \mathcal{G} generated by $\mathcal{Z}_{\mathcal{G}}(\mathcal{S})$ and the \mathcal{U}_{α} with $\alpha \in \Phi_D \cup \Phi^-$. The parabolic subgroup $\bar{\mathcal{P}}_D$ is *opposite* to \mathcal{P}_D in the sense that $\mathcal{P}_D \cap \bar{\mathcal{P}}_D = \mathcal{M}_D$ is a Levi subgroup of both. Moreover

$$\mathrm{Lie}_{\mathbb{F}}(\mathcal{G}) = \mathrm{Lie}_{\mathbb{F}}(\mathcal{R}_{\mathrm{u}}(\mathcal{P}_D)) \oplus \mathrm{Lie}_{\mathbb{F}}(\mathcal{M}_D) \oplus \mathrm{Lie}_{\mathbb{F}}(\mathcal{R}_{\mathrm{u}}(\bar{\mathcal{P}}_D)).$$

We shall also need the *pseudo-parabolic* subgroup

$$(2) \quad P(\chi) := \{p \in G : \lim_{\lambda \rightarrow 0} \chi(\lambda) p \chi(\lambda)^{-1} \text{ exists}\}$$

for an algebraic cocharacter $\chi: \mathbb{F}^{\times} \rightarrow G$. This limit is meant purely algebraically, by definition it exists if and only if the corresponding map $\mathbb{F}^{\times} \rightarrow G$ extends to an algebraic morphism $\mathbb{F} \rightarrow G$. In a reductive group, any pseudo-parabolic subgroup is the group of \mathbb{F} -points of a parabolic subgroup by [20, Lemma 15.1.2].

From now on we assume that the field \mathbb{F} is endowed with a non-trivial discrete valuation $v: \mathbb{F} \rightarrow \mathbb{Q} \cup \{\infty\}$. We fix a real number $q > 1$ and we define a metric on \mathbb{F} by

$$d(\lambda, \mu) = q^{-v(\lambda - \mu)}.$$

Via an embedding $\mathcal{G} \rightarrow \mathrm{GL}_n$, the metric d yields a metric on $G = \mathcal{G}(\mathbb{F})$ as well. Even though there is no unique way to do this, the resulting collection of bounded subsets of G is canonical. This bornology on G is compatible with the group structure, in the sense that $B_1^{-1}B_2$ is bounded for all bounded subsets B_1 and B_2 of G .

It follows directly from the properties of a valuation that every finitely generated subgroup of $(\mathbb{F}, +)$ is bounded, and this implies that every unipotent element of G generates a bounded subgroup.

Following Deligne [6], we assign to any $g \in G$ the *parabolic subgroup contracted by g* ,

$$(3) \quad P_g := \{p \in G : \{g^n pg^{-n} : n \in \mathbb{N}\} \text{ is bounded}\},$$

and

$$(4) \quad M_g := P_g \cap P_{g^{-1}} = \{p \in G : \{g^n pg^{-n} : n \in \mathbb{Z}\} \text{ is bounded}\}.$$

The following result, which will be needed in Section 7.2, was proved in [15, Lemma 2] under the additional assumptions that \mathcal{G} is semisimple and almost \mathbb{F} -simple. Although it is apparently well-known that it holds for general reductive groups, the authors have not found a good reference for this.

Proposition 2.3. *The subgroups P_g and M_g for $g \in G$ have the following properties:*

- (a) P_g is a parabolic subgroup of G .
- (b) $\mathcal{R}_{\mathrm{u}}(P_g) = \{p \in G : \lim_{n \rightarrow \infty} g^n pg^{-n} = 1\}$.
- (c) The parabolic subgroup $P_{g^{-1}}$ is opposite to P_g and M_g is a Levi subgroup of P_g .
- (d) $g\mathcal{Z}(M_g)$ is contained in a bounded subgroup of $M_g / \mathcal{Z}(M_g)$.

Proof. We first establish (a). Clearly, P_g is a subgroup of G that contains g . The difficulty is to show that P_g is an algebraic subgroup of G , although it is defined in topological terms. Choose a finite extension field \mathbb{F}_g of \mathbb{F} which contains the roots of the characteristic polynomial of g . Then we have a Jordan decomposition $g = g_s g_u = g_u g_s$ in $\mathcal{G}(\mathbb{F}_g)$, see [20, Section 2.4]. Let \mathcal{T} be a maximal torus in \mathcal{G} defined over \mathbb{F}_g that contains g_s , and let $\tilde{\mathbb{F}}$ be a finite extension field of \mathbb{F}_g over which \mathcal{T} splits (Proposition 2.1). We may and will assume that $\tilde{\mathbb{F}}$ is normal over \mathbb{F} . According to [19, Section I.4] the valuation v extends to a valuation \tilde{v} on $\tilde{\mathbb{F}}$. We abbreviate $\mathcal{G}(\tilde{\mathbb{F}}) = \tilde{G}$, and similarly for its algebraic subgroups. Let $\tilde{\Phi}$ be the root system of \mathcal{G} with respect to \mathcal{T} .

Since g_u is unipotent, $\tilde{K} := \{g_u^n : n \in \mathbb{Z}\}$ is a bounded subgroup of \tilde{G} , and it centralises g_s . For $\alpha \in \tilde{\Phi}$ and $p \in \tilde{U}_\alpha \setminus \{1\}$, the following are equivalent:

- $\{g^n pg^{-n} : n \in \mathbb{N}\}$ is bounded,
- $\tilde{K} \{g_s^n pg_s^{-n} : n \in \mathbb{N}\} \tilde{K}$ is bounded,
- $\{g_s^n pg_s^{-n} : n \in \mathbb{N}\}$ is bounded,
- $g_s p g_s^{-1} = \lambda p$ with $\{\lambda^n : n \in \mathbb{N}\} \subseteq \tilde{\mathbb{F}}$ bounded,
- $\tilde{v}(\alpha(g_s)) \geq 0$.

We may choose a system of positive roots $\tilde{\Phi}^+$ with $\tilde{v}(\alpha(g_s)) \geq 0$ for all $\alpha \in \tilde{\Phi}^+$. Let $D \subseteq \tilde{\Delta}$ be the set of simple roots with $\tilde{v}(\alpha(g_s)) = 0$. The group \tilde{P}_g is generated by $\tilde{T} := \mathcal{T}(\tilde{\mathbb{F}})$ and all \tilde{U}_α with $\alpha \in \tilde{\Phi}^+ \cup \Phi_D$. Thus \tilde{P}_g is the group of $\tilde{\mathbb{F}}$ -points of the parabolic subgroup \mathcal{P}_D of \mathcal{G} , and the collection of non-zero weights of \tilde{T} in $\text{Lie}_{\tilde{\mathbb{F}}}(\mathcal{P}_D)$ equals

$$(5) \quad \{\alpha \in \tilde{\Phi} : \tilde{v}(\alpha(g_s)) \geq 0\} =: \Phi(\mathcal{P}_g, \mathcal{T}).$$

As mentioned above, \tilde{P}_g is also a pseudo-parabolic subgroup of \tilde{G} , so there is a cocharacter $\tilde{\chi} \in X_*(\mathcal{G})$ with $\tilde{P}_g = \tilde{P}(\tilde{\chi})$. In fact, any $\tilde{\chi} \in X_*(\tilde{\mathcal{T}})$ with

$$(6) \quad \{\alpha \in \tilde{\Phi} : \langle \alpha, \tilde{\chi} \rangle \geq 0\} = \Phi(\mathcal{P}_g, \mathcal{T})$$

will do. To prove that $P_g = \tilde{P}_g \cap G$ is a parabolic subgroup of G , we must find a cocharacter χ that satisfies (6) and is defined over \mathbb{F} . Then $P_g = P(\chi)$ will be pseudo-parabolic and hence parabolic.

Let Γ be the group of field automorphisms of $\tilde{\mathbb{F}}$ over \mathbb{F} . Since $g \in \mathcal{G}(\mathbb{F})$ and Γ acts continuously, the subgroup \tilde{P}_g is Γ -invariant by (3), so that $\gamma \circ \tilde{\chi} \circ \gamma^{-1}$ satisfies (6) for all $\gamma \in \Gamma$. Since the set of solutions of (6) forms a cone in the free abelian group $X_*(\tilde{\mathcal{T}})$, it contains

$$\tilde{\chi}^\Gamma : \lambda \mapsto \prod_{\gamma \in \Gamma} \gamma(\tilde{\chi}(\gamma^{-1} \lambda)).$$

Thus $\tilde{P}_g = \tilde{P}(\tilde{\chi}^\Gamma)$. The cocharacter $\tilde{\chi}^\Gamma$ is defined over $\tilde{\mathbb{F}}^\Gamma$. The field extension $\mathbb{F} \subseteq \tilde{\mathbb{F}}^\Gamma$ is finite and purely inseparable, see for example [10, Section 7.7]. Hence some positive multiple χ of $\tilde{\chi}^\Gamma$ is defined over \mathbb{F} and still satisfies (6). This yields $\tilde{P}_g = \tilde{P}(\chi)$ and finishes the proof of (a).

Now we prove (b). $\text{Lie}_{\tilde{\mathbb{F}}}(\mathcal{P}_g)$ is spanned by the vectors $X \in \text{Lie}_{\tilde{\mathbb{F}}}(\mathcal{G})$ with $\text{Ad}(g_s)X = \lambda X$ with $\tilde{v}(\lambda) \geq 0$. Similarly, $\text{Lie}_{\tilde{\mathbb{F}}}(\mathcal{R}_u(\mathcal{P}_g))$ is spanned by the root subspaces $\text{Lie}_{\tilde{\mathbb{F}}}(\mathcal{U}_\alpha)$ with $\alpha \in \Phi(\mathcal{P}_g, \mathcal{T})$ but $-\alpha \notin \Phi(\mathcal{P}_g, \mathcal{T})$. These are precisely the $\alpha \in \Phi$ with $\tilde{v}(\alpha(g_s)) > 0$. Therefore

$$\lim_{n \rightarrow \infty} g_s^n h g_s^{-n} = 1 \iff h \in \text{R}_u(\tilde{P}_g).$$

Since all powers of g_u are contained in the bounded subgroup \tilde{K} , these statements are also equivalent to $\lim_{n \rightarrow \infty} g^n h g^{-n} = 1$. Now (b) follows because $\text{R}_u(P_g) = \text{R}_u(\tilde{P}_g) \cap P_g$.

Next we establish (c). Let χ be a cocharacter of \mathcal{G} defined over \mathbb{F} with $P_g = P(\chi)$. The same reasoning as in the proof of (a) shows that $P_{g^{-1}} = P(-\chi)$. The assertion (c) now follows by applying [20, Theorem 13.4.2] to P_g and $P_{g^{-1}}$.

Finally, we turn to (d). The eigenvalues of $\text{Ad}(g_s)$ acting on $\text{Lie}_{\tilde{\mathbb{F}}}(\mathcal{M}_g)$ all have valuation 0. Hence $\text{Ad}(g)$ lies in a bounded subgroup of the adjoint group of \tilde{M}_g . Equivalently, the image of g in $\tilde{M}_g / \text{Z}(\tilde{M}_g)$ generates a bounded subgroup. Finally, we note that $M_g / \text{Z}(M_g)$ can be identified with a subgroup of $\tilde{M}_g / \text{Z}(\tilde{M}_g)$. \square

3. SOME BRUHAT–TITS THEORY

We keep the notation from Section 2. Let \mathbb{F} be a non-Archimedean local field with a discrete valuation v . We normalise v by $v(\mathbb{F}^\times) = \mathbb{Z}$. Let $\mathcal{O} \subset \mathbb{F}$ be the ring of integers and $\mathfrak{P} \subset \mathcal{O}$ its maximal ideal. The cardinality q of the residue field \mathcal{O}/\mathfrak{P} is a power of a prime number p . We briefly call \mathbb{F} a p -adic field.

Bruhat and Tits [3, 4, 21] constructed an affine building for any reductive p -adic group $G = \mathcal{G}(\mathbb{F})$. More precisely, they constructed two buildings, one corresponding to G and one corresponding to the maximal semisimple quotient of G . We call the latter the Bruhat–Tits building of G and denote it by $\mathcal{B}(\mathcal{G}, \mathbb{F})$. Relying on [18, § 1.1] and [23, Section 1], we now recall its construction. The main ingredients are certain subgroups $U_{\alpha,r}$ and H_r of G .

3.1. The prolonged valued root datum. Let $\langle \cdot, \cdot \rangle: X_*(\mathcal{S}) \times X^*(\mathcal{S}) \rightarrow \mathbb{Z}$ be the canonical pairing. There is a unique group homomorphism

$$\nu: \text{Z}_G(S) \rightarrow X_*(\mathcal{S}) \otimes_{\mathbb{Z}} \mathbb{R}$$

such that $\langle \nu(z), \chi|_S \rangle = -v(\chi(z))$ for all $\chi \in X^*(\text{Z}_G(S))$. Let

$$H := \ker(\nu) = \{z \in \text{Z}_G(S) : v(\chi(z)) = 0 \text{ for all } \chi \in X^*(\text{Z}_G(S))\}.$$

be the maximal compact subgroup of $\text{Z}_G(S)$.

Bruhat and Tits [4] defined discrete decreasing filtrations of H and U_α by compact open subgroups H_r and $U_{\alpha,r}$, respectively. These groups satisfy the properties of a “prolonged valued root datum” [3, § 6.2]. We first describe these subgroups in the special case where \mathcal{G} splits over \mathbb{F} . Then each U_α is a one-dimensional vector space over \mathbb{F} , and a Chevalley basis of $\text{Lie}_{\mathbb{F}}(\mathcal{G})$ gives rise to an isomorphism $U_\alpha \cong \mathbb{F}$. Chevalley bases are known to exist but they are not unique. We fix one, and we use suitable subsets as bases of $\text{Lie}_{\mathbb{F}}(\mathcal{P}_D)$ and $\text{Lie}_{\mathbb{F}}(\mathcal{M}_D)$, for any standard parabolic subgroup P_D with Levi factor M_D . Thus U_α is endowed with a discrete valuation v_α and one defines

$$(7) \quad U_{\alpha,r} := v_\alpha^{-1}([r, \infty)) \quad \text{for } r \in \mathbb{R}.$$

By assumption, the maximal split torus is a maximal torus, that is, $\mathcal{S} = \mathcal{Z}_{\mathcal{G}}(\mathcal{S})$. For $r < 0$ we may put $H_r = H$, but H_0 is more difficult to define. According to [4, 5.2.1] there is a canonical smooth affine \mathcal{O} -group scheme \mathfrak{Z} such that $\mathfrak{Z}(\mathbb{F}) = \text{Z}_G(S)$. Let \mathfrak{Z}_c be the neutral component of \mathfrak{Z} and put $H_0 := \mathfrak{Z}_c(\mathcal{O})$. The inclusions

$$H_0 \subseteq \mathfrak{Z}(\mathcal{O}) \subseteq H$$

are all of finite index. We define

$$(8) \quad H_r := \{z \in H_0 : v(\chi(z) - 1) \geq r \text{ for all } \chi \in X^*(\mathcal{Z}_{\mathcal{G}}(\mathcal{S}))\}$$

for $r > 0$ as in [18, Proposition I.2.6].

Now we extend the above construction to a non-split group G . Proposition 2.1 provides a finite Galois extension $\tilde{\mathbb{F}}$ of \mathbb{F} over which \mathcal{G} splits. The strategy of descent is explained in [3, Chapitre 9]; the basic idea is to construct the required groups first in $\mathcal{G}(\tilde{\mathbb{F}})$ and then to intersect them with $\mathcal{G}(\mathbb{F})$. This does not work as such because the root system of $\mathcal{G}(\tilde{\mathbb{F}})$ is usually larger than that of $\mathcal{G}(\mathbb{F})$, so that must be

taken into account as well. Bruhat and Tits descend in two steps: first from split to quasi-split, then from there to the general case. This is, in all probability, necessary for the proof, but the conclusions can be written down in one step. Of course it is by no means obvious that the groups we will define below form a (prolonged) valued root datum: proving this is precisely what most of the work in [4] is dedicated to.

If X is any object constructed over \mathbb{F} , then we will denote the corresponding object over $\tilde{\mathbb{F}}$ by \tilde{X} . According to [19, Proposition I.2.3] $\tilde{\mathbb{F}}$ is also a local field, and there is a unique discrete valuation $\tilde{v}: \tilde{\mathbb{F}} \rightarrow \mathbb{Q} \cup \{\infty\}$ that extends v . By definition,

$$\tilde{v}(\tilde{\mathbb{F}}^\times) = e_{\tilde{\mathbb{F}}/\mathbb{F}}^{-1} \mathbb{Z},$$

where $e_{\tilde{\mathbb{F}}/\mathbb{F}} \in \mathbb{N}$ is the ramification index of $\tilde{\mathbb{F}}$ over \mathbb{F} . The constructions above still work for this non-normalised valuation \tilde{v} .

Let $\tilde{S} \subseteq \mathcal{G}(\tilde{\mathbb{F}})$ be a maximal $\tilde{\mathbb{F}}$ -split torus that contains $\mathcal{S}(\tilde{\mathbb{F}})$. Since $\tilde{S} \supseteq S$, restriction of characters defines a surjection

$$(9) \quad \rho_S: \tilde{\Phi} \cup \{0\} \rightarrow \Phi \cup \{0\}.$$

For $\alpha \in \Phi^{\text{red}}$ and $r \in \mathbb{R}$ the descent [4, 4.2.2 and 5.1.16] boils down to

$$(10) \quad \begin{aligned} U_{\alpha,r} &:= U_\alpha \cap \left(\prod_{\beta \in \rho_S^{-1}\{\alpha\}} \tilde{U}_{\beta,r} \times \prod_{\beta \in \rho_S^{-1}\{2\alpha\}} \tilde{U}_{\beta,2r} \right), \\ U_{2\alpha,r} &:= U_{2\alpha} \cap U_{\alpha,r/2}. \end{aligned}$$

These groups do not depend on the chosen ordering of the factors. For a standard Levi subgroup $M_D \subseteq G$ and $\alpha \in \Phi_D$, our consistent choice of Chevalley bases ensures that it does not matter whether we consider the groups $U_{\alpha,r}$ in G or M_D .

We can use (10) to define a valuation on U_α by

$$(11) \quad v_\alpha(u_\alpha) := \sup \{r \in \mathbb{R} : u_\alpha \in U_{\alpha,r}\}.$$

Clearly this reproduces (7) in the split case. Let Γ_α be the set of $r \in \mathbb{R}$ at which $U_{\alpha,r}$ jumps, or equivalently the set of values of v_α (except $v_\alpha(1) = \infty$). By construction, $\tilde{\Gamma}_\beta = e_{\tilde{\mathbb{F}}/\mathbb{F}}^{-1} \mathbb{Z}$ for all $\beta \in \tilde{\Phi}$, which implies

$$\mathbb{Z} \subseteq \Gamma_\alpha \subseteq e_{\tilde{\mathbb{F}}/\mathbb{F}}^{-1} \mathbb{Z} \quad \text{for all } \alpha \in \Phi.$$

More precisely, [3, 6.2.23] and [18, Lemma I.2.10] yield $n_\alpha \in \mathbb{N}$ for $\alpha \in \Phi$ with the following properties:

- $\Gamma_\alpha = n_\alpha^{-1} \mathbb{Z}$;
- $n_{w\alpha} = n_\alpha$ for $w \in W(\Phi)$;
- $n_{2\alpha} = n_\alpha$ or $n_{2\alpha} = n_\alpha/2$ whenever $\alpha, 2\alpha \in \Phi$.

Similar to (10) one defines for $r \in \mathbb{R}$ (see [18, I.2.6] and [23, Section 1]):

$$(12) \quad H_r := Z_G(S) \cap \left(\tilde{H}_r \times \prod_{\beta \in \rho_S^{-1}\{0\}} \tilde{U}_{\beta,r} \right).$$

A particularly useful property of the above groups, which holds more or less by the definition of a prolonged valued root datum [3, Proposition 6.4.41], is as follows. Let $\alpha, \beta \in \Phi \cup \{0\}$ and let $r, s \in \mathbb{R}$, with $r \geq 0$ if $\alpha = 0$ and $s \geq 0$ if $\beta = 0$. Then

$$(13) \quad [U_{\alpha,r}, U_{\beta,s}] \subseteq \text{subgroup generated by } \bigcup_{n,m \in \mathbb{Z}_{>0}} U_{n\alpha+m\beta, nr+ms},$$

where $U_{0,t} = H_t$ and $U_{\delta,t} = \{1\}$ if $\delta \notin \Phi \cup \{0\}$. We will need an iterated version of this, which must have been known already to Bruhat and Tits, but for which the authors did not find a reference.

Lemma 3.1. *Let $\alpha_i \in \Phi^+ \cup \{0\}$, $r_i \in \mathbb{R}$ and $u_i \in U_{\alpha_i, r_i}$ for $i = 1, 2, \dots, n$. Assume that $r_i \geq 0$ whenever $\alpha_i = 0$. Then*

$$[u_1, [u_2, [\dots [u_{n-1}, u_n] \dots]]]$$

lies in the group generated by the $U_{\sum_{i=1}^n k_i \alpha_i, \sum_{i=1}^n k_i r_i}$, where the k_i run over $\mathbb{Z}_{>0}$.

Proof. Let us call the group in question K . Suppose that $y_j \in U_{\sum_{i=2}^n k_i \alpha_i, \sum_{i=2}^n k_i r_i}$ for some $k_i \in \mathbb{Z}_{>0}$ (depending on j). Notice that $\sum_{i=2}^n k_i \alpha_i$ cannot be a negative root, and that $\sum_{i=2}^n k_i r_i \geq 0$ if $\sum_{i=2}^n k_i \alpha_i = 0$. We will show by induction on $l \in \mathbb{N}$ that

$$[u_1, y_1 \dots y_l] \text{ is an element of } K.$$

For $l = 1$ this is (13). For $l \geq 2$ we can rewrite it as

$$[u_1, y_1 \dots y_l] = u_1 y_1 u_1^{-1} [u_1, y_2 \dots y_l] y_1^{-1} = [u_1, y_1] y_1 [u_1, y_2 \dots y_l] y_1^{-1}.$$

By the induction hypothesis all terms on the right are in K .

For the actual lemma we use another induction, with respect to n . The case $n = 1$ is trivial. For $n > 1$, the induction hypothesis provides y_j as above, such that

$$[u_1, [u_2, [\dots [u_{n-1}, u_n] \dots]]] = [u_1, y_1 \dots y_l],$$

which by the above lies in K . \square

3.2. The affine Bruhat–Tits building. The image of any cocharacter $\mathbb{F}^\times \rightarrow Z_c(G)$ lies in $S_\Delta \subseteq S$, the maximal \mathbb{F} -split torus in $Z_c(G)$. Hence $X_*(Z_c(G)) = X_*(S_\Delta)$. The standard *apartment* is

$$A_S := (X_*(S)/X_*(Z_c(G))) \otimes_{\mathbb{Z}} \mathbb{R} = (X_*(\mathcal{S})/X_*(\mathcal{S}_\Delta)) \otimes_{\mathbb{Z}} \mathbb{R}.$$

The affine Bruhat–Tits building $\mathcal{B}(\mathcal{G}, \mathbb{F})$ will be defined as $G \times A_S / \sim$ for a suitable equivalence relation \sim .

Let $\langle \cdot, \cdot \rangle_{A_S}$ be a $W(\Phi)$ -invariant inner product on A_S . Then the different irreducible components Φ_i^\vee of Φ^\vee are orthogonal and on $\mathbb{R}\Phi_i^\vee$ the inner product is unique up to scaling. Thus we may assume that $\langle \alpha^\vee, \alpha^\vee \rangle_{A_S} = 1$ for all short coroots $\alpha^\vee \in \Phi^\vee$.

The centraliser $Z_G(S)$ acts on A_S by

$$g \cdot x = x + \nu(g).$$

This extends to an action of $N_G(S)$ on A_S by affine automorphisms, such that the linear part of $x \mapsto g \cdot x$ is given by the image of $g \in N_G(S)$ in $W(\Phi)$. In particular, the action of g on A_S is a translation if and only if $g \in Z_G(S)$. The affine hyperplanes

$$(14) \quad A_{S,\alpha,k} := \{x \in A_S : \langle x, \alpha \rangle = k\} \quad \text{for } \alpha \in \Phi \text{ and } k \in \Gamma_\alpha$$

turn A_S into a polysimplicial complex. The open polysimplices are called facets, that is, a *facet* in A_S is a non-empty subset $F \subseteq A_S$ such that

- $F \subseteq A_{S,\alpha,k}$ or F lies entirely on one side of $A_{S,\alpha,k}$ for all $\alpha \in \Phi$ and $k \in \Gamma_\alpha$;
- F cannot be extended to a larger set with the first property.

Thus the closure of a facet is a polysimplex, and a facet is closed if and only if it is a single point. Moreover, a facet is open in A_S if and only if it is of maximal dimension, in which case we call it a *chamber*.

The affine action of $N_G(S)$ on A_S respects the polysimplicial structure. In fact, $N_G(S)$ is generated by the translations coming from $Z_G(S)$ and the reflections in the hyperplanes $A_{S,\alpha,k}$:

$$x \mapsto x + (k - \langle x, \alpha \rangle) \alpha^\vee \quad \alpha \in \Phi, k \in \Gamma_\alpha,$$

where $\alpha^\vee \in \Phi^\vee$ is the coroot corresponding to α .

For a non-empty subset $\Omega \subseteq A_S$ we define

$$(15) \quad f_\Omega: \Phi \rightarrow \mathbb{R} \cup \{\infty\}, \quad f_\Omega(\alpha) := -\inf_{x \in \Omega} \langle x, \alpha \rangle = \sup_{x \in \Omega} \langle x, -\alpha \rangle.$$

This gives rise to the following subgroups of G :

$$(16) \quad \begin{aligned} U_\Omega &:= \text{subgroup generated by } \bigcup_{\alpha \in \Phi^{\text{red}}} U_{\alpha, f_\Omega(\alpha)}, \\ N_\Omega &:= \{n \in N_G(S) : n \cdot x = x \text{ for all } x \in \Omega\}, \\ P_\Omega &:= N_\Omega U_\Omega = U_\Omega N_\Omega. \end{aligned}$$

The latter is a group because $nU_\Omega n^{-1} = U_{n\Omega}$ for all $n \in N_G(S)$. For $\Omega = \{x\}$ we abbreviate $U_\Omega = U_x$, which should not be confused with the root subgroups U_α .

Given a partition $\Phi = \Phi^+ \cup \Phi^-$ of $\Phi(\mathcal{G}, \mathbb{F})$ in positive and negative roots, we let U^\pm be the subgroup of G generated by $\bigcup_{\alpha \in \Phi^\pm} U_\alpha$. We write

$$U_\Omega^+ := U_\Omega \cap U^+ \quad \text{and} \quad U_\Omega^- := U_\Omega \cap U^-.$$

Proposition 3.2 ([3, 6.4.9]). *These subgroups have the following properties:*

- (a) $U_\Omega \cap U_\alpha = U_{\alpha, f_\Omega(\alpha)}$ for all $\alpha \in \Phi$.
- (b) The product map

$$\prod_{\alpha \in \Phi^{\text{red}} \cap \Phi^\pm} U_{\alpha, f_\Omega(\alpha)} \rightarrow U_\Omega^\pm$$

is an isomorphism of algebraic varieties, for any ordering of the factors.

- (c) $U_\Omega = U_\Omega^+ U_\Omega^- (U_\Omega \cap N_G(S))$.

We define an equivalence relation \sim on $G \times A_S$ by

$$(g, x) \sim (h, y) \iff \text{there is } n \in N_G(S) \text{ with } nx = y \text{ and } g^{-1}hn \in U_x.$$

As announced, the Bruhat–Tits building of G is

$$\mathcal{B}(\mathcal{G}, \mathbb{F}) = G \times A_S / \sim.$$

The group G acts naturally on $\mathcal{B}(\mathcal{G}, \mathbb{F})$ from the left, and the map

$$A_S \rightarrow \mathcal{B}(\mathcal{G}, \mathbb{F}), \quad x \mapsto (1, x) / \sim$$

is an $N_G(S)$ -equivariant embedding. An apartment of $\mathcal{B}(\mathcal{G}, \mathbb{F})$ is a subset of the form $g \cdot A_S$ with $g \in G$, and $g \cdot A_S = A_S$ if and only if $g \in N_G(S)$. Since all maximal split tori of G are conjugate by [2, Théorème 4.21], there is a bijection between apartments in $\mathcal{B}(\mathcal{G}, \mathbb{F})$ and maximal split tori in G .

A facet of $\mathcal{B}(\mathcal{G}, \mathbb{F})$ is a subset of the form $g \cdot F$, where $g \in G$ and F is a facet of A_S . For a polysimplicial complex Σ , we denote the set of vertices by Σ° and the set of n -dimensional polysimplices in Σ by Σ^n for $n \in \mathbb{N}$.

For any subset $\Omega \subseteq \mathcal{B}(\mathcal{G}, \mathbb{F})$, we denote the pointwise stabiliser of Ω by P_Ω . This is consistent with (16) when $\Omega \subseteq A_S$.

4. FIXED POINTS IN THE BUILDING

An element g of G is called *compact* if its image in $G/Z(G)$ belongs to a compact subgroup of $G/Z(G)$. According to the Bruhat–Tits Fixed Point Theorem (see [3, § 3.2]), the compact elements of G are precisely those that fix a point in the building $\mathcal{B}(\mathcal{G}, \mathbb{F})$. In this section, we study how the fixed point subset $\mathcal{B}(\mathcal{G}, \mathbb{F})^\gamma$ depends on γ .

Let H be a group of polysimplicial automorphisms of $\mathcal{B}(\mathcal{G}, \mathbb{F})$. If $x, y \in \mathcal{B}(\mathcal{G}, \mathbb{F})^H$, then H fixes the geodesic segment $[x, y]$ pointwise by [3, 2.5.4]. Consequently, $\mathcal{B}(\mathcal{G}, \mathbb{F})^H$ is a convex subset of $\mathcal{B}(\mathcal{G}, \mathbb{F})$. Recall that a *chamber complex* is a polysimplicial complex Σ such that:

- all maximal polysimplices of Σ (the chambers) have the same dimension;

- given any two chambers C_1 and C_2 of Σ , there exists a gallery of chambers connecting C_1 and C_2 .

If $g \in G$ is compact and belongs to a maximal split torus S of G , then there is a chamber in the corresponding apartment A_S that is fixed pointwise by g . There exist, however, regular semisimple elements $\gamma \in G$ that fix no chamber in the building pointwise. For such elements the fixed point subcomplex is not necessarily a chamber complex. But once g fixes a chamber, say, because it belongs to a maximal split torus, the fixed point subset is automatically a chamber complex:

Lemma 4.1. *Suppose that H fixes a chamber $C \subseteq \mathcal{B}(\mathcal{G}, \mathbb{F})$ pointwise. Then $\mathcal{B}(\mathcal{G}, \mathbb{F})^H$ is a chamber complex.*

Proof. This is well-known, but we include a proof anyway. Let $x \in \mathcal{B}(\mathcal{G}, \mathbb{F})^H$ and let A_x be an apartment that contains C and x . Since $\dim C = \dim A_x$ and $\mathcal{B}(\mathcal{G}, \mathbb{F})^H$ is convex, it contains an open subset of some chamber $C_x \subseteq A_x$ with $x \in \overline{C_x}$. Thus H fixes C_x pointwise and $\mathcal{B}(\mathcal{G}, \mathbb{F})^H$ is the union of all its closed chambers.

Suppose that \mathcal{C} is any collection of chambers of an apartment A_S of $\mathcal{B}(\mathcal{G}, \mathbb{F})$. Then $\bigcup_{C \in \mathcal{C}} \overline{C}$ is convex if and only if all minimal galleries between elements of \mathcal{C} are contained in \mathcal{C} . Hence $\mathcal{B}(\mathcal{G}, \mathbb{F})^H \cap A_S$ contains all minimal galleries between its chambers. \square

4.1. The split case. Let $S \subset G$ be a split maximal torus and let $\gamma \in S$ be a compact element. Then $v(\chi(\gamma)) = 0$ for all $\chi \in X^*(S)$, so that γ fixes the apartment A_S pointwise. The subcomplex $\mathcal{B}(\mathcal{G}, \mathbb{F})^\gamma \subseteq \mathcal{B}(\mathcal{G}, \mathbb{F})$ is convex and S -invariant. Its core is formed by the apartment A_S and from there “hairs” extend in all directions. This terminology applies quite well to one-dimensional buildings, but in general such a hair is a (not necessarily bounded) chamber complex. Since S acts by translations on A_S , it shifts all these hairs. If $\gamma \in S$ is regular, then $\mathcal{B}(\mathcal{G}, \mathbb{F})^\gamma/S$ is compact by [9, Section 9.1]: the length of the hairs is finite.

Now we study when an arbitrary point $x \in \mathcal{B}(\mathcal{G}, \mathbb{F})$ is fixed by $\gamma \in S$. Choose a chamber $C_0 \subseteq A_S$ and let ρ be the retraction of $\mathcal{B}(\mathcal{G}, \mathbb{F})$ to A_S centred at C_0 . Let Φ^+ be a system of positive roots in Φ such that $f_{\rho(x)}(\alpha) \geq f_{C_0}(\alpha)$ for all $\alpha \in \Phi^+$; equivalently, Φ^+ contains all roots with $f_{\rho(x)}(\alpha) > f_{C_0}(\alpha)$. Let Δ be the basis of Φ corresponding to Φ^+ .

Then $U_{C_0} \cap U_\alpha \subseteq U_{\rho(x)} \cap U_\alpha$ for all $\alpha \in \Phi_-$, so $U_{C_0}^- \subseteq U_{\rho(x)}^-$. Furthermore, $N_{C_0} = N_{\rho(x)}$, which together with Proposition 3.2.(c) shows that $P_{C_0} \subseteq U_{C_0}^+ P_{\rho(x)}$. Since P_{C_0} acts transitively on the set of apartments containing C_0 by [3, 7.4.9], there is $u \in U_{C_0}^+$ with $x = u\rho(x)$. Thus we want to know which part of the apartment uA_S is fixed by γ .

By definition, $u \in U_{C_0}^+$ fixes all $y \in A_S$ satisfying $-\alpha(y) \leq f_{C_0}(\alpha)$ for all $\alpha \in \Phi^+$. These points constitute a cone in $A_S \cap uA_S$, which is fixed by γ . We are interested in the larger subset $(uA_S)^\gamma$, which is a convex subcomplex of $\mathcal{B}(\mathcal{G}, \mathbb{F})^\gamma$. Hence the complex $Y := u^{-1}(uA_S)^\gamma$ is convex as well. Concretely, this means that $Y \subseteq A_S$ is determined by a system of equations $-\alpha(y) \leq r_\alpha$ for certain $r_\alpha \in \mathbb{R}$, $\alpha \in \Phi^+$. We need some notation to make this more explicit. The *singular depth* of γ in the direction $\alpha \in \Phi$ is

$$\text{sd}_\alpha(\gamma) := v(\alpha(\gamma) - 1).$$

We also let $\text{sd}(\gamma) := \max_{\alpha \in \Phi^+} \text{sd}_\alpha(\gamma)$.

Recall that the height of a positive root is defined as follows:

- $\text{ht}(\alpha) = 1$ if $\alpha \in \Phi^+$ is simple;
- $\text{ht}(\alpha + \beta) = \text{ht}(\alpha) + \text{ht}(\beta)$ if $\alpha, \beta, \alpha + \beta \in \Phi^+$.

Since ht extends to a group homomorphism $X^*(G/\mathbb{Z}_c(G)) \rightarrow \mathbb{R}$, we may regard it as a point in the apartment A_S . Since y is contained in the same apartment,

this gives meaning to the linear combination $y + \text{sd}(\gamma) \text{ht}$ for $y \in Y$ appearing in Proposition 4.2.(c) below.

By Proposition 3.2.(b) we can write

$$(17) \quad u = \prod_{\alpha \in \Phi^+} u_\alpha \quad \text{with} \quad u_\alpha \in U_{\alpha, f_{C_0}(\alpha)}.$$

Proposition 4.2. *Let $y \in u^{-1}(uA_S)^\gamma$.*

- (a) *The compact element $\gamma \in S$ fixes $x = u\rho(x)$ if and only if $[\gamma, u^{-1}] \in U_{\rho(x)}^+$.*
- (b) *$u_\alpha \in U_{\alpha, -\alpha(y) - \text{sd}_\alpha(\gamma)}$ for all simple roots $\alpha \in \Delta$.*
- (c) *$u \in U_{y + \text{sd}(\gamma) \text{ht}}^+$, where $\text{sd}(\gamma) \text{ht} \in A_S$.*

Proof. (a) Since $\gamma \in S$ fixes $\rho(x) \in A_S$,

$$\gamma(x) = \gamma u \rho(x) = \gamma u \gamma^{-1} \rho(x).$$

This point equals $x = u\rho(x)$ if and only if $\gamma u^{-1} \gamma^{-1} u \rho(x) = \rho(x)$, which is equivalent to $[\gamma, u^{-1}] \in P_{\rho(x)}$. As $u \in U^+$ and γ normalises U^+ , this is equivalent to $[\gamma, u^{-1}] \in P_{\rho(x)} \cap U^+ = U_{\rho(x)}^+$.

(b) The decomposition (17) is unique once we fix an ordering on Φ^+ , but the terms u_α may depend on this ordering. Let $\Phi^* := \Phi^+ \setminus \Delta$ be the set of non-simple positive roots. Then $\bigcup_{\alpha \in \Phi^*} (U_\alpha \cap U_{C_0})$ generates a normal subgroup $U_{C_0}^*$ of $U_{C_0}^+$. The quotient $U_{C_0}^+ / U_{C_0}^*$ is abelian and can be identified with a lattice in the \mathbb{F} -vector space $\prod_{\alpha \in \Delta} U_\alpha$. The image of u in $U_{C_0}^+ / U_{C_0}^*$ is $\prod_{\alpha \in \Delta} u_\alpha$, which shows that the ingredients u_α of (17) for $\alpha \in \Delta$ are independent of the ordering of Φ^+ .

Suppose now that γ fixes $uy \in uA_S$. By part (a), we have $[\gamma, u^{-1}] \in U_y^+$. Since γ normalises the groups $U_{\alpha, r}$ for $\alpha \in \Phi^+$, $r \in \mathbb{R}$, this implies

$$(18) \quad [\gamma, u^{-1}]U_y^* = \prod_{\alpha \in \Delta} [\gamma, u_\alpha^{-1}]U_y^* \in U_y^+ / U_y^*.$$

But on the vector space U_α the map $a \mapsto [\gamma, a]$ can be identified with multiplication by $\alpha(\gamma) - 1$. Hence (18) is equivalent to

$$(19) \quad u_\alpha \in (\alpha(\gamma) - 1)^{-1} U_{\alpha, -\alpha(y)}$$

for all $\alpha \in \Delta$. Together with (7) implies the statement (b).

(c) We fix an ordering $\Phi^+ = \{\alpha_1, \alpha_2, \dots, \alpha_k\}$ with $\text{ht}(\alpha_i) \leq \text{ht}(\alpha_{i+1})$ for all i , and we get a unique decomposition $u = \prod_{i=1}^k u_{\alpha_i}$ in $U_{C_0}^+$. Similarly, Proposition 3.2.(b) yields a unique decomposition

$$(20) \quad \prod_{i=1}^k [\gamma, u^{-1}]_{\alpha_i} = [\gamma, u^{-1}] = \gamma u_{\alpha_k}^{-1} u_{\alpha_{k-1}}^{-1} \cdots u_{\alpha_1}^{-1} \gamma^{-1} u_{\alpha_1} u_{\alpha_2} \cdots u_{\alpha_k}.$$

By construction $[\gamma, u^{-1}]_\alpha \in U_{\alpha, f_{C_0}(\alpha)}$, and γ fixes uy if and only if, even more,

$$(21) \quad [\gamma, u^{-1}]_\alpha \in U_{\alpha, -\alpha(y)} \quad \text{for all } \alpha \in \Phi^+.$$

Assuming (21), we will show by induction on $\text{ht}(\alpha)$ that

$$(22) \quad [\gamma, u_\alpha^{-1}] \in U_{\alpha, -\alpha(y) + (1 - \text{ht}(\alpha))\text{sd}(\gamma)} \quad \text{for all } \alpha \in \Phi^+.$$

Like in (19), this statement is equivalent to $u_\alpha \in U_{\alpha, -\alpha(y) - \text{sd}_\alpha(\gamma) + (1 - \text{ht}(\alpha))\text{sd}(\gamma)}$, which for roots α of height 1 is part (b).

Let us assume (22) for roots of height less than k . Let $N_{>k}$ be the product of the groups U_α for roots α of height greater than k . This is a normal subgroup of the Borel group SU^+ , and the subgroups $U_\alpha \subseteq U^+$ for a root α of height k become central in the quotient $U^+ / N_{>k}$. We may determine the α -component for a root α of height k by computations in $U^+ / N_{>k}$ because of the uniqueness of the decomposition (17).

Now we split u up as $u_{<k}u_ku_{>k}$, where the factors $u_{<k}$, u_k and $u_{>k}$ contain the contributions u_α of positive roots α with height less than k , equal to k , and greater than k , respectively. In the quotient $U^+/N_{>k}$, we may drop $u_{>k}$, and u_k becomes central. Hence

$$(23) \quad \begin{aligned} [\gamma, u^{-1}] &= \gamma u_{>k}^{-1} u_k^{-1} u_{<k}^{-1} \gamma^{-1} u_{<k} u_k u_{>k} \equiv \gamma u_k^{-1} \gamma^{-1} \gamma u_{<k}^{-1} \gamma^{-1} u_{<k} u_k \\ &\equiv \gamma u_k^{-1} \gamma^{-1} u_k \gamma u_{<k}^{-1} \gamma^{-1} u_{<k} = [\gamma, u_k^{-1}] [\gamma, u_{<k}^{-1}] \equiv \left(\prod_{\text{ht}(\alpha)=k} [\gamma, u_\alpha^{-1}] \right) [\gamma, u_{<k}^{-1}], \end{aligned}$$

where we compute in the quotient $U^+/N_{>k}$. We will use the induction hypothesis and the estimate on $[\gamma, u_\alpha]$ to estimate $[\gamma, u_\alpha^{-1}]$ when $\text{ht}(\alpha) = k$.

We first rewrite a commutator $[\gamma, z_1 \cdots z_j]$ as a product of iterated commutators

$$(24) \quad C(z_{i_1}, \dots, z_{i_k}) := [z_{i_1}, [z_{i_2}, \dots, [z_{i_{k-1}}, [\gamma, z_{i_k}]] \dots]].$$

We claim that $[\gamma, z_1 \cdots z_j]$ is a product of the factors $C(z_{i_1}, \dots, z_{i_k})$ with $1 \leq i_1 < i_2 < \dots < i_k \leq j$, each factor appearing exactly once. The proof is by induction on j , the case $j = 1$ being clear. For the induction step, we use

$$\begin{aligned} [\gamma, z_1 \cdots z_j] &= \gamma z_1 \gamma^{-1} [\gamma, z_2 \cdots z_j] z_1^{-1}, \\ \gamma z_1 \gamma^{-1} x_1 \cdots x_k z_1^{-1} &= [\gamma, z_1] \cdot [z_1, x_1] x_1 \cdot [z_1, x_2] x_2 \cdots [z_1, x_k] x_k. \end{aligned}$$

By the induction hypothesis, $[\gamma, z_2 \cdots z_j]$ is the product in some order of the factors $C(z_{i_1}, \dots, z_{i_k})$ for all $2 \leq i_1 < \dots < i_k \leq j$. Plugging this into the second equation above shows that $[\gamma, z_1 \cdots z_j]$ is the product in some order of the factors $C(z_{i_1}, \dots, z_{i_k})$ for all $1 \leq i_1 < \dots < i_k \leq j$. By the way, a more careful induction argument also yields the order of the factors: it is the reverse lexicographic order for the words $(j - i_k, i_{k-1}, i_{k-2}, \dots, i_1)$.

Now we apply this to $u_{<k}^{-1} = u_{\alpha_1}^{-1} \cdots u_{\alpha_1}^{-1} = z_1 \cdots z_l$. By the induction hypothesis and by Lemma 3.1, all the occurring $C(u_{\alpha_{i_1}}^{-1}, \dots, u_{\alpha_{i_k}}^{-1})$ lie in the group generated by the $U_{\alpha, r}$ with $\alpha = \sum_{j=1}^k k_j \alpha_{i_j}$ and $r = \text{sd}_{\alpha_{i_k}}(\gamma) + \sum_{j=1}^k k_j r_{i_j}$, where $k_j \in \mathbb{Z}_{>0}$ and

$$r_{i_j} = -\alpha_{i_j}(y) - \text{sd}_{\alpha_{i_j}}(\gamma) + (1 - \text{ht}(\alpha_{i_j}))\text{sd}(\gamma).$$

For such $\alpha \in \Phi^+$ and $r \in \mathbb{R}$ we have

$$(25) \quad \begin{aligned} r &= \text{sd}_{\alpha_{i_k}}(\gamma) + \sum_{j=1}^k k_j (-\alpha_{i_j}(y) - \text{sd}_{\alpha_{i_j}}(\gamma) + (1 - \text{ht}(\alpha_{i_j}))\text{sd}(\gamma)) \geq \\ &\quad -\alpha(y) + (1 - \text{ht}(\alpha))\text{sd}(\gamma) + \left(-1 + \sum_{j=1}^k k_j \right) (\text{sd}(\gamma) - \max_j \text{sd}_{\alpha_{i_j}}(\gamma)) \geq \\ &\quad -\alpha(y) + (1 - \text{ht}(\alpha))\text{sd}(\gamma). \end{aligned}$$

For a root α of height k , (21) and (23) show that $[\gamma, u_\alpha^{-1}]$ must lie in the largest of the groups $U_{\alpha, -\alpha(y)}$ and $U_{\alpha, r}$. Now we see from (25) that in any case $[\gamma, u_\alpha^{-1}] \in U_{\alpha, -\alpha(y) + (1 - \text{ht}(\alpha))\text{sd}(\gamma)}$, so

$$u_\alpha^{-1}, u_\alpha \in U_{\alpha, -\alpha(y) - \text{ht}(\alpha)\text{sd}(\gamma)} = U_\alpha \cap U_{y + \text{sd}(\gamma)\text{ht}}^+. \quad \square$$

Given an arbitrary point $y \in A_S$, the condition in Proposition 4.2.(c) does not imply that γ fixes uy . Counterexamples exist whenever Φ contains an irreducible root system of rank greater than one.

Proposition 4.2 only applies to fixed points of semisimple elements that lie in a split maximal torus. (We will not consider the fixed points of non-semisimple elements of G in this article.) For elements of non-split maximal tori we need yet another aspect of Bruhat–Tits theory.

4.2. The non-split case. The construction of the Bruhat–Tits building over p -adic fields is functorial with respect to finite field extensions by [3, 9.1.17]. For any such extension $\tilde{\mathbb{F}}/\mathbb{F}$, the group

$$\Gamma := \{\sigma \in \text{Aut}(\tilde{\mathbb{F}}) : \sigma|_{\mathbb{F}} = \text{id}_{\mathbb{F}}\}$$

acts naturally on $\mathcal{B}(\mathcal{G}, \tilde{\mathbb{F}})$, and $\mathcal{B}(\mathcal{G}, \mathbb{F})$ is contained in $\mathcal{B}(\mathcal{G}, \tilde{\mathbb{F}})^{\Gamma}$. In particular, for every $g \in \mathcal{G}(\mathbb{F})$ we have an inclusion

$$(26) \quad \mathcal{B}(\mathcal{G}, \mathbb{F})^g = \mathcal{B}(\mathcal{G}, \tilde{\mathbb{F}})^g \cap \mathcal{B}(\mathcal{G}, \mathbb{F}) \subseteq \mathcal{B}(\mathcal{G}, \tilde{\mathbb{F}})^{\Gamma \times \langle g \rangle},$$

where $\langle g \rangle \subseteq \mathcal{G}(\tilde{\mathbb{F}})$ denotes the subgroup generated by g .

In general, $\mathcal{B}(\mathcal{G}, \mathbb{F})$ is strictly smaller than $\mathcal{B}(\mathcal{G}, \tilde{\mathbb{F}})^{\Gamma}$, even if $\tilde{\mathbb{F}}/\mathbb{F}$ is a Galois extension (in which case Γ is its Galois group). Rousseau [17] proved that $\mathcal{B}(\mathcal{G}, \mathbb{F}) = \mathcal{B}(\mathcal{G}, \tilde{\mathbb{F}})^{\Gamma}$ if $\tilde{\mathbb{F}}/\mathbb{F}$ is a tamely ramified Galois extension, see also [16]. Consequently, (26) is an equality for such extensions.

Let $T = \mathcal{T}(\mathbb{F})$ be a maximal torus and $\tilde{\mathbb{F}}/\mathbb{F}$ a finite Galois extension over which T splits, as in Proposition 2.1. Since T is defined over \mathbb{F} , it is Γ -stable, and hence the corresponding apartment $\tilde{A}_{\mathcal{T}(\tilde{\mathbb{F}})}$ of $\mathcal{B}(\mathcal{G}, \tilde{\mathbb{F}})$ is Γ -stable. The action of Γ on $\tilde{A}_{\mathcal{T}(\tilde{\mathbb{F}})}$ is linear, so that the origin of $\tilde{A}_{\mathcal{T}(\tilde{\mathbb{F}})}$ is fixed. Thus Rousseau’s above result implies that

$$(27) \quad \mathcal{B}(\mathcal{G}, \mathbb{F}) \cap \tilde{A}_{\mathcal{T}(\tilde{\mathbb{F}})} \neq \emptyset \quad \text{if } \tilde{\mathbb{F}}/\mathbb{F} \text{ is tamely ramified.}$$

Any $g \in G$ acts on $\text{Lie}_{\mathbb{F}}(\mathcal{G}) / \text{Lie}_{\mathbb{F}}(\mathcal{Z}_{\mathcal{G}}(g))$ by the adjoint representation. The collection $E(g)$ of eigenvalues (in some algebraic closure of \mathbb{F}) is finite and does not contain 1. Assume that G is not a torus and that g is regular, that is, $\mathcal{Z}_{\mathcal{G}}(g)$ has the smallest possible dimension. The number

$$\text{sd}(g) := \max_{\lambda \in E(g)} v(\lambda - 1)$$

is well-defined because every eigenvalue lies in a finite field extension of \mathbb{F} . For irregular $g \in G$ we put $\text{sd}(g) = \infty$, because in that case the multiplicity of the eigenvalue 1 of $\text{Ad}(g) \in \text{End}_{\mathbb{F}}(\text{Lie}_{\mathbb{F}}(\mathcal{G}))$ is too high. Finally, if G is a torus, then we define $\text{sd}(g) = 0$ for all $g \in G$. This definition stems from [1, Section 4], where $\text{sd}(g)$ is called the *singular depth* of γ . We note that

$$(28) \quad \text{sd}(gz) = \text{sd}(g) = \text{sd}(hgh^{-1}) \quad \text{for } z \in Z(G) \text{ and } h \in G.$$

Let T and $\tilde{\mathbb{F}}$ be as above and let $\tilde{\Phi} = \Phi(\mathcal{G}(\tilde{\mathbb{F}}), \mathcal{T}(\tilde{\mathbb{F}}))$ be the corresponding root system. Let \tilde{v} be the discrete valuation that extends v and suppose $\gamma \in T$. Then

$$\text{sd}(\gamma) = \max_{\alpha \in \tilde{\Phi}} \text{sd}_{\alpha}(\gamma),$$

which agrees with the notation from Proposition 4.2.(c). Notice that $\text{sd}(\gamma) \geq 0$, for if $\text{sd}_{\alpha}(\gamma) < 0$ then $\tilde{v}(\alpha(\gamma)) < 0$, so $\tilde{v}(\alpha(\gamma)^{-1}) > 0$ and $\text{sd}_{-\alpha}(\gamma) = 0$.

Now we specialise to a compact regular semisimple element $\gamma \in T$. Then $\mathcal{B}(\mathcal{G}, \mathbb{F})^{\gamma}$ is non-empty by the Bruhat–Tits Fixed Point Theorem. If $T/Z_c(G)$ is anisotropic, then $\mathcal{B}(\mathcal{G}, \mathbb{F})^{\gamma}$ is a finite polysimplicial complex (see [18, p. 53]) and there is an open neighbourhood U of γ in G such that $\mathcal{B}(\mathcal{G}, \mathbb{F})^U = \mathcal{B}(\mathcal{G}, \mathbb{F})^{\gamma}$.

If $T/Z_c(G)$ is not anisotropic, we have a weaker substitute. Since $\mathcal{B}(\mathcal{G}, \mathbb{F})^{\gamma}/T$ is compact, there exists an open neighborhood V of γ in T such that $\mathcal{B}(\mathcal{G}, \mathbb{F})^g = \mathcal{B}(\mathcal{G}, \mathbb{F})^{\gamma}$ for all $g \in V$. Let \tilde{H}_r be as in (12), but with respect to $(\mathcal{G}(\tilde{\mathbb{F}}), \mathcal{T}(\tilde{\mathbb{F}}))$. First the authors believed that one could take $V = \gamma \tilde{H}_r \cap T$ for any $r > \text{sd}(\gamma)$, but this turns out to be incorrect in general. We thank the referee for pointing out the weakness in our former argument.

Lemma 4.3. *Write $\text{ht}(\Phi) := \max_{\alpha \in \Phi^+} \text{ht}(\alpha)$ and let $r > \text{ht}(\Phi)\text{sd}(\gamma)$. Then $\mathcal{B}(\mathcal{G}, \mathbb{F})^{\gamma h} = \mathcal{B}(\mathcal{G}, \mathbb{F})^\gamma$ for all $h \in \tilde{H}_r \cap T$.*

Proof. In view of (26) it suffices to prove the corresponding statement for fixed points in the building $\mathcal{B}(\mathcal{G}, \tilde{\mathbb{F}})$. We use the notation from the proof of Proposition 4.2, but with some additional tildes. We want to know when γ fixes uy , for some point $y \in \tilde{A}_S$. According to (21), this is equivalent to

$$(29) \quad [\gamma, u^{-1}]_\alpha \in \tilde{U}_{\alpha, -\alpha(y)} \quad \text{for all } \alpha \in \tilde{\Phi}.$$

From (23) we know that apart from $[\gamma, u_\alpha^{-1}]$, all the contributions to $[\gamma, u^{-1}]_\alpha$ come from commutators of elements u_β^{-1} with $\text{ht}(\beta) < \text{ht}(\alpha)$. Supposing that u_β has already been fixed for all roots β of smaller height than α , (29) determines which $u_\alpha \in \tilde{U}_\alpha$ can give rise to fixed points uy .

Recall from Section 3.1 that we have a Chevalley basis of $\text{Lie}_{\tilde{\mathbb{F}}}(\mathcal{G})$ and corresponding isomorphisms of algebraic groups $\tilde{U}_\alpha \cong \tilde{\mathbb{F}}$. These restrict to

$$\tilde{U}_{\alpha, r} \cong \{\lambda \in \tilde{\mathbb{F}} : \tilde{v}(\lambda) \geq r\} \quad \text{for all } r \in \mathbb{R},$$

and if u_α corresponds to $\lambda_\alpha \in \tilde{\mathbb{F}}$, then $[\gamma, u_\alpha^{-1}]$ becomes $(1 - \alpha(\gamma))\lambda_\alpha$. Because we are interested in uy , the component u_α is determined only modulo $\tilde{U}_{\alpha, -\alpha(y)}$, that is, λ_α modulo $\{\lambda \in \tilde{\mathbb{F}} : \tilde{v}(\lambda) \geq -\alpha(y)\}$ is all that matters.

Now we compare γ with γh . We note that for all $\beta \in \Phi$

$$(30) \quad \begin{aligned} \tilde{v}((1 - \beta(\gamma)) - (1 - \beta(\gamma h))) &= \tilde{v}(\beta(\gamma)(\beta(h) - 1)) \\ &= \tilde{v}(\beta(h) - 1) = \text{sd}_\beta(h) \geq r > \text{ht}(\Phi)\text{sd}(\gamma). \end{aligned}$$

By (25) the valuation of a contribution from $C(u_{\alpha_{i_1}}^{-1}, \dots, u_{\alpha_{i_k}}^{-1})$ to $[\gamma, u^{-1}]_\alpha$ is at least

$$(31) \quad -\alpha(y) + (1 - \text{ht}(\alpha))\text{sd}(\gamma).$$

Recall that $C(u_{\alpha_{i_1}}^{-1}, \dots, u_{\alpha_{i_k}}^{-1})$ also involves $[\gamma, u_{i_k}^{-1}]$. If we use γh instead of γ , then by (30) and (31) we get a new element whose v_α -value differs only in the fractional ideal of $\tilde{\mathbb{F}}$ where the valuation is at least

$$-\alpha(y) + (1 - \text{ht}(\alpha))\text{sd}(\gamma) + \text{ht}(\Phi)\text{sd}(\gamma) \geq -\alpha(y) + \text{sd}(\gamma).$$

So, if the u_β with $\text{ht}(\beta) < \text{ht}(\alpha)$ have already been fixed, then the condition (29) for both γ and γh leads to two sets of solutions for λ_α , and these sets differ only in the parts of valuation at least

$$-\alpha(y) + \text{sd}(\gamma) - \text{sd}_\alpha(\gamma) \geq -\alpha(y).$$

But these parts do not influence the point uy . Hence γh fixes such a point uy if and only if γ does. Since this holds for all $y \in \tilde{A}_S$ we conclude that

$$\mathcal{B}(\mathcal{G}, \tilde{\mathbb{F}})^{\gamma h} = \mathcal{B}(\mathcal{G}, \tilde{\mathbb{F}})^\gamma. \quad \square$$

5. THE GROUPS $U_\Omega^{(e)}$

Schneider and Stuhler introduced an important system of compact subgroups of G , which they used to derive several interesting results on complex smooth G -representations in [18]. These subgroups were also studied by Moy and Prasad in [12, 13] for their theory of unrefined minimal types, and by Vignéras in [23] in the context of G -representations on vector spaces over general fields.

Let $\tilde{\mathbb{R}}$ be the set $\mathbb{R} \cup \{r+ : r \in \mathbb{R}\} \cup \{\infty\}$ endowed with the ordering

$$r < r+ < s < s+ < \infty \quad \text{if } r < s.$$

We define addition and multiplication with positive numbers on $\tilde{\mathbb{R}}$ in the obvious way, so that they respect the ordering. For example

$$r + (s+) = (r + s) + \quad \text{and} \quad 2 \cdot r+ = (2r)+.$$

Starting with the filtrations (10) and (12) we define for $\alpha \in \Phi$ and $r \in \mathbb{R}$:

$$(32) \quad \begin{aligned} U_{\alpha, r+} &:= \bigcup_{s > r} U_{\alpha, s}, & U_{\alpha, \infty} &:= \{1\}, \\ H_{r+} &:= \bigcup_{s > r} H_s, & H_{\infty} &:= \{1\}. \end{aligned}$$

Since the filtrations are discrete, we have $U_{\alpha, r+} = U_{\alpha, r+\epsilon}$ for sufficiently small $\epsilon > 0$, and similarly for H_{r+} .

For a function $f: \Phi \cup \{0\} \rightarrow \tilde{\mathbb{R}}$, let U_f be the subgroup of G generated by $\bigcup_{\alpha \in \Phi} U_{\alpha, f(\alpha)} \cup H_{f(0)}$. For non-empty $\Omega \subseteq A_S$ we vary on (15) by

$$(33) \quad f_{\Omega}^*: \Phi \cup \{0\} \rightarrow \tilde{\mathbb{R}}, \quad \alpha \mapsto \begin{cases} \langle \Omega, -\alpha \rangle + & \text{if } \alpha \text{ is constant on } \Omega, \\ \sup_{x \in \Omega} \langle x, -\alpha \rangle & \text{otherwise.} \end{cases}$$

For $e \in \mathbb{R}_{\geq 0}$, we define

$$U_{\Omega}^{(e)} := U_{f_{\Omega}^* + e}.$$

Notice that the closure $\overline{\Omega}$ of Ω yields $f_{\overline{\Omega}}^* = f_{\Omega}^*$ and hence $U_{\overline{\Omega}}^{(e)} = U_{\Omega}^{(e)}$.

Example 5.1. Let $G = \mathrm{GL}_n(\mathbb{F})$. We identify the standard apartment A_S of $\mathcal{B}(\mathrm{GL}_n, \mathbb{F})$ with $\mathbb{R}^n / \mathbb{R}(1, 1, \dots, 1)$, such that the set of vertices is the image of \mathbb{Z}^n . Denote the smallest integer larger than $r+ \in \tilde{\mathbb{R}}$ by $[r+]$. Recall the fractional ideals \mathfrak{P}^m in \mathbb{F} for $m \in \mathbb{Z}$. For a point $x = (x_1, \dots, x_n) \in A_S$ and $e \in \mathbb{R}_{\geq 0}$ we have

$$U_x^{(e)} = \begin{pmatrix} 1 + \mathfrak{P}^{[e+]} & \mathfrak{P}^{[x_2 - x_1 + e+]} & \dots & \mathfrak{P}^{[x_n - x_1 + e+]} \\ \mathfrak{P}^{[x_1 - x_2 + e+]} & 1 + \mathfrak{P}^{[e+]} & \dots & \mathfrak{P}^{[x_n - x_{n-1} + e+]} \\ \vdots & \vdots & \ddots & \vdots \\ \mathfrak{P}^{[x_1 - x_n + e+]} & \mathfrak{P}^{[x_{n-1} - x_n + e+]} & \dots & 1 + \mathfrak{P}^{[e+]} \end{pmatrix}$$

If $e \in \mathbb{Z}_{\geq 0}$ and $\Omega \subset A_S$ is the standard chamber, defined by $x_1 > x_2 > \dots > x_n > x_1 - 1$, then

$$U_{\Omega}^{(e)} = \begin{pmatrix} 1 + \mathfrak{P}^{e+1} & \mathfrak{P}^e & \dots & \mathfrak{P}^e \\ \mathfrak{P}^{e+1} & 1 + \mathfrak{P}^{e+1} & \dots & \mathfrak{P}^e \\ \vdots & \vdots & \ddots & \vdots \\ \mathfrak{P}^{e+1} & \mathfrak{P}^{e+1} & \dots & 1 + \mathfrak{P}^{e+1} \end{pmatrix}.$$

Notice that $U_{\Omega}^{(0)}$ is contained in the standard Iwahori subgroup of $\mathrm{GL}_n(\mathbb{F})$, and that they are not equal because the diagonal entries differ.

The groups $U_{\Omega}^{(e)}$ satisfy the following *unique decomposition property*.

Proposition 5.2 ([3, 6.4.48]). *For any ordering of Φ^{red} the product map*

$$H_{e+} \times \prod_{\alpha \in \Phi^{\mathrm{red}}} (U_{\Omega}^{(e)} \cap U_{\alpha}) \rightarrow U_{\Omega}^{(e)}$$

is a diffeomorphism. Moreover $U_\Omega^{(e)} \cap N_G(S) = H_{e+}$ and for $\alpha \in \Phi^{\text{red}}$

$$U_\Omega^{(e)} \cap U_\alpha = \begin{cases} U_{\alpha, f_\Omega^*(\alpha)+e} & \text{if } 2\alpha \notin \Phi, \\ U_{\alpha, f_\Omega^*(\alpha)+e} \cdot U_{2\alpha, f_\Omega^*(2\alpha)+e} & \text{if } 2\alpha \in \Phi. \end{cases}$$

By a diffeomorphism between p -adic algebraic varieties we mean a homeomorphism f , such that f and f^{-1} are given locally by convergent power series. The above product map is obviously algebraic, but its inverse need not be.

There is a version of the unique decomposition property with $\Phi^{\text{red}} \cup \{0\}$ instead of Φ^{red} . It follows easily from Proposition 5.2, since H_{e+} normalises $U_{\alpha,r}$.

The above decomposition implies that the subgroups $U_\Omega^{(e)}$ behave well with respect to field extensions and Levi subgroups.

Lemma 5.3. *Let $\tilde{\mathbb{F}}/\mathbb{F}$ be a finite field extension and let $\tilde{U}_\Omega^{(e)} \subseteq \mathcal{G}(\tilde{\mathbb{F}})$ be defined like $U_\Omega^{(e)} \subseteq \mathcal{G}(\mathbb{F})$. Then $U_\Omega^{(e)} = \tilde{U}_\Omega^{(e)} \cap \mathcal{G}(\mathbb{F})$.*

Proof. Let \tilde{S} and ρ_S be as on page 7 and let $\tilde{A}_{\tilde{S}} \supseteq A_S$ be the corresponding apartment of $\mathcal{B}(\mathcal{G}, \tilde{\mathbb{F}})$. Then $\tilde{f}_\Omega^*(\alpha) = f_\Omega^*(\rho_S(\alpha))$ for all $\alpha \in \tilde{\Phi}$. Now apply Proposition 5.2 and Equations (10) and (12). \square

Let $M_D = \mathcal{M}_D(\mathbb{F})$ be a standard Levi subgroup of G . Then a maximal split torus S of G is a maximal split torus of M_D as well, and the standard apartment of $\mathcal{B}(\mathcal{M}_D, \mathbb{F})$ is

$$A_D := (X_*(S) / X_*(Z_c(M_D))) \otimes_{\mathbb{Z}} \mathbb{R} = (X_*(S) / X_*(\mathcal{S}_D)) \otimes_{\mathbb{Z}} \mathbb{R}.$$

Since $S_\Delta \subseteq S_D$, there is a quotient map between the apartments

$$(34) \quad A_S \rightarrow A_{S_D}, \quad x \mapsto x_D,$$

in the buildings for G and M_D .

Lemma 5.4. *Let Ω_D be the image of Ω in the standard apartment A_D of the building for M_D . Then $U_{\Omega_D}^{(e)} = U_\Omega^{(e)} \cap M_D$ and*

$$U_\Omega^{(e)} = (U_\Omega^{(e)} \cap R_u(P_D)) (U_\Omega^{(e)} \cap M_D) (U_\Omega^{(e)} \cap R_u(\bar{P}_D)).$$

Proof. For $\Omega \subseteq A_S$ and $\alpha \in \Phi_D$ we clearly have $f_{\Omega_D}^*(\alpha) = f_\Omega^*(\alpha)$. As the groups $U_{\alpha,r}$ and H_r are the same in M_D and in G , the statement follows from Proposition 5.2. \square

We are mainly interested in the cases where Ω is a point, a facet or a polysimplex.

Theorem 5.5. *For a point x , a polysimplex σ , and a general subset Ω of an apartment A_S , the following hold:*

- (a) $U_\Omega^{(e)}$ is open if Ω is bounded.
- (b) $U_\Omega^{(e)}$ is compact.
- (c) $U_\Omega^{(e)}$ is normal in P_Ω .
- (d) $U_x^{(e)}$ fixes the star of x pointwise.
- (e) $U_\sigma^{(e)} = \prod_{x \text{ vertex of } \sigma} U_x^{(e)}$ if $e \in \mathbb{Z}_{\geq 0}$.
- (f) If x is an interior point of σ and $e \in \mathbb{Z}_{\geq 0}$, then $U_x^{(e)} = U_\sigma^{(e)}$.
- (g) $U_\Omega^{(e)} \supseteq U_\Omega^{(e')}$ whenever $e \leq e'$.
- (h) The groups $U_\sigma^{(e)}$ for $e \in \mathbb{N}$ form a neighbourhood basis of 1 in G .
- (i) The group generated by the commutators $[U_\Omega^{(e)}, U_\Omega^{(e')}]$ is contained in $U_\Omega^{(e+e')}$.

Since $U_{\alpha,r} = \{1\}$ if and only if $r = \infty$, (a) follows from Proposition 5.2. Statements (c) and (d) show that the order of the product in (e) does not matter. The proofs of (b)–(e) and (g)–(h) may be found in [18, Section I.2]. Property (f) is [23, Proposition

1.1], whereas (i) follows from [3, 6.4.41]. Notice that so far these properties hold only for subsets of the standard apartment A_S . However, (c) allows us to define

$$(35) \quad U_\Omega^{(e)} := g U_{g^{-1}\Omega}^{(e)} g^{-1}$$

for any non-empty subset Ω of an apartment gA_S . Now Theorem 5.5 holds in the entire building $\mathcal{B}(\mathcal{G}, \mathbb{F})$.

We need one more important property. We define the hull $\mathcal{H}(\sigma, \tau)$ of two polysimplices σ and τ as the intersection of all apartments containing $\sigma \cup \tau$. This finite polysimplicial complex is a combinatorial approximation to the closed convex hull of $\sigma \cup \tau$. Similarly, we can define the hull $\mathcal{H}(x, z)$ of two arbitrary points $x, z \in \mathcal{B}(\mathcal{G}, \mathbb{F})$. The proof of [23, Lemma 1.28] yields

$$(a) \text{ If } x, z \in \mathcal{B}(\mathcal{G}, \mathbb{F}) \text{ and } y \in \mathcal{H}(x, z), \text{ then } U_y^{(e)} \subseteq U_x^{(e)} U_z^{(e)}.$$

The fixed points of the groups $U_{\alpha, k}$ in the standard apartment are described by [3, 7.44]:

$$(36) \quad \begin{aligned} A_S^{U_{\alpha, k}} &= \{x \in A_S : \langle x, \alpha \rangle \geq -k\}, \\ A_S^{U_{\alpha, k+}} &= \{x \in A_S : \langle x, \alpha \rangle \geq -k - n_\alpha^{-1}\}. \end{aligned}$$

for all $\alpha \in \Phi$ and $k \in \Gamma_\alpha$. Let $\lfloor r \rfloor_{\Gamma_\alpha}$ for $r \in \mathbb{R}$ denote the largest element of Γ_α that is *strictly* smaller than r . For $x \in A_S$, (36), Proposition 5.2 and Theorem 5.5.(c) yield

$$(37) \quad \begin{aligned} A_S^{U_x^{(e)}} &= \{y \in A_S : \langle y, \alpha \rangle \geq \lfloor \alpha(x) - e \rfloor_{\Gamma_\alpha} \text{ for all } \alpha \in \Phi\}, \\ \mathcal{B}(\mathcal{G}, \mathbb{F})^{U_x^{(e)}} &= P_x \cdot A_S^{U_x^{(e)}}. \end{aligned}$$

5.1. The level of representations. The system of subgroups $(U_x^{(e)})_{x \in \mathcal{B}(\mathcal{G}, \mathbb{F})^\circ}$ for fixed $e \in \mathbb{Z}_{\geq 0}$ is a “consistent equivariant system of subgroups” in the terminology of [11, § 2.2] because of properties (b), (e), and (a) in Theorem 5.5 and (35). The main result of [11], which was inspired by [9, Section 7.1], uses these subgroups to construct resolutions of G -representations and suitable subsets thereof. We now describe this in greater detail.

Let π be a representation of G on a $\mathbb{Z}[1/p]$ -module V , where p is the characteristic of the residue field of \mathbb{F} . For any polysimplicial subcomplex $\Sigma \subseteq \mathcal{B}(\mathcal{G}, \mathbb{F})$ we define

$$C_n(\Sigma; V) := \bigoplus_{\sigma \in \Sigma^n} V^{U_\sigma^{(e)}} \otimes_{\mathbb{Z}} \mathbb{Z}\{\sigma\}.$$

If τ is a face of σ , then $U_\tau^{(e)} \subseteq U_\sigma^{(e)}$ by Theorem 5.5.(e) above, so that $V^{U_\tau^{(e)}} \supseteq V^{U_\sigma^{(e)}}$. Fix any orientation of $\mathcal{B}(\mathcal{G}, \mathbb{F})$ and declare σ endowed with the opposite orientation to be equal to $-\sigma \in \mathbb{Z}\{\sigma\}$. We define a boundary map

$$(38) \quad \partial_n : C_n(\Sigma; V) \rightarrow C_{n-1}(\Sigma; V), \quad v \otimes \sigma \mapsto v \otimes \partial(\sigma).$$

Here $\partial(\sigma)$ is the usual boundary of σ , a weighted sum of codimension-one faces of σ . This yields a chain complex $(C_*(\Sigma; V), \partial_*)$, that is, $\partial^2 = 0$. We augment it by

$$(39) \quad \partial_0 : C_0(\Sigma; V) \rightarrow V, \quad v \otimes x \mapsto v.$$

If $g \in G$ and $g \cdot \Sigma \subseteq \Sigma$, then g acts on $C_*(\Sigma; V)$ by

$$g \cdot (v \otimes \sigma) = \pi(g)v \otimes g \cdot \sigma,$$

where $g \cdot \sigma$ is endowed with the orientation coming from σ .

Theorem 5.6 ([11, Theorem 2.4]). *Let Σ be a convex subcomplex of $\mathcal{B}(\mathcal{G}, \mathbb{F})$, let $e \in \mathbb{Z}_{\geq 0}$, and let $\pi: G \rightarrow \text{Aut}(V)$ be a representation as above. Then $(C_*(\Sigma; V), \partial_*)$ is exact in all positive degrees, and the augmentation map ∂_0 induces a bijection*

$$H_0(\Sigma; V) \cong \sum_{x \in \Sigma^\circ} V^{U_x^{(e)}}.$$

Definition 5.7. A (smooth) G -representation V has *level* $e \in \mathbb{Z}_{\geq 0}$ if

$$V = \sum_{x \in \mathcal{B}(\mathcal{G}, \mathbb{F})^\circ} V^{U_x^{(e)}}.$$

This level is similar to the depth of a representation defined by Vignéras in [22, II.5.7], generalising [12]. More precisely, if V is irreducible and e is the smallest integer such that V has level e , then the depth of V lies in $(e-1, e]$. The category of G -representations of level e is studied in [11, Section 3]. If V is a complex G -representation of level e and $\Sigma = \mathcal{B}(\mathcal{G}, \mathbb{F})$, then Theorem 5.6 recovers a result of Schneider and Stuhler [18, II.3.1]. As we will see later, Theorem 5.6 for finite subcomplexes has independent significance.

Let P be a parabolic subgroup of G with unipotent radical $R_u(P)$. We let

$$V(R_u(P)) := \text{span}\{\pi(g)v - v : g \in R_u(P)\}, \quad V_{R_u(P)} := V / V(R_u(P)).$$

The representation $(\pi_{R_u(P)}, V_{R_u(P)})$ of P or $P/R_u(P)$ is called the (unnormalised) *parabolic restriction* of V .

Let (ρ, W) be a smooth representation of $P/R_u(P)$. Inflate it to a representation of P and construct the smoothly induced G -representation $\text{Ind}_P^G(W)$. This is known as the (unnormalised) *parabolic induction* of W .

Proposition 5.8. *Let $P \subseteq G$ be a parabolic subgroup.*

- (a) *If V is a G -representation of level e , then $V_{R_u(P)}$ is a representation of $P/R_u(P)$ of level e .*
- (b) *If W is a representation of $P/R_u(P)$ of level e , then $\text{Ind}_P^G(W)$ has level e .*

Proof. We first establish (a). We may assume that $P = P_D$ is a standard parabolic subgroup. Then $U^+ \subseteq P_D$. [3, Proposition 7.3.1] yields $G = P_D N_G(S) U_C$ for any chamber $C \subseteq A_S$. Since \overline{C} is a fundamental domain for the action of G on $\mathcal{B}(\mathcal{G}, \mathbb{F})$,

$$\mathcal{B}(\mathcal{G}, \mathbb{F})^\circ = G \cdot \overline{C}^\circ = P_D N_G(S) U_C \overline{C}^\circ = P_D N_G(S) \overline{C}^\circ = P_D A_S^\circ.$$

The definition of the level and Lemma 5.4 yield

$$\begin{aligned} V &= \sum_{x \in \mathcal{B}(\mathcal{G}, \mathbb{F})^\circ} V^{U_x^{(e)}} = \sum_{p \in P_D} \sum_{x \in A_S^\circ} p \cdot V^{U_x^{(e)}} \\ &\subseteq \sum_{p \in P_D} \sum_{x \in A_S^\circ} p \cdot V^{U_x^{(e)} \cap M_D} = \sum_{p \in P_D} \sum_{x_D \in A_D^\circ} p \cdot V^{U_{x_D}^{(e)}}. \end{aligned}$$

This implies that $V_{R_u(P_D)}$ has level e as well:

$$V_{R_u(P_D)} = \sum_{p \in P_D} \sum_{x_D \in A_D^\circ} p \cdot V_{R_u(P_D)}^{U_{x_D}^{(e)}} = \sum_{x_D \in \mathcal{B}(M_D, \mathbb{F})^\circ} V_{R_u(P_D)}^{U_{x_D}^{(e)}}.$$

Now we establish (b). For notational convenience, we assume that $P = P_D$ is standard parabolic, so that we may identify $P/R_u(P)$ with $M_D = \mathcal{M}_D(\mathbb{F})$. A representation of M_D has level e if and only if it is a quotient of a direct sum of copies of the regular representation on $C_c^\infty(M_D / U_{x_D}^{(e)})$ for points x_D in the building of M_D ; here C_c^∞ denotes the space of locally constant functions with compact support. Since Jacquet induction preserves direct sums and quotients, it suffices to prove that the

Jacquet induction of $C_c^\infty(M_D/U_{x_D}^{(e)})$ has level e . Inspection shows that this Jacquet induction is isomorphic to the regular representation on $C_c^\infty(G/R_u(P_D)U_{x_D}^{(e)})$.

The subgroup $R_u(P_D)U_{x_D}^{(e)}$ of G is an inductive limit of compact subgroups because $U_{x_D}^{(e)}$ is compact and $R_u(P_D)$ is unipotent. It is useful to choose a special sequence of compact subgroups exhausting $R_u(P_D)$, namely,

$$K_n := \gamma^n(U_{x_D}^{(e)} \cap R_u(P_D))\gamma^{-n},$$

where γ is a central element of M_D that is strictly positive, that is, $\bigcup K_n = R_u(P_D)$. We also consider the subgroups $\bar{K}_n := \gamma^n(U_{x_D}^{(e)} \cap R_u(\bar{P}_D))\gamma^{-n}$ in the opposite unipotent group; then $\bigcap \bar{K}_n = \{1\}$.

The space $C_c^\infty(G/R_u(P_D)U_{x_D}^{(e)})$ is the coinvariant space for the right action of $R_u(P_D)U_{x_D}^{(e)}$ on $C_c^\infty(G)$. This coinvariant space for an increasing union of compact subgroups is the inductive limit

$$C_c^\infty(G/R_u(P_D)U_{x_D}^{(e)}) \cong \varinjlim C_c^\infty(G/K_n U_{x_D}^{(e)}) \cong \varinjlim C_c^\infty(G / \gamma^n(U_x^{(e)} \cap P_D)\gamma^{-n}).$$

Here x is a pre-image of x_D in the building for G for the map in (34). Thus $U_x^{(e)} \cap M_D = U_{x_D}^{(e)}$ and

$$U_x^{(e)} = (U_x^{(e)} \cap R_u(P_D)) \cdot (U_x^{(e)} \cap M_D) \cdot (U_x^{(e)} \cap R_u(\bar{P}_D)).$$

Any smooth compactly supported function on $G/\gamma^n(U_x^{(e)} \cap P_D)\gamma^{-n}$ is invariant under right translation by \bar{K}_m for sufficiently large m because $\bigcap \bar{K}_m = 1$. Hence we may rewrite

$$\begin{aligned} C_c^\infty(G/R_u(P_D)U_{x_D}^{(e)}) &\cong \varinjlim_{n,m} C_c^\infty(G / \bar{K}_m \gamma^n(U_x^{(e)} \cap P_D)\gamma^{-n}) \\ &\cong \varinjlim_n C_c^\infty(G / \bar{K}_n \gamma^n(U_x^{(e)} \cap P_D)\gamma^{-n}) \cong \varinjlim_n C_c^\infty(G / \gamma^n U_x^{(e)} \gamma^{-n}). \end{aligned}$$

Since the regular representations on $C_c^\infty(G/\gamma^n U_x^{(e)} \gamma^{-n}) \cong C_c^\infty(G/U_x^{(e)})$ have level e , so has their inductive limit. Hence $C_c^\infty(G/R_u(P_D)U_{x_D}^{(e)})$ has level e as asserted. \square

6. CHARACTERS OF ADMISSIBLE REPRESENTATIONS

We define the character of an admissible representation first as a distribution and then describe how to interpret it as a locally constant function on suitable open subsets. Our discussion is purely algebraic and also works for representations over arbitrary fields whose characteristic is different from the characteristic p of the residue field of \mathbb{F} .

There is a Haar measure μ on G such that $\mu(K) \in \mathbb{Z}[1/p]$ for all compact open subgroups $K \subseteq G$ by [11, Lemma 1.1]. Let $\mathcal{H}(G, \mathbb{Z}[1/p])$ be the $\mathbb{Z}[1/p]$ -module of locally constant functions $G \rightarrow \mathbb{Z}[1/p]$ with compact support. Define the convolution product of $f_1, f_2 \in \mathcal{H}(G, \mathbb{Z}[1/p])$ by

$$(f_1 * f_2)(h) = \int_G f_1(g) f_2(g^{-1}h) d\mu(g).$$

We call $\mathcal{H}(G, \mathbb{Z}[1/p])$ endowed with this multiplication the *Hecke algebra*. It is an associative idempotent, non-unital $\mathbb{Z}[1/p]$ -algebra. Every element of G naturally defines a multiplier of $\mathcal{H}(G, \mathbb{Z}[1/p])$, but is not contained in $\mathcal{H}(G, \mathbb{Z}[1/p])$. Given a pro- p compact open subgroup $K \subseteq G$, we let

$$\langle K \rangle = \mu(K)^{-1} 1_K \in \mathcal{H}(G, \mathbb{Z}[1/p])$$

be the corresponding idempotent.

A smooth representation π of G on a $\mathbb{Z}[1/p]$ -module V becomes a $\mathcal{H}(G, \mathbb{Z}[1/p])$ -module in a natural way, and we have $\langle K \rangle V = V^K$, the module of K -invariant

vectors in V . We call an $\mathcal{H}(G, \mathbb{Z}[1/p])$ -module W *smooth* if $W = \varinjlim \langle K \rangle W$, where the limit runs over all pro- p compact open subgroups K of G . There is a natural equivalence between the following categories:

- smooth representations of G on $\mathbb{Z}[1/p]$ -modules,
- smooth $\mathcal{H}(G, \mathbb{Z}[1/p])$ -modules

(see [11, Proposition 1.3]). We say that a representation G on a \mathbb{K} -vector space V has *good characteristic* if the characteristic of the field \mathbb{K} does not equal p .

In good characteristic, we may define the algebra $\mathcal{H}(G, \mathbb{K})$, whose smooth modules are in bijection with smooth representations of G on \mathbb{K} -vector spaces. Such a representation (π, V) is called *admissible* if V^K has finite dimension for all compact open subgroups $K \subseteq G$. An admissible representation in good characteristic gives rise to a distribution

$$\theta_\pi: \mathcal{H}(G, \mathbb{K}) \rightarrow \mathbb{K}, \quad f \mapsto \text{tr}(\pi(f), V).$$

If $\mathbb{K} = \mathbb{C}$, then Harish-Chandra's Theorem 1.1 shows that this distribution is associated to a locally integrable function, that is, $\theta_\pi(f) = \int f(g) \cdot \text{tr}_\pi(g) d\mu(g)$ for all $f \in \mathcal{H}(G, \mathbb{C})$ and a locally integrable function tr_π . Furthermore, tr_π is locally constant on the subset of regular semisimple elements. Since this subset has full measure, the distribution θ_π is determined by the values of tr_π on regular semisimple elements. If V has infinite dimension, then tr_π is not locally constant near a unipotent element u because the closure of the conjugacy class of u contains 1 and $\text{tr}_\pi(1) = \dim V = \infty$.

Since integration requires analysis, the notion of a locally integrable function is unclear for a general field \mathbb{K} . The following definition of a character function makes sense for any field \mathbb{K} :

Definition 6.1. Let (π, V) be an admissible \mathbb{K} -linear representation of G and let $g \in G$. We write $\text{tr}_\pi(g) = \tau \in \mathbb{K}$ if there is a compact open subgroup K such that $\text{tr}(\pi(f), V) = \tau \cdot \int_G f(g) d\mu(g)$ for all $f \in \mathcal{H}(G, \mathbb{Z}[1/p])$ that are supported in KgK .

By definition, the domain of definition $\text{dom } \text{tr}_\pi$ of tr_π is open in G , and tr_π is locally constant on $\text{dom } \text{tr}_\pi$. Moreover, the trace property of θ_π forces the function tr_π to be a class function, that is, $\text{dom } \text{tr}_\pi$ is invariant under conjugation and $\text{tr}_\pi(gxg^{-1}) = \text{tr}_\pi(x)$ for all $g \in G$ and $x \in \text{dom } \text{tr}_\pi$.

In the following sections, we will show that $\text{dom } \text{tr}_\pi$ contains all regular semisimple elements, and given such an element g , we will describe a subgroup K for which tr_π is locally constant on KgK . We begin with some preparatory results. First we describe the trace distribution as a limit of locally constant functions and relate the latter to the trace function.

Let K be a compact open pro- p subgroup of G (these exist by [11, Lemma 1.1]). Since the space V^K of K -invariants in V is finite-dimensional, the linear operator $\pi(\langle K \rangle g \langle K \rangle)$ has finite rank for all $g \in G$. Hence

$$\chi_K(g) := \text{tr}(\pi(\langle K \rangle g \langle K \rangle), V) = \mu(K)^{-1} \text{tr}(\pi(1_{Kg}), V) = \mu(K)^{-1} \text{tr}(\pi(1_{gK}), V)$$

defines a K -biinvariant function on G ; here we used that $\pi(g \langle K \rangle)$, $\pi(\langle K \rangle g \langle K \rangle)$, and $\pi(\langle K \rangle g)$ have the same trace. By construction,

$$(40) \quad \text{tr}(\pi(f), V) = \int_G f(g) \chi_K(g) d\mu(g)$$

for all K -biinvariant compactly supported functions f on G . Let $(K_n)_{n \in \mathbb{N}}$ be a decreasing sequence of compact open pro- p subgroups with $\bigcap K_n = \{1\}$. Then any locally constant, compactly supported function is K_n -biinvariant for some $n \in \mathbb{N}$, so that (40) holds for $K = K_n$ for all sufficiently large n . In this sense, the trace

distribution is the limit of the locally constant functions χ_K in a distributional sense. The following lemma is trivial:

Lemma 6.2. *The trace function exists at $\gamma \in G$ and has value τ if and only if there is $n_0 \in \mathbb{N}$ with $\chi_{K_n}(g) = \tau$ for all $g \in K_{n_0} \gamma K_{n_0}$ and all $n \geq n_0$. Furthermore, then tr_π is defined and constant on $K_{n_0} \gamma K_{n_0}$.*

Let $\gamma \in G$ be a regular semisimple element. Then γ is contained in some maximal torus T . Let $T^{\text{rss}} \subseteq T$ be the subset of regular elements. It is well-known that the map

$$(41) \quad \psi: G/T \times T^{\text{rss}} \rightarrow G, \quad (gT, t) \mapsto gtg^{-1},$$

is open. We are going to quantify this statement by providing compact open subgroups $K, K_G \subseteq G$, and $K_T \subseteq T$ such that $\psi(K_G T \times K_T \gamma)$ contains $K \gamma K$ for a given regular element γ of T . We first consider the split case.

Lemma 6.3. *Suppose that T contains the maximal split torus S of G . Then the map*

$$U^+ \rightarrow U^+: u \mapsto [u, \gamma]$$

is a diffeomorphism.

Proof. For $\alpha, \beta, \alpha + \beta \in \Phi \cup \{0\}$, we have $[U_\alpha, U_\beta] \subseteq U_{\alpha+\beta}$, where we interpret U_0 as $Z_G(T)$. Let $U^{(n)}$ be the group generated by the U_α with $\alpha \in \Phi^+$ of height at least n . Then

$$U^+ = U^{(1)} \supseteq U^{(2)} \supseteq \cdots \supseteq U^{(\text{ht}(\Phi))} \supseteq \{1\}$$

is a filtration of U^+ by normal subgroups. Moreover, as algebraic groups

$$U^{(n)}/U^{(n+1)} \cong \prod_{\alpha \in \Phi^{(n)}} U_\alpha/U_{2\alpha}$$

where $\Phi^{(n)}$ denotes the set of roots of height n . The group $U_\alpha/U_{2\alpha}$ carries a canonical \mathbb{F} -vector space structure, so we can speak of λu_α for $\lambda \in \mathbb{F}$ and $u_\alpha \in U_\alpha/U_{2\alpha}$.

Given $v \in U^+$, we recursively construct $u_n \in U^{(n)}$ such that

$$[u_n \cdots u_2 \cdot u_1, \gamma] \in vU^{(n+1)}.$$

Then $u := u_{\text{ht}(\Phi)} \cdots u_2 \cdot u_1$ belongs to U^+ and satisfies $[u, \gamma] = v$. The construction will show that the u_n and hence u depend algebraically on v and that the class of u_n in $U^{(n)}/U^{(n+1)}$ is unique. It follows that the map $u \mapsto [u, \gamma]$ is bijective and that the inverse map is algebraic.

Let $w_n := [u_n \cdots u_2 \cdot u_1, \gamma]$ and define $w_0 := 1$. These elements satisfy the recursive relation

$$\begin{aligned} w_n &= u_n u_{n-1} \cdots u_1 \gamma (u_{n-1} \cdots u_1)^{-1} \gamma^{-1} \gamma u_n^{-1} \gamma^{-1} \\ &= w_{n-1} w_{n-1}^{-1} u_n w_{n-1} u_n^{-1} u_n \gamma u_n^{-1} \gamma^{-1} = w_{n-1} [w_{n-1}^{-1}, u_n] [u_n, \gamma]. \end{aligned}$$

If $u_n \in U^{(n)}$, then $[u_n, \gamma] \in U^{(n)}$ and $[w_{n-1}^{-1}, u_n] \in U^{(n+1)}$ because $[U^+, U^{(n)}] \subseteq U^{(n+1)}$. Since $U^{(n)}/U^{(n+1)}$ is commutative, we have $w_n \in w_{n-1} [u_n, \gamma] U^{(n+1)}$. Hence u_n must solve the equation $[u_n, \gamma] \in w_{n-1}^{-1} v U^{(n+1)}$. As

$$[u_\alpha, \gamma] = (1 - \alpha(\gamma)) u_\alpha \quad \text{for } u_\alpha \in U_\alpha/U_{2\alpha},$$

the map $[\cdot, \gamma]: U^{(n)}/U^{(n+1)} \rightarrow U^{(n)}/U^{(n+1)}$ is invertible. Since $w_{n-1}^{-1} v \in U^{(n)}$ by induction assumption, there is a unique coset $u_n U^{(n+1)}$ with $w_{n-1} [u_n, \gamma] U^{(n+1)} = v U^{(n+1)}$, and it depends algebraically on $w_{n-1}^{-1} v$. We may pick a representative in this coset by an algebraic map. If we do this in each step, then the final result u depends algebraically on v and satisfies $[u, \gamma] = v$. In each step, there is a unique way of lifting a solution of the equation $[u, \gamma] = v$ from $U^+/U^{(n)}$ to $U^+/U^{(n+1)}$;

in the first step, there is a unique solution in $U^+/U^{(2)}$. Hence there is a unique $u \in U^+$ with $[u, \gamma] = v$. \square

Proposition 6.4. *Suppose that the maximal torus T containing γ is split, so that G is split. Let A_S be the apartment corresponding to $S = T$, let $x \in A_S$, and let $r \in \mathbb{R}_{\geq \text{sd}(\gamma)}$. Then the map ψ in (41) restricts to an injective map from $(U_x^{(0)}/H_{0+}) \times H_{r+\gamma}$ onto a neighbourhood of γ that contains $U_x^{(r)}\gamma$.*

Proof. First we prove injectivity on the indicated domain. Assume $\psi(g_1T, t_1) = \psi(g_2T, t_2)$. Then $g_2^{-1}g_1t_1g_1^{-1}g_2 = t_2 \in T$. Since t_1 is regular, this implies $g_2^{-1}g_1 \in N_G(T)$. But $N_G(T) \cap U_x^{(0)} = Z_G(T) = T$, so that $g_1T = g_2T$ and therefore $t_1 = t_2$.

Since \mathcal{G} splits, the definition (8) yields $H_{r+} \subseteq T$. As $\psi(u, h\gamma) = [u, h\gamma]h\gamma$, Lemma 6.3 shows that $\psi(G/T \times H_{r+\gamma})$ contains $U^+H_{r+\gamma}$ for any positive system $\Phi^+ \subset \Phi$. We may decompose any element of $U_x^{(r)}\gamma$ as $y = y_+ \cdot y_- \cdot y_0$ with $y_{\pm} \in U^{\pm} \cap U_x^{(r)}$ and $y_0 \in H_{r+\gamma}$. There are $u_+ \in U_+$ and $u_- \in U_-$ such that

$$y_+y_0 = u_+y_0u_+^{-1} \quad \text{and} \quad y_-y_0 = u_-y_0u_-^{-1}.$$

Now $\text{sd}(y_0) = \text{sd}(\gamma) \geq 0$ and $[u_+, y_0] = y_+ \in U_x^{(r)}$ force $u_+ \in U_x^{(r-\text{sd}(\gamma))} \subseteq U_x^{(0)}$. For the same reason, $u_- \in U_x^{(r-\text{sd}(\gamma))}$. A good approximation for $\psi^{-1}(y)$ is (u_-u_+, y_0) :

$$\begin{aligned} (42) \quad \psi(u_-u_+, y_0) &= u_-u_+y_0u_+^{-1}u_-^{-1} = u_-y_+y_0u_-^{-1} \\ &= u_-y_+u_-^{-1}y_-y_0 = [u_-, y_+]y_+y_-y_0 = [u_-, y_+]y. \end{aligned}$$

Theorem 5.5.(i) yields

$$[u_-, y_+] \in [U_x^{(r-\text{sd}(\gamma))}, U_x^{(r)}] \subseteq U_x^{(2r-\text{sd}(\gamma))},$$

but we can be more precise. Let $r' > r$ the smallest number with $U_x^{(r')} \neq U_x^{(r)}$. Choose $\epsilon \in (0, r' - r)$ such that $U_x^{(\epsilon)} = U_x^{(0)}$ (this is possible because the filtrations (10) and (12) are discrete). Now Theorem 5.5.(i) yields

$$[u_-, y_+] \in U_x^{(r')}.$$

In other words, $\psi(u_-u_+, y_0) = y$ in $P_x/U_x^{(r')}$.

Next we try to find a solution of the form $\psi(u_-u_+g, ty_0) = y$. By (42) this is equivalent to

$$\psi(g, ty_0) = u_+^{-1}u_-^{-1}yu_-u_+ = (u_-u_+)^{-1}[y_+, u_-](u_-u_+)y_0.$$

Since $u_-u_+ \in U_x^{(0)} \subseteq N_G(U_x^{(r')})$, the right hand side lies in $U_x^{(r')}y_0$. Thus we transformed the original problem

$$\psi(U_x^{(0)}/H_{0+} \times H_{r+\gamma}) \supseteq U_x^{(r)}\gamma$$

to the problem

$$\psi(U_x^{(0)}/H_{0+} \times H_{r+\gamma}) \supseteq U_x^{(r')}y_0.$$

Since $H_{r+\gamma} = H_{r+y_0}$, $r' > r$ and

$$U_x^{(r')}y_0 \subseteq U_x^{(r')}H_{r+\gamma} \subsetneq U_x^{(r)}\gamma,$$

repetition of this process yields a solution $\psi^{-1}(y)$. \square

Now we consider a regular element γ of a non-split maximal torus $T = \mathcal{T}(\mathbb{F})$. Furthermore, we want to generalise the statement by allowing the choice of an arbitrary $x \in A_S$. Let $\tilde{\mathbb{F}}$ be a splitting field of \mathcal{T} , let $\tilde{G} = \mathcal{G}(\tilde{\mathbb{F}})$, and let $\tilde{T} := \mathcal{T}(\tilde{\mathbb{F}})$. This is a split maximal torus in \tilde{G} , which therefore corresponds to an apartment $\tilde{A}_{\tilde{T}}$ in the building $\mathcal{B}(\mathcal{G}, \tilde{\mathbb{F}})$. Recall the subgroups $\tilde{H}_r \subseteq \mathcal{Z}_{\mathcal{G}(\tilde{\mathbb{F}})}(\mathcal{T}(\tilde{\mathbb{F}}))$.

For $x \in \mathcal{B}(\mathcal{G}, \mathbb{F})$, let $\pi_T(x)$ be the point of $\tilde{A}_{\tilde{T}}$ that is nearest to x . Let Ψ be the root system corresponding to an apartment of $\mathcal{B}(\mathcal{G}, \tilde{\mathbb{F}})$ that contains x and $\pi_T(x)$. We define

$$(43) \quad d_T(x) := \max_{\beta \in \Psi^{\text{red}}} |\beta(\pi_T(x)) - \beta(x)|.$$

If $\tilde{\mathbb{F}}/\mathbb{F}$ is tamely ramified, then (27) shows that $\tilde{A}_{\tilde{T}} \cap \mathcal{B}(\mathcal{G}, \mathbb{F})$ is non-empty, that is, there is x with $d_T(x) = 0$.

Alternatively, let $\tilde{C} \subseteq \tilde{A}_{\tilde{T}}$ be a chamber containing $\pi_T(x)$, let $\rho_{\tilde{A}_{\tilde{T}}, \tilde{C}}: \mathcal{B}(\mathcal{G}, \tilde{\mathbb{F}}) \rightarrow \tilde{A}_{\tilde{T}}$ be the associated retraction. Then

$$d_T(x) = \max_{\alpha \in \tilde{\Phi}^{\text{red}}} |\alpha(\pi_T(x)) - \alpha(\rho_{\tilde{A}_{\tilde{T}}, \tilde{C}}(x))|.$$

Proposition 4.2.(c) yields

$$(44) \quad d_T(x) \leq \text{ht}(\Phi) \text{sd}(\gamma) \quad \text{for all } x \in \mathcal{B}(\mathcal{G}, \tilde{\mathbb{F}})^\gamma.$$

Lemma 5.3 and (33) yield

$$(45) \quad U_x^{(r+d_T(x))} = \tilde{U}_x^{(r+d_T(x))} \cap \mathcal{G}(\mathbb{F}) \subseteq \tilde{U}_{\pi_T(x)}^{(r)} \cap U_x.$$

Lemma 6.5. *Let $\gamma \in T$ be regular and let $r \in \mathbb{R}_{\geq \text{sd}(\gamma)}$. Let $x \in \mathcal{B}(\mathcal{G}, \mathbb{F})$ and abbreviate $K_x = \tilde{U}_{\pi_T(x)}^{(0)} \cap G$. Then $U_x^{(r+d_T(x))}\gamma$ is contained in $\psi(K_x \times (\tilde{H}_{r+\gamma} \cap T))$.*

Proof. Equation (45) and Proposition 6.4 show that every element of $\tilde{U}_x^{(r+d_T(x))}\gamma$ is conjugate in $\mathcal{G}(\tilde{\mathbb{F}})$ to an element of $\tilde{H}_{r+\gamma} \cap \mathcal{T}(\tilde{\mathbb{F}})$. Since the maps

$$\tilde{\psi}: (\mathcal{G}(\tilde{\mathbb{F}})/\mathcal{T}(\tilde{\mathbb{F}})) \times \mathcal{T}(\tilde{\mathbb{F}}) \rightarrow \mathcal{G}(\tilde{\mathbb{F}}) \quad \text{and} \quad \psi: (\mathcal{G}(\mathbb{F})/\mathcal{T}(\mathbb{F})) \times \mathcal{T}(\mathbb{F}) \rightarrow \mathcal{G}(\mathbb{F})$$

are injective and open, respectively on $\tilde{U}_{\pi_T(x)}^{(0)}/\tilde{H}_{0+} \times (\tilde{H}_{r+\gamma} \cap \mathcal{T}(\tilde{\mathbb{F}}))$ and on the intersection of this set with G ,

$$\tilde{\psi}(K_x \times (\tilde{H}_{r+\gamma} \cap T)) = \tilde{\psi}(\tilde{U}_{\pi_T(x)}^{(0)} \times (\tilde{H}_{r+\gamma} \cap \mathcal{T}(\tilde{\mathbb{F}}))) \cap G.$$

Moreover, by Proposition 6.4 the right hand side contains

$$(46) \quad \tilde{U}_{\pi_T(x)}^{(r)}\gamma \cap G \supseteq \tilde{U}_x^{(r+d_T(x))}\gamma \cap G = U_x^{(r+d_T(x))}\gamma. \quad \square$$

There is a decreasing sequence $(K_n)_{n \in \mathbb{N}}$ of *normal* compact open subgroups in K_x with $\bigcap K_n = \{1\}$. Since K_x is open in G , we may use this sequence to approximate the trace distribution as in (40). Since K_n is normal in K_x , then the space of K_n -biinvariant functions is invariant under conjugation by elements of K_x . This implies that the function χ_{K_n} is invariant under conjugation by elements of K_x . Therefore, Lemma 6.5 shows that χ_{K_n} is constant on $U_x^{(r+d_T(x))}\gamma$ once it is constant on $\tilde{H}_{r+\gamma} \cap T$. In the following, we may therefore restrict attention to elements of a torus in G .

7. THE LOCAL CONSTANCY OF CHARACTERS

Let (π, V) be an admissible representation of G in good characteristic, of level $e \in \mathbb{Z}_{\geq 0}$. Let γ be a regular semisimple element of a maximal torus $T \subseteq G$ and let $x \in \mathcal{B}(\mathcal{G}, \mathbb{F})^\circ$ be a vertex in the building of G . We are going to find $r(\gamma) \in \mathbb{N}$ depending only on γ and the level e of the representation, such that tr_π is defined and constant on $U_x^{(r(\gamma)+d_T(x))}$ with $d_T(x)$ as in (43).

7.1. Local constancy for compact elements. First we assume, in addition, that γ is a compact element, so that γ fixes some point in the affine building. The assertions for general elements are reduced to the compact case in Section 7.2.

Our definition of $r(\gamma)$ is somewhat complicated and probably not optimal. It is likely that $r(\gamma) = \max\{\text{sd}(\gamma), e\}$ works, but we can only prove this if T has a subtorus S that is a maximal \mathbb{F} -split torus of G .

Let $T = \mathcal{T}(\mathbb{F}) \subseteq G$ be a maximal torus containing γ and let $\tilde{\mathbb{F}}/\mathbb{F}$ be a finite Galois extension splitting \mathcal{T} . Recall the subgroups $\tilde{U}^+ \subset \mathcal{G}(\tilde{\mathbb{F}})$ and $\tilde{H}_r \subseteq \mathcal{Z}_{\mathcal{G}(\tilde{\mathbb{F}})}(\mathcal{T}(\tilde{\mathbb{F}}))$. Let \tilde{B} be a Borel subgroup of $\mathcal{G}(\tilde{\mathbb{F}})$ containing $\mathcal{T}(\tilde{\mathbb{F}})$ and let \tilde{U}_T^+ be its unipotent radical. We define $U_T^+ = \tilde{U}_T^+ \cap G$. Since \tilde{B} need not be defined over \mathbb{F} , $T \ltimes U_T^+$ need not be a Borel subgroup of G . In fact U_T^+ can be reduced to $\{1\}$, for example if T is anisotropic.

Definition 7.1. For $x \in \mathcal{B}(\mathcal{G}, \mathbb{F})$ define $d_T(x)$ as in (43) and let $d(\gamma) \in \mathbb{R}$ be the smallest number such that

$$(47) \quad \mathcal{B}(\mathcal{G}, \mathbb{F})^\gamma \subseteq U_T^+ \cdot \{x \in \mathcal{B}(\mathcal{G}, \mathbb{F}) : d_T(x) \leq d(\gamma)\}.$$

We have $d(\gamma) < \infty$ because $\mathcal{B}(\mathcal{G}, \mathbb{F})^\gamma/T$ is compact.

Theorem 7.2. Define $r(\gamma) := \max\{\text{ht}(\Phi)\text{sd}(\gamma), e + d(\gamma)\}$.

- (a) *The function tr_π is defined and constant on $\gamma\tilde{H}_{r(\gamma)}^+ \cap T$, and on all G -conjugacy classes intersecting this set.*
- (b) *The function tr_π is constant on $U_x^{(r(\gamma)+d_T(x))}\gamma$, for any $x \in \mathcal{B}(\mathcal{G}, \mathbb{F})$.*
- (c) *If T has a subtorus S that is a maximal \mathbb{F} -split torus of G , then $d(\gamma) = 0$ and we may omit the factor $\text{ht}(\Phi)$ in the definition of $r(\gamma)$, that is, tr_π is constant on $\gamma\tilde{H}_{\max\{\text{sd}(\gamma), e\}}^+ \cap T$.*

If $\tilde{\mathbb{F}}/\mathbb{F}$ is tamely ramified, then (27) shows that there is a point $x \in \mathcal{B}(\mathcal{G}, \mathbb{F})$ with $d_T(x) = 0$, so that tr_π is constant on $U_x^{(r(\gamma))}\gamma$.

The number $r(\gamma)$ will reappear frequently in the following. We will not need the definition of $r(\gamma)$ but only Theorem 7.2.(a). That is, the following results remain true for a smaller value of $r(\gamma)$ provided Theorem 7.2.(a) can be established for it.

Proof. (a) Theorem 5.6 implies a formula for $\text{tr}(\pi(f), V)$, which is worked out in [11, Proposition 4.1]. We need some notation to state this trace formula. For $g \in G$, let Σ^g be the set of all polysimplices $\sigma \in \Sigma$ with $g\sigma = \sigma$ and let $\epsilon_\sigma(g) = \pm 1$, depending on whether the automorphism of σ induced by g preserves or reverses orientation. For a locally constant function f supported in P_x , [11, Proposition 4.1] asserts

$$(48) \quad \text{tr}(\pi(f), V) = \lim_{\Sigma} \int_{g \in P_x} f(g) \sum_{\sigma \in \Sigma^g} (-1)^{\dim \sigma} \epsilon_\sigma(g) \text{tr}(\pi(g), V^{U_\sigma^{(e)}}) \, d\mu(g),$$

where the limit means that there is a finite convex subcomplex Σ_0 such that the right hand side is the same for all P_x -invariant finite convex subcomplexes Σ of $\mathcal{B}(\mathcal{G}, \mathbb{F})$ with $\Sigma \supseteq \Sigma_0$. Thus we want to show that the function

$$(49) \quad \tau_\Sigma : g \mapsto \sum_{\sigma \in \Sigma^g} (-1)^{\dim \sigma} \epsilon_\sigma(g) \text{tr}(\pi(g), V^{U_\sigma^{(e)}})$$

is constant on $U_x^{(r(\gamma)+d_T(x))}\gamma$ for all sufficiently large P_x -invariant finite convex subcomplexes Σ . The function τ_Σ is invariant under conjugation by elements of P_x because Σ is P_x -invariant.

Lemma 4.3 yields $\mathcal{B}(\mathcal{G}, \mathbb{F})^g = \mathcal{B}(\mathcal{G}, \mathbb{F})^\gamma$ for all $g \in \tilde{H}_{r(\gamma)}^+ \gamma \cap T$, because $r(\gamma) \geq \text{ht}(\Phi)\text{sd}(\gamma)$. Since

$$(50) \quad \tilde{H}_{e+d_T(x)}^+ \subseteq \tilde{U}_{\pi_T(x)}^{(e+d_T(x))} \subseteq \tilde{U}_x^{(e)},$$

the operator $\pi(g^{-1}\gamma)$ restricts to the identity on $V^{U_x^{(e)}}$, for all x with $d_T(x) \leq d(\gamma)$.

Let \mathcal{D} be a set of simplices in $\mathcal{B}(\mathcal{G}, \mathbb{F})^\gamma$, such that \mathcal{D} is a fundamental domain for the action of U_T^+ on $U_T^+ \cdot \mathcal{B}(\mathcal{G}, \mathbb{F})^\gamma$ and every $\sigma \in \mathcal{D}$ contains an interior point x with $d_T(x) \leq d(\gamma)$. Equation (49) becomes

$$(51) \quad \tau_\Sigma(g) = \sum_{u\sigma \in \Sigma^g} \epsilon_{u\sigma}(g) \operatorname{tr}(\pi(g), V^{U_{u\sigma}^{(e)}}) = \sum_{u\sigma \in \Sigma^g} \epsilon_\sigma(u^{-1}gu) \operatorname{tr}(\pi(u^{-1}gu), V^{U_\sigma^{(e)}}),$$

where the sum runs over all polysimplices $u\sigma \in \Sigma^g = \Sigma^\gamma$ with $\sigma \in \mathcal{D}$ and $u \in U_T^+$. Notice that we pick only one u for each such polysimplex. Given another $u_1 \in U_T^+$ with $u_1\sigma = u\sigma$, we have $u_1^{-1}u \in P_\sigma$, so $\theta(u_1, g) = \theta(u, g)$, where

$$\theta(u, g) := \epsilon_\sigma(u^{-1}gu) \operatorname{tr}(\pi(u^{-1}gu), V^{U_\sigma^{(e)}}).$$

Recall that

$$\{u \in \tilde{U}_T^+ : [u^{-1}, g] \in \tilde{P}_\sigma\} = \{u \in \tilde{U}_T^+ : g(u\sigma) = u\sigma\}.$$

The equality $\Sigma^g = \Sigma^\gamma$ implies that

$$\{u \in \tilde{U}^+ : [u^{-1}, g] \in \tilde{P}_\sigma\} = \{u \in \tilde{U}^+ : [u^{-1}, \gamma] \in \tilde{P}_\sigma\}.$$

We denote this set by $\tilde{U}_T^+(\sigma)$ and we write $U_T^+(\sigma) = \tilde{U}_T^+(\sigma) \cap G$. Let μ be a Haar measure on U_T^+ , normalised so that $\mu(U_T^+(\sigma)) = 1$. Now we can rewrite (51) as $\tau_\Sigma(g) = \sum_{\sigma \in \mathcal{D}} \tau_{\Sigma, \sigma}(g)$, where

$$(52) \quad \tau_{\Sigma, \sigma}(g) = \sum_{u\sigma \in \Sigma^g \cap U_T^+ \sigma} \theta(u, g) = \sum_{u \in U_T^+(\sigma) / U_T^+ \cap P_\sigma} \theta(u, g) = \int_{U_T^+(\sigma)} \theta(u, g) d\mu(u).$$

By Lemma 6.3 the map

$$\phi_g : \tilde{U}^+ \rightarrow \tilde{U}^+ : u \mapsto [u^{-1}, g]$$

is a diffeomorphism. We claim that $\phi_g^{-1}(U_T^+) = U_T^+$.

It is clear that $\phi_g^{-1}(U_T^+) \supset U_T^+$. Suppose that $u \in \phi_g^{-1}(U_T^+)$, that is, $u^{-1}gug^{-1} \in \tilde{U}_T^+ \cap G$. For any $\tau \in \operatorname{Gal}(\tilde{\mathbb{F}}/\mathbb{F})$ we have

$$G \ni u^{-1}gu = \tau(u^{-1}gu) = \tau(u^{-1})g\tau(u),$$

so $\tau(u)u^{-1} \in Z_{\mathcal{G}(\tilde{\mathbb{F}})}(g)$. Since \tilde{U}_T^+ is unipotent and $\tilde{\mathbb{F}}$ -split, the same holds for λu with $\lambda \in \tilde{\mathbb{F}}$. Hence

$$\tau(u)u^{-1} \in Z_{\mathcal{G}(\tilde{\mathbb{F}})}(g)^\circ = \mathcal{T}(\tilde{\mathbb{F}}).$$

Also $\tau(u)$ is unipotent and $\tau(u) \in \mathcal{T}(\tilde{\mathbb{F}})u \subset \tilde{B}$, we must have $\tau(u) \in \tilde{U}_T^+$. Then $\tau(u)u^{-1} \in \tilde{U}_T^+ \cap \mathcal{T}(\tilde{\mathbb{F}}) = \{1\}$, which means that $\tau(u) = u$. Hence $u \in \tilde{U}_T^+ \cap \mathcal{G}(\tilde{\mathbb{F}})^{\operatorname{Gal}(\tilde{\mathbb{F}}/\mathbb{F})} = U_T^+$.

As $\tilde{U}_T^+(\sigma) = \phi_g^{-1}(\tilde{U}_T^+(\sigma) \cap \tilde{P}_\sigma)$, ϕ_g restricts to a diffeomorphism

$$\phi_g : U_T^+(\sigma) \rightarrow P_\sigma \cap U_T^+.$$

The same statements hold for $\phi_\gamma : \tilde{U}_T^+ \rightarrow \tilde{U}_T^+$. For every $u_1 \in U_T^+(\sigma)$, thought of as appearing in $\tau_{\Sigma, \sigma}(g)$, the element $u_2 = \phi_\gamma^{-1}(\phi_g(u_1))$ appears in $\tau_{\Sigma, \sigma}(\gamma)$. We do not know whether $u_1\sigma$ equals $u_2\sigma$ or not, fortunately that is not needed. Now

$$\begin{aligned} \theta(u_2, \gamma) &= \epsilon_\sigma([u_2^{-1}, \gamma]\gamma) \operatorname{tr}(\pi([u_2^{-1}, \gamma]\gamma), V^{U_\sigma^{(e)}}) \\ &= \epsilon_\sigma([u_1^{-1}, g]g(g^{-1}\gamma)) \operatorname{tr}(\pi([u_1^{-1}, g])\pi(g)\pi(g^{-1}\gamma), V^{U_\sigma^{(e)}}). \end{aligned}$$

Since $\Sigma^g = \Sigma^\gamma$, $g^{-1}\gamma$ fixes σ pointwise, while in view of (50) and the definition of \mathcal{D} , $\pi(g^{-1}\gamma)$ acts as the identity on $V^{U_\sigma^{(e)}}$. Therefore

$$(53) \quad \theta(u_2, \gamma) = \epsilon_\sigma([u_1^{-1}, g]g) \operatorname{tr}(\pi([u_1^{-1}, g]g), V^{U_\sigma^{(e)}}) = \theta(u_1, g),$$

which shows that every term of the integral (51) also occurs in $\tau_{\Sigma,\sigma}(\gamma)$.

The proof of Lemma 6.3 shows that (with respect to any basepoint in \tilde{U}_T^+) the generalised eigenvalues of the differentials

$$D\phi_\gamma, D\phi_g : \text{Lie}_{\tilde{\mathbb{F}}}(\tilde{U}_T^+) \rightarrow \text{Lie}_{\tilde{\mathbb{F}}}(\tilde{U}_T^+)$$

are $\{1 - \alpha(\gamma) : \alpha \in \tilde{\Phi}^+\}$ and $\{1 - \alpha(g) : \alpha \in \tilde{\Phi}^+\}$, and they occur with multiplicity $d_\alpha := \dim \text{Lie}_{\tilde{\mathbb{F}}}(\mathcal{U}_\alpha / \mathcal{U}_{2\alpha})$. The restriction $h = \gamma^{-1}g \in \tilde{H}_{\text{sd}(\gamma)+} \cap \mathcal{T}(\tilde{\mathbb{F}})$ implies

$$v(1 - \alpha(g)) = v(1 - \alpha(\gamma)\alpha(h)) = v(1 - \alpha(\gamma) + \alpha(\gamma)(1 - \alpha(h))) = v(1 - \alpha(\gamma))$$

for all $\alpha \in \tilde{\Phi}$. Hence $\phi_g, \phi_\gamma : \tilde{U}_T^+ \rightarrow \tilde{U}_T^+$ both have Jacobian $\prod_{\alpha \in \tilde{\Phi}^+} \|1 - \alpha(g)\|_{\tilde{\mathbb{F}}}^{d_\alpha}$, at any point of \tilde{U}_T^+ .

The group U_T^+ need not be defined over \mathbb{F} , but it always equals the $\text{Gal}(\tilde{\mathbb{F}}/\mathbb{F})$ -invariants in the algebraic $\tilde{\mathbb{F}}$ -group \tilde{U}_T^+ . The Galois group permutes the roots of $(\mathcal{G}, \mathcal{T})$, so the invariants are contained in the product \tilde{U}_T^\sharp of the root groups $\mathcal{U}_\alpha(\tilde{\mathbb{F}})$ for which $\text{Gal}(\tilde{\mathbb{F}}/\mathbb{F})\alpha \subset R(\tilde{B}, \tilde{T})$. The action of $\text{Gal}(\tilde{\mathbb{F}}/\mathbb{F})$ on \tilde{U}_T^\sharp is given by permutations of these roots and the Galois action on $\tilde{\mathbb{F}}$. It follows that the Galois invariant elements U_T^+ in \tilde{U}_T^\sharp form an analytic \mathbb{F} -variety, diffeomorphic to a vector space.

Knowing this, a direct calculation shows that (at any point of U_T^+) both ϕ_g and $\phi_\gamma : U_T^+ \rightarrow U_T^+$ have Jacobian $\prod_\alpha \|1 - \alpha(g)\|_{\tilde{\mathbb{F}}}^{d_\alpha/[\tilde{\mathbb{F}}:\mathbb{F}]}$, where the product runs over all $\alpha \in \tilde{\Phi}^+$ such that the restriction of α to T appears in $\text{Lie}_{\mathbb{F}}(U_T^+)$. As the diffeomorphisms $\phi_g, \phi_\gamma : U_T^+(\sigma) \rightarrow P_\sigma \cap U_T^+$ have the same Jacobian, and by (53),

$$\begin{aligned} \tau_{\Sigma,\sigma}(g) &= \int_{U_T^+(\sigma)} \theta(u, g) d\mu(u) = \int_{U_T^+(\sigma)} \theta(\phi_\gamma^{-1}(\phi_g(u)), \gamma) d\mu(u) \\ &= \int_{U_T^+(\sigma)} \theta(u, \gamma) d\mu(u) = \tau_{\Sigma,\sigma}(\gamma). \end{aligned}$$

This holds for all $\sigma \in \mathcal{D}$, so

$$\tau_\Sigma(g) = \sum_{\sigma \in \mathcal{D}} \tau_{\Sigma,\sigma}(g) = \sum_{\sigma \in \mathcal{D}} \tau_{\Sigma,\sigma}(\gamma) = \tau_\Sigma(\gamma).$$

(b) Lemma 6.5 shows that any element of $U_x^{(r(\gamma)+d_T(x))}\gamma$ is P_x -conjugate to one of $\gamma \tilde{H}_{r(\gamma)+} \cap T$. Hence (b) follows from (a).

(c) To a large extent we will copy the proof of part (a), but we take advantage of $U^+ \cdot A_S = \mathcal{B}(\mathcal{G}, \mathbb{F})$. This clearly implies $d(\gamma) = 0$, so that \mathcal{D} is a collection of simplices of A_S . This \mathcal{D} works for both γ and $g = \gamma h$. We may replace $\text{ht}(\Phi)\text{sd}(\gamma)$ by $\text{sd}(\gamma)$ in the definition on $r(\gamma)$, because the factor $\text{ht}(\Phi)$ was only needed to ensure that $\mathcal{B}(\mathcal{G}, \mathbb{F})^g$ equals $\mathcal{B}(\mathcal{G}, \mathbb{F})^\gamma$.

With these choices the proof of (a) mostly goes through, even though we do not know whether or not g and γ have the same fixed points. We now have two possibly different sets $U_T^+(\sigma)$, one for g and one for γ , but the map $\phi_\gamma^{-1} \circ \phi_g$ still provides a diffeomorphism between them. \square

7.2. Local constancy for non-compact elements. We would like to generalise Theorem 7.2 to all regular semisimple elements. This is possible using Jacquet modules and parabolic restriction as in [5]. Although the methods in [5] are algebraic and not restricted to complex coefficients, Casselman refers to earlier work which was written with complex representations in mind. This makes it hard to judge whether Casselman's proofs work for representations in good characteristic. Fortunately, Vignéras [22] proved the required results in this generality.

Let $\gamma \in T$ be a semisimple element and let $P_\gamma \subseteq G$ be the parabolic subgroup contracted by γ , which is defined in (3). Since \mathbb{F} is complete with respect to the valuation v , Proposition 2.3.(d) shows that γ is compact in M_γ . It follows from Proposition 2.3.(b) that $\text{Lie}_{\mathbb{F}}(\mathcal{R}_u(\mathcal{P}_\gamma)) \subseteq \text{Lie}_{\mathbb{F}}(\mathcal{G})$ is the sum of all eigenspaces of $\text{Ad}(\gamma)$ corresponding to eigenvalues with strictly positive valuation. (Although the eigenvalues may lie in a field extension of \mathbb{F} , this subspace is defined over \mathbb{F} .) Similarly, $R_u(P_{\gamma^{-1}})$ corresponds to the γ -eigenvalues with strictly negative valuation.

The description of (standard) parabolic subgroups in Definition 2.2 shows that M_γ contains a maximal split torus of G , say S_γ . It may happen that $\gamma \notin S_\gamma$. Let x be a point of the apartment A_γ of $\mathcal{B}(\mathcal{G}, \mathbb{F})$ corresponding to S_γ . Proposition 5.2 implies

$$(54) \quad U_x^{(e)} = (U_x^{(e)} \cap R_u(P_{\gamma^{-1}})) (U_x^{(e)} \cap M_\gamma) (U_x^{(e)} \cap R_u(P_\gamma)),$$

or, in other words, $U_x^{(e)}$ is well-placed with respect to (P_γ, M_γ) . The collection

$$X = \{gx \in \mathcal{B}(\mathcal{G}, \mathbb{F}) : g \text{ lies in the maximal compact subgroup of } T\}$$

is finite and γ -invariant. Since $T \subset M_\gamma$, the subgroup $U_{x'}^{(e)}$ is well-placed with respect to (P_γ, M_γ) for every $x' \in X$. The group $K^{(e)} := \bigcap_{x' \in X} U_{x'}^{(e)}$ is also well-placed:

$$K^{(e)} = (K^{(e)} \cap R_u(P_{\gamma^{-1}})) (K^{(e)} \cap M_\gamma) (K^{(e)} \cap R_u(P_\gamma)) =: K_-^{(e)} K_0^{(e)} K_+^{(e)}.$$

It follows that

$$\gamma K_-^{(e)} \gamma^{-1} \supseteq K_-^{(e)}, \quad \gamma K_0^{(e)} \gamma^{-1} = K_0^{(e)}, \quad \gamma K_+^{(e)} \gamma^{-1} \subsetneq K_+^{(e)},$$

so that the sequence $K^{(e)}$ for $e \in \mathbb{N}$ has all the properties claimed in [6].

Theorem 7.3 ([22, II.3.7]). *Let (π, V) be an admissible smooth G -representation in good characteristic and let $g \in G$ be such that $P_g = P_\gamma$. There exist increasing sequences of finite-dimensional vector spaces $V^{(e)} \subseteq V^{K^{(e)}}$ and $V_{R_u(P_\gamma)}^{(e)} \subseteq V_{R_u(P_\gamma)}^{K_0^{(e)}}$ such that*

- (a) $\bigcup_e V^{(e)} \oplus V(R_u(P_\gamma)) = V$ and $\bigcup_e V_{R_u(P_\gamma)}^{(e)} = V_{R_u(P_\gamma)}$,
- (b) The quotient map $V \rightarrow V/V(R_u(P_\gamma)) = V_{R_u(P_\gamma)}$ restricts to bijections $V^{(e)} \rightarrow V_{R_u(P_\gamma)}^{(e)}$ and $(\bigcup_r V^{(r)})^{K^{(e)}} \rightarrow V_{R_u(P_\gamma)}^{K_0^{(e)}}$,
- (c) $V^{(e)}$ is stable under $\pi(1_{K^{(e)} g K^{(e)}})$.

This setup allows us to use the (elementary) arguments from [5, page 104], which result in

$$(55) \quad \text{tr}(\mu(K^{(e)} g K^{(e)})^{-1} \pi(1_{K^{(e)} g K^{(e)}}), V) = \text{tr}(\pi_{R_u(P_\gamma)}(g), V_{R_u(P_\gamma)}^{K_0^{(e)}})$$

for all $g \in G$ with $P_g = P_\gamma$. Notice that the set of such g is contained in M_γ , so it is not open in G unless γ is compact in G .

Theorem 7.4. *Let γ be a regular semisimple element. Then $\text{tr}_\pi(\gamma)$ and $\text{tr}_{\pi_{R_u(P_\gamma)}}(\gamma)$ are both defined, and they are equal.*

Proof. Since γ is compact in M_γ , Theorem 7.2 tells us that $\text{tr}_{\pi_{R_u(P_\gamma)}}$ is well-defined and constant near γ . Pick an $e \in \mathbb{N}$ such that it is constant on $\gamma K_0^{(e)}$. Now (55) yields

$$\begin{aligned} \text{tr}_{\pi_{R_u(P_\gamma)}}(\gamma) &= \text{tr}(\pi_{R_u(P_\gamma)}(\gamma * \langle K_0^{(e)} \rangle), V_{R_u(P_\gamma)}) = \text{tr}(\pi_{R_u(P_\gamma)}(\gamma), V_{R_u(P_\gamma)}^{K_0^{(e)}}) \\ &= \text{tr}(\mu(K^{(e)} \gamma K^{(e)})^{-1} \pi(1_{K^{(e)} g K^{(e)}}), V). \end{aligned}$$

As the subsets $K^{(e)} \gamma K^{(e)}$ form a neighbourhood basis of γ in G , taking the limit $e \rightarrow \infty$ and invoking Lemma 6.2 shows that $\text{tr}_\pi(\gamma)$ is well-defined and equals $\text{tr}_{\pi_{R_u(P_\gamma)}}(\gamma)$. \square

This theorem, which Casselman [5] proved for complex representations, enables us to reduce the computation of traces from general semisimple elements to compact semisimple elements. Theorem 7.2 tells us on which neighbourhood of γ the function $\text{tr}_{\pi_{R_u(P_\gamma)}}$ is constant. But this is only a neighbourhood in M_γ . We also want to know on which neighbourhood in G the function tr_π is constant. Let $r(\gamma)$ be such that Theorem 7.2.(a) holds when we view γ as a compact element in M_γ .

Theorem 7.5. *Let γ be a regular element of a (not necessarily split) maximal torus T of G . Let (π, V) be an admissible representation of G of level e in good characteristic.*

- (a) *The function tr_π is defined and constant on $\tilde{H}_{r(\gamma)+\gamma} \cap T$, and on all G -conjugacy classes intersecting this set.*
- (b) *The function tr_π is constant on $U_x^{(r(\gamma)+d_T(x))}\gamma$, for any $x \in \mathcal{B}(\mathcal{G}, \mathbb{F})$.*

Proof. For every root $\alpha \in \Phi(\mathcal{G}(\tilde{\mathbb{F}}), \mathcal{T}(\tilde{\mathbb{F}}))$ and every $g \in \tilde{H}_{r(\gamma)+\gamma} \cap T$ we have $\tilde{v}(\alpha(g)) = \tilde{v}(\alpha(\gamma))$ because $g\gamma^{-1}$ is compact. Together with (5), this implies $P_g = P_\gamma$, so that Theorem 7.4 applies to all $g \in \tilde{H}_{r(\gamma)+\gamma} \cap T$ and tells us that $\text{tr}_\pi(g) = \text{tr}_{\pi_{R_u(P_\gamma)}}(g)$. Theorem 7.2 and Proposition 5.8 show that $\text{tr}_{\pi_{R_u(P_\gamma)}}$ is constant on $\tilde{H}_{r(\gamma)+\gamma} \cap T$, so the same goes for tr_π . This proves (a), from which (b) follows upon applying Lemma 6.5. \square

This theorem is similar to [1, Corollary 12.11], which was proved only for complex representations and “tame” elements γ . Our neighbourhoods of constancy are usually smaller than those in [1], because Theorem 7.2.(a) is not optimal. The results of Adler and Korman suggest that Theorem 7.2.(c) could be valid whenever the maximal torus T splits over a tamely ramified extension of \mathbb{F} . Possibly this has something to do with Rousseau’s result (26).

8. A BOUND FOR THE DIMENSION OF V^K

In this section, we will use the resolutions of [11] to estimate the dimension of $V_x^{U_x^{(e)}}$ for an admissible representation (π, V) of G in good characteristic. We abbreviate $K_e := U_x^{(e)}$.

First we estimate the growth of some related double coset spaces in order to show that our later estimates are optimal, at least for GL_n .

Since every irreducible smooth representation is a subquotient of a parabolically induced one, the essential case is $V = \text{Ind}_P^G(W)$, where P is a parabolic subgroup of G and (ρ, W) is a supercuspidal representation of $P/R_u(P)$. There is a natural isomorphism

$$(56) \quad V^{K_e} \cong \bigoplus_{PgK_e} W^{P \cap gK_e g^{-1}},$$

where the sum runs over all double (P, K_e) -cosets. The space $P \backslash G / K_e$ is finite because $P \backslash G$ is a complete algebraic variety (and hence compact in the p -adic topology) and K_e is open. We will discuss how $|P \backslash G / K_e|$ grows as e increases, under some simplifications. If P is a Borel subgroup and ρ is a character, then $|P \backslash G / K_e|$ and $\dim V^{K_e}$ have equivalent growth rates.

Suppose that G is split. Let S be a split maximal torus of G and let P_D be a standard parabolic subgroup of G . The dimension of $P_D \backslash G$ is

$$\dim(P_D \backslash G) = \dim_{\mathbb{F}}(\text{Lie}_{\mathbb{F}}(G) / \text{Lie}_{\mathbb{F}}(P_D)) = \sum_{\alpha \in \Phi^- \setminus \Phi_D^-} \dim_{\mathbb{F}} \text{Lie}_{\mathbb{F}}(U_\alpha) = |\Phi^-| - |\Phi_D^-|.$$

Let $x \in A_S$. By construction, the groups K_e decrease equally fast in every direction; if K_e corresponds to a lattice $L^{(e)}$ in $\text{Lie}_{\mathbb{F}}(G)$, then K_{e+1} corresponds to $\mathfrak{P}L^{(e)}$,

where \mathfrak{P} is the maximal ideal in the maximal compact subring of \mathbb{F} . Hence a double coset $P_D g K_e$ contains approximately $q^{\dim(P_D \backslash G)}$ double (P_D, K_{e+1}) -cosets. Therefore, $|P_D \backslash G/K_e|$ grows, in first approximation, like $q^{e \dim(P_D \backslash G)}$.

Now we focus on the easier example $G = \mathrm{GL}_n$ and let P and S be the standard Borel subgroup and the standard maximal torus in $\mathrm{GL}_n(\mathbb{F})$. The irreducible representations of $S = P/R_u(P)$ are characters. Let (ρ, \mathbb{C}) be such a character and let V be the parabolically induced representation of G . Since any character is trivial on $K_e \cap S$ for large enough e , $\mathbb{C}^{P \cap g K_e g^{-1}} \cong \mathbb{C}$ for large enough e , so that $\dim(V^{K_e}) = |P \backslash G/K_e|$ for large e . These numbers are routine to compute:

$$(57) \quad |P \backslash G/K_e| \approx e^{n-1} q^{en(n-1)/2}$$

in the sense that the quotient of both sides tends towards a constant as $e \rightarrow \infty$.

For complex representations, we may use the growth rate of $\dim V^{K_e}$ to estimate the growth of the character. It will, however, turn out that these estimates are far from optimal. The idea is simple enough: if tr_π is constant on $K_e \gamma$, then

$$\mathrm{tr}_\pi(\gamma) = \frac{1}{|K_e \gamma|} \int_{K_e \gamma} \mathrm{tr}_\pi(\gamma) d\mu(\gamma) = \mathrm{tr}(\pi(\langle K_e \rangle \gamma)).$$

Equip the finite-dimensional vector space V^{K_0} with some norm. Since the range of $\langle K_e \rangle \gamma$ is contained in $V^{K_e} \subseteq V^{K_0}$ and the largest eigenvalue of $\langle K_e \rangle \gamma$ is controlled by the operator norm $\|\langle K_0 \rangle \gamma \langle K_0 \rangle\|_\infty$, we get the estimate

$$(58) \quad |\mathrm{tr}_\pi(\gamma)| \leq \|\langle K_0 \rangle \gamma \langle K_0 \rangle\|_\infty \cdot \dim V^{K_e}.$$

Since the function $\gamma \mapsto \langle K_0 \rangle \gamma \langle K_0 \rangle$ is locally constant, the *local* growth of the right hand side is equivalent to that of $\dim V^{K_e}$. This depends on γ via e . For x sufficiently close to the set of singular elements (namely, for $\mathrm{sd}(\gamma) > e + d(\gamma)$) we may take $e = \mathrm{sd}(\gamma)$ by Theorem 7.2.

Unfortunately, a direct computation for GL_n shows that

$$\sum_{e=0}^{\infty} \dim V^{K_e} \cdot \mu\{g \in K_0 : \mathrm{sd}(g) = e\}$$

diverges, already for GL_2 . Hence the estimate (58) does not imply the local integrability of tr_π . The authors have not been able to detect the additional cancellation in our trace formula that makes the character locally integrable.

Instead, we estimate of the growth of $\dim V^{K_e}$. For convenience, we assume that $x = o$ is the origin of the apartment A_S and that $e \in \mathbb{Z}_{\geq 0}$.

Theorem 5.6 assigns to every convex subcomplex Σ of $\mathcal{B}(G, \mathbb{F})$ a subspace of V , namely the image $\sum_{x \in \Sigma^\circ} V^{U_x^{(e)}}$ of $\partial_0: C_0(\Sigma, V) \rightarrow V$. This space admits an important alternative description if Σ is finite.

Theorem 8.1 ([11, Theorem 2.12]). *The elements*

$$u_\Sigma^{(e)} := \sum_{\sigma \in \Sigma} (-1)^{\dim \sigma} \langle U_\sigma^{(e)} \rangle \in \mathcal{H}(G, \mathbb{Z}[1/p])$$

are *idempotent* and

$$\begin{aligned} u_\Sigma^{(e)} \mathcal{H}(G, \mathbb{Z}[1/p]) &= \sum_{x \in \Sigma^\circ} \langle U_x^{(e)} \rangle \mathcal{H}(G, \mathbb{Z}[1/p]), \\ (1 - u_\Sigma^{(e)}) \mathcal{H}(G, \mathbb{Z}[1/p]) &= \bigcap_{x \in \Sigma^\circ} (1 - \langle U_x^{(e)} \rangle) \mathcal{H}(G, \mathbb{Z}[1/p]). \end{aligned}$$

In particular,

$$\mathrm{im}(\partial_0: C_0(\Sigma, V) \rightarrow V) = \sum_{x \in \Sigma^\circ} V^{U_x^{(e)}} = u_\Sigma^{(e)} V.$$

It is shown in [11] that there is a convex subcomplex Σ_0 such that $\langle U_o^{(r)} \rangle u_\Sigma^{(e)} = \langle U_o^{(r)} \rangle u_{\Sigma_0}^{(e)}$ for all convex subcomplexes Σ with $\Sigma \supseteq \Sigma_0$. The following lemma describes Σ_0 explicitly. To state it, we need some notation. For $\alpha \in \Phi$ we define

$$\begin{aligned} A_{S,r}^{\alpha+} &:= \{x \in A_S : \langle x, \alpha \rangle > r\}, \\ A_{S,r}^{\alpha 0} &:= \{x \in A_S : \langle x, \alpha \rangle \in [-r, r]\}, \\ A_{S,r}^{\alpha-} &:= \{x \in A_S : \langle x, \alpha \rangle < -r\}, \end{aligned}$$

and for any map $\epsilon: \Phi \rightarrow \{+, 0, -\}$ we write

$$A_{S,r}^\epsilon := \bigcap_{\alpha \in \Phi} A_{S,r}^{\alpha, \epsilon(\alpha)}.$$

Most of the sets $A_{S,r}^\epsilon$ are empty, some are compact, and the others are unbounded. The non-empty $A_{S,r}^\epsilon$ partition A_S . Let $A_{S,r}^b$ be the union of the bounded $A_{S,r}^\epsilon$; this is a polysimplicial subcomplex of A_S which is star-shaped around o . The subcomplex $B_r := P_o \cdot A_{S,r}^b$ of $\mathcal{B}(\mathcal{G}, \mathbb{F})$ is obviously stable under the action of all the groups $U_o^{(s)}$ for $s \in \mathbb{R}_{\geq 0}$. We may think of B_r as a combinatorial approximation to a ball of radius r around o .

Lemma 8.2. *Let $r \in \mathbb{Z}_{\geq e}$ and let $\Sigma \subseteq \mathcal{B}(\mathcal{G}, \mathbb{F})$ be any finite convex subcomplex that contains B_{r-e} . Then*

$$\langle U_o^{(r)} \rangle u_\Sigma^{(e)} = \langle U_o^{(r)} \rangle u_{B_{r-e}}^{(e)} = \sum_{\sigma \in B_{r-e}} (-1)^{\deg \sigma} \langle U_o^{(r)} \rangle \langle U_\sigma^{(e)} \rangle.$$

Proof. Fix $\epsilon: \Phi \rightarrow \{+, 0, -\}$ such that $A_{S,r-e}^\epsilon$ is unbounded. First we establish $\langle U_o^{(r)} \rangle \langle U_F^{(e)} \rangle = \langle U_o^{(r)} \rangle \langle U_{F'}^{(e)} \rangle$ for certain facets $F, F' \subseteq A_{S,r-e}^\epsilon$. The coroots $\alpha^\vee \in \Phi^\vee$ with $\epsilon(\alpha) = 0$ span a proper subspace $A_{S,\perp}^\epsilon \subsetneq A_S$. We may pick a non-zero vector $\delta^\epsilon \in A_S$ such that

- (1) δ^ϵ is orthogonal to $A_{S,\perp}^\epsilon$,
- (2) $A_{S,r-e}^\epsilon + \mathbb{R}_{\geq 0} \delta^\epsilon \subseteq A_{S,r-e}^\epsilon$,
- (3) δ^ϵ lies in the span of an irreducible root subsystem Ψ^\vee of Φ^\vee (here we decompose Φ^\vee as a direct sum of irreducible root systems).

For every facet $F \subseteq A_{S,r-e}^\epsilon$ let $M(F) \subseteq A_{S,r-e}^\epsilon$ be the unique facet such that for all $x \in F$ there exists $\lambda > 0$ with $x + \lambda \cdot \delta^\epsilon \subseteq M(F)$. We claim that

$$(59) \quad \langle U_o^{(r)} \rangle \langle U_F^{(e)} \rangle = \langle U_o^{(r)} \rangle \langle U_{M(F)}^{(e)} \rangle \quad \text{for } F \subseteq A_{S,r-e}^\epsilon.$$

In view of the unique decomposition property (Proposition 5.2) this is equivalent to

$$(U_o^{(r)} \cup U_F^{(e)}) \cap U_\alpha = (U_o^{(r)} \cup U_{M(F)}^{(e)}) \cap U_\alpha \quad \text{for all } \alpha \in \Phi^{\text{red}}.$$

By definition, $U_o^{(r)} \cap U_\alpha = U_{\alpha,r+}$ and $U_F^{(e)} \cap U_\alpha = U_{\alpha,-\alpha(x)+e+}$ for $x \in F$. If $\epsilon(\alpha) = -$, then $-\alpha + e > r$ on $F \cup M(F)$, so that

$$U_{\alpha,r+} \supseteq U_\alpha \cap (U_o^{(r)} \cup U_{M(F)}^{(e)}).$$

If $\epsilon(\alpha) \neq -$, then $\sup_{x \in F} -\alpha(x) \leq \sup_{x \in M(F)} -\alpha(x)$, which combined with $U_F^{(e)} \subseteq U_{M(F)}^{(e)}$ yields $U_F^{(e)} \cap U_\alpha = U_{M(F)}^{(e)} \cap U_\alpha$. This finishes the proof of (59).

Now we use (59) to establish some cancellation. Every facet F in A_S can be written uniquely as $F = F_\Psi \times F_\perp$, where F_Ψ and F_\perp are facets in $\mathbb{R}\Psi^\vee$ and $\Psi^\perp \subseteq A_S$, respectively. Consider a facet $F \subseteq A_{S,r-e}^\epsilon$ such that $M^{-1}(F)$ is not empty. Then

$M(F) = F$, and $M^{-1}(F)$ consists of facets of \overline{F} . Property (3) above shows that $F'_\perp = F_\perp$ for any $F' \in M^{-1}(F)$. Hence

$$\bigcup_{F' \in M^{-1}(F)} F' = \tau \times F_\perp,$$

where $\tau \subseteq \mathbb{R}\Psi^\vee$ consists of the facets of $\overline{F_\Psi}$ that contain points of the form $x + \lambda\delta^\epsilon$ with $x \in F$ and $\lambda \geq 0$. In particular, τ is diffeomorphic to

$$(-1, 1]\delta^\epsilon + \{x \in F : \langle x, \delta^\epsilon \rangle = c\}$$

for some $c \in \mathbb{R}$, so that the Euler characteristic of τ is zero. Therefore,

$$(60) \quad \sum_{F' \in M^{-1}(F)} (-1)^{\dim F'} = \sum_{F' \in M^{-1}(F)} (-1)^{\dim F'_\Psi} (-1)^{\dim F_\perp} = \sum_{\tau' \text{ facet in } \tau} (-1)^{\dim \tau'} (-1)^{\dim F_\perp} = 0,$$

which together with (59) yields

$$(61) \quad \sum_{F' \in M^{-1}(F)} (-1)^{\dim F'} \langle U_o^{(r)} \rangle \langle U_{F'}^{(e)} \rangle = 0 \in \mathcal{H}(G, \mathbb{Z}[1/p]).$$

Suppose that A_S is any apartment of $\mathcal{B}(\mathcal{G}, \mathbb{F})$ that contains o and at least one facet $F' \in M^{-1}(F)$. As δ^ϵ points away from o , the apartment A_S contains points of F , so that $\overline{F} \subseteq A_S$. This enables us to extend the map M to all facets of $\mathcal{B}(\mathcal{G}, \mathbb{F})$. Recall that any Weyl chamber $A_S^+ \subseteq A_S$ is a fundamental domain for the action of P_o on $\mathcal{B}(\mathcal{G}, \mathbb{F})$. On A_S^+ we define M according to the above recipe and by $M(F) := F$ if $F \subseteq A_{S,r-e}^b \cap A_S^+$. The properties (1)–(3) of δ^ϵ ensure that $M(F)$ and F have the same isotropy group in P_o , so we can extend M P_o -equivariantly to $\mathcal{B}(\mathcal{G}, \mathbb{F})$.

Since Σ contains o and is a convex subcomplex of $\mathcal{B}(\mathcal{G}, \mathbb{F})$, its collection of facets is stable under M . By definition

$$\begin{aligned} \langle U_o^{(r)} \rangle u_\Sigma^{(e)} &= \langle U_o^{(r)} \rangle \sum_{\sigma \in \Sigma} (-1)^{\deg \sigma} \langle U_\sigma^{(e)} \rangle \\ &= \langle U_o^{(r)} \rangle \sum_{F \text{ facet of } \Sigma} \sum_{F' \in M^{-1}(F)} (-1)^{\dim F'} \langle U_{F'}^{(e)} \rangle. \end{aligned}$$

Now (61) (which only holds for facets of unbounded $A_{S,r-e}^b$) shows that the facets of $\Sigma \setminus B_{r-e}$ do not contribute to this sum. As M is the identity on facets of B_{r-e} , we remain with $\langle U_o^{(r)} \rangle u_\Sigma^{(e)} = \langle U_o^{(r)} \rangle u_{B_{r-e}}^{(e)}$. \square

Remark 8.3. Lemma 8.2 provides a direct proof of the special case of [11, Proposition 3.6] where the consistent system of idempotents is $\langle U_x^{(e)} \rangle$; this proof does not use the fact that the Hecke algebra is Noetherean.

We turn to the space of invariants $V^{U_o^{(r)}}$. Since it has finite dimension, it is contained in the range of $u_\Sigma^{(e)}$ for some finite convex subcomplex $\Sigma \subseteq \mathcal{B}(\mathcal{G}, \mathbb{F})$. We may as well assume that Σ contains B_{r-e} , so that Lemma 8.2 yields

$$V^{U_o^{(r)}} = \langle U_o^{(r)} \rangle u_\Sigma^{(e)} V = \left(\sum_{\sigma \in B_{r-e}} (-1)^{\deg \sigma} \langle U_o^{(r)} \rangle \langle U_\sigma^{(e)} \rangle \right) V.$$

The right hand side is contained in $\sum_{x \in B_{r-e}^\circ} \langle U_o^{(r)} \rangle \langle U_x^{(e)} \rangle V$ by Theorem 5.5.(e). It is the space of $U_o^{(r)}$ -invariants in $\sum_{x \in B_{r-e}^\circ} \langle U_x^{(e)} \rangle V$ because $\sum_{x \in B_{r-e}^\circ} \langle U_x^{(e)} \rangle V$ is

P_o -invariant. Let $P_o \supseteq \langle U_o^{(r)} \rangle$ act on $\bigoplus_{x \in B_{r-e}^{\circ}} \langle U_x^{(e)} \rangle V$ by $g \cdot (x, v) = (g \cdot x, \pi(g)v)$. Then $\sum_{x \in B_{r-e}^{\circ}} \langle U_o^{(r)} \rangle \langle U_x^{(e)} \rangle V$ is a quotient of $\bigoplus_{x \in B_{r-e}^{\circ}} \langle U_x^{(e)} \rangle V$. The addition map

$$\left(\bigoplus_{x \in B_{r-e}^{\circ}} \langle U_x^{(e)} \rangle V \right)^{U_o^{(r)}} \rightarrow \left(\sum_{x \in B_{r-e}^{\circ}} \langle U_x^{(e)} \rangle V \right)^{U_o^{(r)}}$$

is surjective because $U_o^{(r)}$ is compact and we are working in good characteristic. Since there are only finitely many G -orbits of vertices in $\mathcal{B}(\mathcal{G}, \mathbb{F})$,

$$(62) \quad m_V := \max_{x \in \mathcal{B}(\mathcal{G}, \mathbb{F})} \dim V^{U_x^{(e)}}$$

exists. The dimension of $(\bigoplus_{x \in B_{r-e}^{\circ}} \langle U_x^{(e)} \rangle V)^{U_o^{(r)}}$ is at most $m_V |B_{r-e}^{\circ}/U_o^{(r)}|$.

It remains to estimate the number of $U_o^{(r)}$ -orbits of vertices in B_{r-e} . For $\alpha \in \Phi$ let d_α be the dimension of $\text{Lie}_{\mathbb{F}}(\mathcal{U}_\alpha/\mathcal{U}_{2\alpha})$ and let d_0 be the dimension of $\text{Lie}_{\mathbb{F}}(\mathcal{Z}_{\mathcal{G}}(\mathcal{S}))$. Recall that $q = |\mathcal{O}/\mathfrak{P}|$ and that $n_\alpha^{-1}\mathbb{Z}$ is the set of jumps of the filtration of U_α .

Lemma 8.4. *The number of $U_o^{(r)}$ -orbits on B_{r-e}° is of order $O(r^{\dim A_S} Q^r)$, where*

$$Q := \exp \left(\log(q) \sum_{\alpha \in \Phi^{\text{red}}} \frac{d_\alpha n_\alpha}{2} + \frac{d_{2\alpha} n_{2\alpha}}{4} \right).$$

Proof. Recall from (16) and Proposition 3.2.(c) that

$$P_o = U_o N_o = U_o^+ U_o^- (P_o \cap N_G(S)),$$

for any positive root system Φ^+ of Φ . Hence every facet of $B_{r-e} = P_o \cdot A_{S, r-e}^b$ is of the form $u \cdot F$ with $u \in U_o^+ U_o^-$ and a facet F of A_S . Fix F and choose a positive root system Φ^+ such that $\alpha(F) \geq 0$ for all $\alpha \in \Phi^+$. Then $U_o^- \subseteq U_F^-$ fixes F pointwise, so that we only need $u \in U_o^+$. By Propositions 3.2.(b) and 5.2 the product maps

$$\prod_{\alpha \in \Phi^{\text{red}} \cap \Phi^+} U_{\alpha,0} \rightarrow U_o^+, \quad \prod_{\alpha \in \Phi^{\text{red}} \cap \Phi^+} (U_\alpha \cap U_o^{(r)}) \rightarrow U^+ \cap U_o^{(r)}$$

are diffeomorphisms. Together with the conventions (10) we get

$$(63) \quad [U_o^+ : U_o^+ \cap U_o^{(r)}] = \prod_{\alpha \in \Phi^{\text{red}} \cap \Phi^+} [U_{\alpha,0} : U_\alpha \cap U_o^{(r)}] \\ = \prod_{\alpha \in \Phi^{\text{red}} \cap \Phi^+} [U_{\alpha,0} U_{2\alpha,0} : U_{\alpha,r+} U_{2\alpha,2r+}] \\ = \prod_{\alpha \in \Phi^{\text{red}} \cap \Phi^+} [U_{\alpha,0}/U_{2\alpha,0} : U_{\alpha,r+}/U_{2\alpha,2r+}] \cdot [U_{2\alpha,0} : U_{2\alpha,2r+}].$$

Since we are dealing with unipotent pro- p -groups, these indices can be read off from the Lie algebras. For $\alpha \in \Phi$ and $s \in n_\alpha^{-1}\mathbb{Z}$, the construction from (7) and (10) shows that $U_{\alpha,s} \supsetneq U_{\alpha,s+}$ corresponds to multiplying a lattice in $\text{Lie}_{\mathbb{F}}(\mathcal{U}_\alpha)$ with the maximal ideal \mathfrak{P} of \mathcal{O} , see also [21, 3.5.4]. Hence

$$[U_{\alpha,s}/U_{2\alpha,2s} : U_{\alpha,s+}/U_{2\alpha,2s+}] = q^{d_\alpha}, \\ [U_{\alpha,0}/U_{2\alpha,0} : U_{\alpha,r+}/U_{2\alpha,2r+}] = q^{d_\alpha \lceil n_\alpha r+ \rceil},$$

where $\lceil y+ \rceil$ denotes the smallest integer larger than $y+ \in \tilde{\mathbb{R}}$. Similarly

$$[U_{2\alpha,0} : U_{2\alpha,r+}] = q^{d_{2\alpha} \lceil n_{2\alpha} r/2+ \rceil},$$

from which we conclude that

$$(64) \quad [U_o^+ : U^+ \cap U_o^{(r)}] = \prod_{\alpha \in \Phi^{\text{red}} \cap \Phi^+} q^{d_\alpha \lceil n_\alpha r + \rceil} q^{d_{2\alpha} \lceil n_{2\alpha} r / 2 + \rceil} \\ \leq \prod_{\alpha \in \Phi^{\text{red}}} q^{d_\alpha (n_\alpha r + 1) / 2} q^{d_{2\alpha} (n_{2\alpha} r + 2) / 4}.$$

This number is an upper bound for the number of $U_o^{(r)}$ -orbits in $U_o \cdot F$. Since it does not depend on F , we only need to multiply it with the number of facets of $A_{S, r-e}^b$. While this number is not easily expressible in a formula, it clearly grows like $r^{\dim A_S}$. \square

Theorem 8.5. *Let (π, V) be an admissible G -representation of level $e \in \mathbb{Z}_{\geq 0}$ in good characteristic. Let $r \in \mathbb{R}_{\geq e}$ and define Q and m_V as in Lemma 8.4 and (62). Then*

$$\dim V_o^{U_o^{(r)}} = O(m_V r^{\dim A_S} Q^r) \\ \mu(U_o^{(r)}) \dim V_o^{U_o^{(r)}} = O(m_V r^{\dim A_S} q^{-rd_0} Q^{-r})$$

with constants independent of V and r .

Proof. The first estimate follows from Lemma 8.4 and the arguments above. Proposition 5.2 yields

$$[U_o^{(s)} : U_o^{(r+s)}] = [H_{s+} : H_{r+s+}] \prod_{\alpha \in \Phi^{\text{red}}} [U_{\alpha, s+} U_{2\alpha, s+} : U_{\alpha, r+s+} U_{2\alpha, r+s+}]$$

for all $s \in \mathbb{Z}_{\geq 0}$. A calculation like the one in (63) and (64) shows that this index is at least

$$q^{rd_0} \prod_{\alpha \in \Phi^{\text{red}}} q^{rn_\alpha d_\alpha} q^{rn_{2\alpha} d_{2\alpha} / 2}.$$

(We cannot be exact because we do not know at which points the filtration of H jumps.) This yields the second estimate. \square

These estimates are sharp in some examples: (57) shows that (a) and (c) cannot be improved for GL_n . Here all n_α and d_α are 1, Φ is reduced, and there are $n(n-1)/2$ positive roots, so that $Q = q^{n(n-1)/2}$.

9. CONCLUSION

Let G be a reductive p -adic group and let (ρ, V) be an admissible representation of G on a vector space V of characteristic not equal to p . We have seen that the character of (ρ, V) is a locally constant function on the set of regular semi-simple elements, and we have described explicit open subsets on which it is constant. Furthermore, we have estimated the growth of the dimensions of the fixed-point subspaces $V_x^{U_x^{(e)}}$ for $e \rightarrow \infty$. Both results are based on the main result of [11] about the acyclicity of certain coefficient systems on the affine Bruhat–Tits building.

It is still unclear whether Harish-Chandra’s theorem about the local integrability of the character function for complex representations can be established using these resolutions. This may depend on a better understanding of the character formulas. While the resolution in [11] does provide an explicit formula for the character, more work is required to understand and simplify this formula.

REFERENCES

- [1] Jeffrey D. Adler and Jonathan Korman, *The local character expansion near a tame, semisimple element*, Amer. J. Math. **129** (2007), no. 2, 381–403. MR **2306039**
- [2] Armand Borel and Jacques Tits, *Groupes réductifs*, Inst. Hautes Études Sci. Publ. Math. **27** (1965), 55–150 (French). MR **0207712**
- [3] François Bruhat and Jacques Tits, *Groupes réductifs sur un corps local I. Données radicielles valuées*, Inst. Hautes Études Sci. Publ. Math. **41** (1972), 5–251 (French). MR **0327923**
- [4] ———, *Groupes réductifs sur un corps local. II. Schémas en groupes. Existence d'une donnée radicielle valuée*, Inst. Hautes Études Sci. Publ. Math. **60** (1984), 197–376 (French). MR **756316**
- [5] William A. Casselman, *Characters and Jacquet modules*, Math. Ann. **230** (1977), no. 2, 101–105. MR **0492083**
- [6] Pierre Deligne, *Le support du caractère d'une représentation supercuspidale*, C. R. Acad. Sci. Paris Sér. A-B **283** (1976), no. 4, Aii, A155–A157 (French, with English summary). MR **0425033**
- [7] Harish-Chandra, *Collected papers. Vol. IV*, Springer-Verlag, New York, 1984. 1970–1983; Edited by V. S. Varadarajan. MR **726026**
- [8] ———, *Admissible invariant distributions on reductive p -adic groups*, University Lecture Series, vol. 16, American Mathematical Society, Providence, RI, 1999. Preface and notes by Stephen DeBacker and Paul J. Sally, Jr. MR **1702257**
- [9] Jonathan Korman, *A character formula for compact elements (the rank one case)* (2004), eprint. arXiv: math.RT/0409292.
- [10] Serge Lang, *Algebra*, 3rd ed., Graduate Texts in Mathematics, vol. 211, Springer-Verlag, New York, 2002. MR **1878556**
- [11] Ralf Meyer and Maarten Solleveld, *Resolutions for representations of reductive p -adic groups via their buildings* (2009), eprint. arXiv: 0902.4856.
- [12] Allen Moy and Gopal Prasad, *Unrefined minimal K -types for p -adic groups*, Invent. Math. **116** (1994), no. 1–3, 393–408. MR **1253198**
- [13] ———, *Jacquet functors and unrefined minimal K -types*, Comment. Math. Helv. **71** (1996), no. 1, 98–121. MR **1371680**
- [14] Takashi Ono, *Arithmetic of algebraic tori*, Ann. of Math. (2) **74** (1961), 101–139. MR **0124326**
- [15] Gopal Prasad, *Elementary proof of a theorem of Bruhat-Tits-Rousseau and of a theorem of Tits*, Bull. Soc. Math. France **110** (1982), no. 2, 197–202 (English, with French summary). MR **667750**
- [16] ———, *Galois-fixed points in the Bruhat–Tits building of a reductive group*, Bull. Soc. Math. France **129** (2001), no. 2, 169–174 (English, with English and French summaries). MR **1871292**
- [17] G. Rousseau, *Immeubles des groupes réductifs sur les corps locaux*, Thesis, Université de Paris-Sud, 1977 (French).
- [18] Peter Schneider and Ulrich Stuhler, *Representation theory and sheaves on the Bruhat–Tits building*, Inst. Hautes Études Sci. Publ. Math. **85** (1997), 97–191. MR **1471867**
- [19] Jean-Pierre Serre, *Local fields*, Graduate Texts in Mathematics, vol. 67, Springer-Verlag, New York, 1979. Translated from the French by Marvin Jay Greenberg. MR **554237**
- [20] T. A. Springer, *Linear algebraic groups*, 2nd ed., Progress in Mathematics, vol. 9, Birkhäuser Boston Inc., Boston, MA, 1998. MR **1642713**
- [21] J. Tits, *Reductive groups over local fields*, Automorphic forms, representations and L -functions (Proc. Sympos. Pure Math., Oregon State Univ., Corvallis, Ore., 1977), Part 1, Proc. Sympos. Pure Math., XXXIII, Amer. Math. Soc., Providence, R.I., 1979, pp. 29–69. MR **546588**
- [22] Marie-France Vignéras, *Représentations l -modulaires d'un groupe réductif p -adique avec $l \neq p$* , Progress in Mathematics, vol. 137, Birkhäuser Boston Inc., Boston, MA, 1996 (French, with English summary). MR **1395151**
- [23] ———, *Cohomology of sheaves on the building and R -representations*, Invent. Math. **127** (1997), no. 2, 349–373. MR **1427623**
- [24] Marie-France Vignéras and Jean-Loup Waldspurger, *Premiers réguliers de l'analyse harmonique mod l d'un groupe réductif p -adique*, J. Reine Angew. Math. **535** (2001), 165–205. MR **1837098**

E-mail address: rameyer@uni-math.gwdg.de

E-mail address: maarten@uni-math.gwdg.de

MATHEMATISCHES INSTITUT AND COURANT CENTRE “HIGHER ORDER STRUCTURES”, GEORG-AUGUST UNIVERSITÄT GÖTTINGEN, BUNSENSTRASSE 3–5, 37073 GÖTTINGEN, GERMANY